

Increased leukotriene B₄ release from ileal pouch mucosa in ulcerative colitis compared with familial adenomatous polyposis

D J Gertner, D S Rampton, M V Madden, I C Talbot, R J Nicholls, J E Lennard-Jones

Abstract

Pouchitis may complicate the construction of an ileal pouch after colectomy for ulcerative colitis (UC) but not familial adenomatous polyposis (FAP). To examine whether differences in eicosanoid metabolism might explain why pouchitis is largely confined to UC patients, this study compared arachidonic acid stimulated release of immunoreactive leukotriene B₄ (LTB₄) and prostaglandin E₂ (PGE₂) from macroscopically uninfamed pouch mucosal biopsy specimens incubated in vitro from patients with UC and FAP. The study also compared eicosanoid release from inflamed and uninfamed pouches in patients with UC. In uninfamed pouches, median LTB₄ release was nearly twice as high in UC as in FAP (p=0.001), but there was no significant difference in PGE₂ production. In UC, stimulated eicosanoid release from uninfamed functioning pouch mucosa was not significantly different from that from either ileostomy or defunctioned pouch mucosa. LTB₄ and PGE₂ release were significantly greater from inflamed than uninfamed pouch mucosa in UC (p=0.001 and 0.01, respectively). Leukotriene synthesis inhibition or receptor antagonism, or both merit therapeutic evaluation in pouchitis. Increased release of LTB₄ from endoscopically normal pouch mucosa suggests increased 5-lipoxygenase activity in patients with UC and could contribute to their predisposition to pouchitis.

(Gut 1994; 35: 1429-1432)

Pouchitis is a syndrome produced by acute inflammation of the mucosa of an ileal reservoir. It is the commonest longterm complication of a restorative proctocolectomy for ulcerative colitis (UC), but is very rare after the same operation for familial adenomatous polyposis (FAP).^{1,2} The cause is unknown.

Mucosal overproduction of eicosanoids seems to play a part in the pathogenesis of UC.^{3,4} Increased synthesis of the lipoxygenase products of arachidonic acid metabolism, in particular, may contribute to the inflammatory process through their effects on migration and degranulation of neutrophils and on vascular permeability.³⁻⁶ Such changes, once initiated, tend to be self perpetuating through the further recruitment of inflammatory cells into the mucosa, and their activation.

As patients with UC have a particular predilection for developing pouch mucosal inflammation and FAP patients do not, we tested the hypothesis that this might result from differences in the metabolism of eicosanoids. We used a short term in vitro incubation technique to measure the release of leukotriene B₄ (LTB₄) and prostaglandin E₂ (PGE₂) from biopsy specimens of uninfamed pouch mucosa, comparing patients with UC with those with FAP.^{6,7} To see if changes in eicosanoid release in pouchitis mimic those in rectal mucosa in UC,³⁻⁷ we also assessed eicosanoid release from inflamed pouch mucosa.

Methods

The study was approved by the ethical committee of the City and Hackney Health Authority.

PATIENTS

We studied 28 patients with UC 4-91 months after restorative proctocolectomy, and eight patients with FAP 3-24 months after the operation.

Patients with uninfamed pouches - none of these patients had symptoms, or signs on examination with a paediatric sigmoidoscope, of pouchitis. In the patients with UC, the pouch was in continuity with the faecal stream in 16, of whom six were taking loperamide, while in the remaining six patients it was defunctioned by a proximal loop ileostomy. In all the patients with FAP, the pouch was in continuity with the faecal stream, and two were taking loperamide.

Patients with inflamed pouches - six of the UC patients had one or more episodes (16 episodes in all) of pouchitis diagnosed clinically, endoscopically, and histologically during the study.¹ In all these patients, the pouch was in continuity with the faecal stream. During five episodes, the patient was taking metronidazole, and during two loperamide.

BIOPSY SITES

In each patient studied, three biopsy specimens were taken from adjacent sites in the pouch or on the surface of the loop ileostomy. Two specimens were used to measure the release of LTB₄ and PGE₂, and one was examined histologically.

St Mark's Hospital,
City Road, London
D J Gertner
M V Madden
I C Talbot
R J Nicholls
J E Lennard-Jones

Gastrointestinal
Science Research Unit,
London Hospital
Medical College,
London
D S Rampton

Correspondence to:
Dr D J Gertner, Basildon
Hospital, Nether Mayne,
Basildon, Essex SS16 5NL.

Accepted for publication
9 February 1994

IN VITRO INCUBATIONS

A modification of Peskar's technique was used.^{6,7} Two specimens (20–40 mg) were separately washed in Tyrode's solution, blot dried, and weighed. Within five minutes, each was placed in a 5 ml glass container for preincubation for 20 minutes in 1 ml continuously gassed (95% O₂/5% carbon dioxide) Tyrode's solution at 37°C. The supernatant was then replaced with 1 ml Tyrode's solution containing arachidonic acid (sodium salt, Sigma, 10 µg/ml) and the specimen incubated for a further 20 minutes. The supernatant was then removed and frozen at -20°C for subsequent assay of its eicosanoid content.

Viability of the incubated specimens was confirmed by their continuing and unaltered release of LTB₄ and PGE₂ with added arachidonic acid during successive 20 minute periods over a total of four hours. In otherwise identical experiments using colorectal mucosal specimens from patients with UC, the coefficient of variation of duplicate adjacent biopsy specimens incubated with arachidonic acid for 20 minutes was 6% (n=6) for LTB₄ release and 8% (n=6) for PGE₂.⁷

ASSAYS

All assays were performed in duplicate on unextracted supernatant within three months of the incubation experiment.

LTB₄ concentration was measured by radioimmunoassay (Amersham International kit no TRK 940). Sensitivity was 0.02 ng/ml, cross reactivity with arachidonic acid <0.05% and with other eicosanoids <0.05%, and the interassay variation 5%. Recovery of LTB₄ added to the supernatant was 79% (72–84) (median (interquartile range)) (n=6). Serial dilution of selected samples showed parallelism with standard solutions. Four random samples were validated by high performance liquid chromatography after extraction and purification of supernatant⁸: the coefficient of variation of the duplicate high performance liquid chromatography and radioimmunoassay measurements was 4%.

PGE₂ was also measured by radioimmunoassay using standard solutions of PGE₂ (Sigma), PGE₂ antisera (gift of Dr J A Salmon) and PGE₂ tracer (Amersham International).⁹ The sensitivity of the method was 0.1 ng/ml, and cross reactivity with 6-keto-PGF₁α, arachidonic acid, and other eicosanoids was 4.8%, <0.1%, and <1%, respectively. The interassay variation was 6%. Recovery of PGE₂ was 71% (63–80) (n=6), and parallelism of serial dilutions of supernatant with standard solutions was confirmed. Six random samples were validated by rat stomach strip bioassay¹⁰; the coefficient of variation of the bioassay and radioimmunoassay results was 3%.

Preliminary studies showed that the presence of arachidonic acid in a concentration of 10 µg/ml did not affect the LTB₄ or PGE₂ radioimmunoassays.

The values shown for LTB₄ and PGE₂ release for each patient are the means of the

concentrations measured in the supernatants of the two biopsy specimens incubated for each subject.

HISTOLOGICAL ASSESSMENT

The third specimen was used to provide a formalin fixed section stained with haematoxylin and eosin. A pathologist unaware of the group to which the patient belonged provided a semi-quantitative light microscopic score, from 0–3, of the extent of polymorphonuclear leucocyte infiltrate.¹¹

STATISTICS

Results are shown as median (interquartile range). Differences among groups were assessed by the Kruskal-Wallis test, and subsequently between groups using Wilcoxon's sum of ranks test (two tailed p values), as appropriate. Correlations were sought using Spearman's rank correlation test.

Results

UNINFLAMED POUCHES IN CONTINUITY WITH THE FAECAL STREAM

The pouch mucosa of patients with UC released significantly more LTB₄ (57 ng/g wet wt/20 min (44–89)) than did the pouch mucosa of patients with FAP (30 (27–35), p=0.001) (Fig 1). There was a trend towards increased release of PGE₂ in patients with UC (263 ng/g wet wt/20 min (169–446) compared with FAP (158 (123–287), but this did not attain statistical significance (Fig 2).

In UC pouches, eicosanoid release did not correlate with either the score for polymorphonuclear leucocyte infiltration (for LTB₄, r=-0.16, p=0.59; for PGE₂, r=0.08, p=0.79, or with the time interval since restorative proctocolectomy (LTB₄, r=-0.37, p=0.17; PGE₂, r=-0.28, p=0.13). There was also no difference in mucosal infiltration of neutrophils in the pouches of patients with




Figure 1: LTB₄ release by mucosal biopsy specimens from endoscopically uninflamed pouches (functioning and defunctioned) and ileostomies in patients with UC and FAP. Horizontal bars denote median values.




Figure 2: PGE₂ release by mucosal biopsy specimens from endoscopically uninflamed pouches (functioning and defunctioned) and ileostomies in patients with UC and FAP. Horizontal bars denote median values.

UC and FAP (scores 0 (0–1), and 0 (0–1), respectively).

UNINFLAMED DEFUNCTIONED POUCHES AND ILEOSTOMIES

There were no significant differences in release of LTB₄ from defunctioned pouch, ileostomy, and functioning pouch mucosa in patients with UC (Fig 1). There were also no differences in the release of PGE₂ in these categories of specimens (Fig 2). The median release of LTB₄ from defunctioned pouches ($p=0.001$), but not ileostomies, in patients with UC was significantly greater than that from pouches of patients with FAP. Although the median release of PGE₂ from defunctioned pouches and from ileostomies in patients with UC was greater than that from the pouches of patients with FAP, the differences were not statistically significant (Figs 1 and 2).

INFLAMED POUCHES IN CONTINUITY WITH THE FAECAL STREAM

Inflamed pouch mucosa released significantly more LTB₄ ($p=0.001$) and PGE₂ ($p=0.01$) than uninflamed pouch mucosa (Table). Median release of LTB₄, but not PGE₂, from inflamed (and uninflamed) ileal pouch mucosa was also significantly higher than we have previously reported from rectal mucosa in patients with active (and inactive) UC⁷ (Table).

Discussion

It is well established that colorectal mucosa in patients with active UC produces more LTB₄

and PGE₂ than does uninflamed mucosa.^{3–7} We have shown here that ileal pouch mucosa when inflamed also releases more of these eicosanoids, confirming previous preliminary data.¹² As in UC,^{3–7} increased synthesis of leukotrienes probably contributes to pouch mucosal inflammation largely by increasing the recruitment and activation of neutrophils. Excess prostaglandin production may worsen diarrhoea by inducing mucosal secretion of water and electrolytes; conversely, increased prostaglandin synthesis may have a mucoprotective role by increasing mucus release, changing mucosal blood flow, and suppressing immune and inflammatory cell function.^{3,4,13} While, in patients with UC, inhibition of prostaglandin synthesis has been shown to be therapeutically ineffective, and indeed possibly harmful,^{3,4} limited data suggest that inhibition of mucosal leukotriene synthesis with A-64077 (Zileuton),¹⁴ may have some benefit, at least in patients not already taking aminosalicylates.¹⁵ By analogy, a clinical trial of Zileuton, or, alternatively, an LTB₄ receptor antagonist, would seem appropriate in patients with pouchitis.

It is not known why pouchitis occurs in the ileal mucosa of patients with UC but not those with FAP.¹² In this study, we report that patients with UC release more LTB₄ from the mucosa of uninflamed functioning pouches than do patients with FAP.

Polymorphonuclear leucocytes are an important source of LTB₄. In patients with uninflamed pouches, however, there was no difference in the intensity of mucosal infiltration by neutrophils in UC and FAP; furthermore there was no correlation between neutrophil infiltration and LTB₄ release in uninflamed UC pouches. The ability of apparently normal pouch mucosa in patients with UC to release increased amounts of LTB₄ shows that the 5-lipoxygenase enzyme system has been activated. Studies of polymorphonuclear leucocytes in the peripheral blood of patients with UC and Crohn's disease have shown an increased release of LTB₄ compared with healthy volunteers.¹⁶ Our findings may thus reflect activation of 5-lipoxygenase within mucosal neutrophils.

The lack of a significant correlation between the time since restorative proctocolectomy and the mucosal release of LTB₄ suggests that operative trauma, residual 'backwash' ileitis, and drugs used during the perioperative period are unlikely to explain the increase in LTB₄ release by uninflamed pouch mucosa in patients with UC.

Our data are consistent with a predisposition to acute inflammation in the mucosa of

Eicosanoid release in uninflamed and inflamed pouch mucosa and rectal mucosa in patients with UC. Data for rectal mucosa are taken from Gertner et al.⁷ Results are shown as median (interquartile range) (n)

	LTB ₄ release (ng/g wet wt/20 min) Pouch	Rectum	PGE ₂ release (ng/g wet wt/20 min) Pouch	Rectum
Uninflamed	57 (44–89) (16)	30 (24–40) (27)*	263 (169–446) (16)	284 (205–342) (27)
Inflamed	87 (70–94) (16)*	58 (45–73) (16)†‡	668 (583–769) (6)†	492 (396–598) (16)§

* $p=0.001$ from uninflamed pouch mucosa, † $=0.01$ from inflamed pouch mucosa, ‡ $=0.01$ from uninflamed pouch mucosa, § $=<0.05$ from uninflamed rectal mucosa.

functioning ileal pouches in patients with UC. They do not, however, show whether this predisposition requires exposure to the faecal stream, as there was no difference between the amounts of LTB₄ released from functioning and defunctioned pouches in patients with UC. Nevertheless, increased LTB₄ synthesis in UC patients with uninfamed pouches suggests a possible role for new drugs inhibiting the synthesis of leukotrienes,^{14 15} or antagonising their receptors, for the prevention of pouchitis.

The method used in this study exposed the mucosal biopsy specimens to an excess of arachidonic acid, the substrate for leukotriene production. It is possible that in vivo, without such an excess of substrate the activated 5-lipoxygenase enzyme system would not synthesise excess leukotrienes and would not therefore cause mucosal inflammation. Our failure to detect, in the mucosa of uninfamed pouches, a difference in the infiltrate of polymorphonuclear leucocytes between patients with UC and FAP suggests that excess LTB₄ was not being synthesised in vivo, as it is a powerful chemotactic mediator. It is possible, however, that synthesis of excess LTB₄ (and other leukotrienes), and the subsequent onset of pouchitis, could be triggered by any of a number of hormonal, inflammatory, and immunological factors, which have been shown to promote release of arachidonic acid.¹⁷ Conversely, reducing mucosal arachidonic acid by changing the diet, for example using fish oil,^{18 19} may merit evaluation in patients with recurrent pouchitis.

It has been suggested that increased mucosal eicosanoid production in active UC could be explained simply by greater availability of substrate in inflamed than control mucosa.²⁰ Our results, in which the pouch mucosa of patients with UC released more LTB₄ than the mucosa of FAP patients incubated in a similar excess of arachidonic acid, argue against this proposal, albeit less directly than in analogous studies using colonic mucosa.⁷

Release of LTB₄, but not PGE₂, from both inflamed and uninfamed pouch mucosa exceeded that from rectal mucosa in active and quiescent UC (Table). These comparisons raise the possibility of an inherent difference between LTB₄ release from small and large bowel, for example because of differences in the underlying cell populations, or lumenally derived stimuli, responsible for LTB₄ production in the ileum and colon. Previous work, however, in which operative specimens of uninfamed mucosa were stimulated in vitro by calcium ionophore (A23187), showed no difference in eicosanoid release between the ileum and colon.²¹ In our study, arachidonic acid was used as the stimulus for LTB₄ release; furthermore, the ileal tissue sampled had been previously transposed to an abnormal site (neorectum or stoma). These factors may explain the discrepancy between the present and previous results²¹: clearly, a study of eicosanoid release from normal ileal and

colorectal mucosa obtained colonoscopically is desirable for any further investigation of possible regional differences in leukotriene synthesis in the intestine.

The reason for increased 5-lipoxygenase activity in the uninfamed ileal mucosa of patients with function ileal pouches who have UC, which is thought to be primarily a disease of the colonic mucosa, is not clear. Explaining this may clarify not only the mechanism of pouchitis but also of ulcerative colitis itself.

We are grateful to the British Digestive Foundation for financial support (DJG).

Part of this work has been published in abstract form (*Gut* 1989; 30: A1481).

- Mortensen NJ, Madden MV. Pouchitis - acute inflammation in ileal pouches. In: Nicholls RJ, Bartolo DC, Mortensen NJ, eds. *Restorative proctocolectomy*. Oxford: Blackwell Scientific 1993: 119-31.
- de Silva HJ, Kettlewell MGW, Mortensen NJ, Jewell DP. Acute inflammation in ileal pouches (review). *Euro J Gastroenterol Hepatol* 1991; 3: 343-9.
- Rampton DS, Hawkey CJ. Prostaglandins and ulcerative colitis. *Gut* 1984; 25: 1399-1413.
- Lauritsen K, Laursen LS, Bukhave K, Rask-Madsen J. Inflammatory intermediaries in inflammatory bowel disease. *Int J Colorectal Dis* 1989; 4: 75-90.
- Sharon P, Stenson WF. Enhanced synthesis of leukotriene B₄ by colonic mucosa in inflammatory bowel disease. *Gastroenterology* 1984; 86: 453-60.
- Peskar BM, Dreyling KW, Peskar BA, May B, Goebell H. Enhanced formation of sulfido-leucotrienes in ulcerative colitis and Crohn's disease: inhibition by sulphasalazine and 5-aminosalicylic acid. *Agents Actions* 1986; 18: 381-3.
- Gertner DJ, Rampton DS, DeNucci G, Cynk E, Lennard-Jones JE. Eicosanoid release by rectal mucosa in vitro in ulcerative colitis: effects of conventional and potential new therapies. *Eur J Gastroenterol Hepatol* 1992; 4: 837-41.
- Galton SA, Piper PJ. The role of blood vessels in the bioconversion of leukotrienes in the pig. *Br J Pharmacol* 1987; 90: 617-22.
- Salmon JA. A radioimmunoassay for 6-ketoprostaglandin F₁alpha. *Prostaglandins* 1978; 15: 383-97.
- Vane JR. A sensitive method for the assay of 5-hydroxytryptamine. *Br J Pharmacol Chemotherap* 1957; 12: 344-9.
- Shepherd NA, Jass JR, Duval I, Moskowitz RL, Nicholls RJ. Restorative proctocolectomy with ileal reservoir: pathological and histochemical study of mucosal biopsy specimens. *J Clin Pathol* 1987; 40: 601-7.
- Belluzzi A, Gionchetti P, Campieri M, Boschi S, Tampieri M, Bertinelli E, et al. Eicosanoids, arachidonic acid and interleukin-1beta in pouchitis. *Gut* 1992; 33 (Suppl 2): S41.
- Knudson PJ, Dinarello CJ, Strom TB. Prostaglandins post-transcriptionally inhibit monocyte expression of interleukin-1 activity by increasing intracellular cyclic adenosine monophosphate. *J Immunol* 1986; 137: 3189-94.
- Laursen LS, Naesdal J, Lauritsen K, Rask-Madsen J. Selective 5-lipoxygenase inhibition in ulcerative colitis. *Lancet* 1990; 335: 683-5.
- Stenson WF, Lauritsen K, Laursen LS, Rask-Madsen J, Jacobsen O, Naesdal J, et al. A clinical trial of Zileuton, a specific inhibitor of 5-lipoxygenase, in ulcerative colitis. *Gastroenterology* 1991; 100: A253.
- Nielsen OH, Ahnfelt-Ronne, Elmgreen J. Abnormal metabolism of arachidonic acid in chronic inflammatory bowel disease: enhanced release of leukotriene B₄ from activated neutrophils. *Gut* 1987; 28: 181-5.
- Deby C. Metabolism of polyunsaturated fatty acids, precursors of eicosanoids. In: P B Curtis-Prior, ed. *Prostaglandins: biology and chemistry of prostaglandins and related eicosanoids*. Edinburgh: Churchill Livingstone, 1988: 11-36.
- Hillier K, Jewell R, Dorrell L, Smith CL. Incorporation of fatty acids from fish oil and olive oil into colonic mucosal lipids and effects upon eicosanoid synthesis in inflammatory bowel disease. *Gut* 1991; 32: 1151-5.
- Hawthorne AB, Daneshmend TK, Hawkey CJ, Belluzzi A, Everitt SJ, Holmes GKT, et al. Treatment of ulcerative colitis with fish oil supplementation: a prospective 12 month randomised controlled trial. *Gut* 1992; 33: 922-8.
- Nishida T, Miwa H, Shigematsu A, Yamamoto M, Iida M, Fujishima M. Increased arachidonic acid composition of phospholipids in colonic mucosa from patients with active ulcerative colitis. *Gut* 1987; 28: 1002-7.
- Dreyling KW, Hoppe U, Peskar BA, Morgenroth K, Kozuschek W, Peskar BM. Leukotriene synthesis by human gastrointestinal tissues. *Biochim Biophys Acta* 1986; 878: 184-93.