

similar.¹ Moreover, it has been known for several years that these bacterial protein toxins are also structurally similar to ricin and abrin.^{1,2} These plant toxins are, respectively, present in seeds of the castor bean plant (*Ricinus communis*) and the jequirity bean (*Abrus precatorius*), which are not closely related to each other.³ There is compelling evidence that the cytotoxic effect of all these toxins is mediated by an identical mechanism. The toxin molecules consist of an A subunit and one or more B subunits. After binding of the B subunit(s) to a cell surface receptor, the A subunit enters the cell, where it cleaves an adenine residue from ribosomal RNA molecules.^{1,2} The end result is that protein synthesis stops and the affected cells die.

This mechanism almost certainly explains the diarrhoea (with or without rectal bleeding) that follows ingestion of castor or jequirity beans,^{3,4} the colonic inflammation that occurs in *S dysenteriae* type 1 and *E coli* O157:H7 infections, and the endothelial cell damage that accompanies the haemolytic uraemic syndrome.⁵ The traditional equation of bacterial haemorrhagic colitis with bacterial invasion of the mucosa is clearly an over simplification. At least in the case of *E coli* O157:H7 and *S dysenteriae* type 1 infections, the important issue is the production of a cytotoxic toxin, irrespective of whether the bacteria invade the mucosa.

To date, ricin achieved its greatest notoriety in the 1978 'umbrella murder' of the Bulgarian broadcaster Georgi Markov.⁶ It is intriguing that the systemic effect of ricin in this unfortunate episode can be directly compared with the pathophysiology of bacterial colitis.

Finally, polymerase chain reaction amplification of *E coli* O157:H7 DNA encoding part of the cytotoxic toxin molecule may facilitate the detection of this bacterium in clinical specimens and in contaminated food.⁷

M F HEYWORTH
VA Medical Center (111)
2101 Elm Street, Fargo,
ND 58102, USA

- Jackson MP. Structure-function analyses of Shiga toxin and the Shiga-like toxins. *Microb Pathog* 1990; 8: 235-42.
- Endo Y, Mitsui K, Motizuki M, Tsurugi K. The mechanism of action of ricin and related toxic lectins on eukaryotic ribosomes: the site and the characteristics of the modification in 28 S ribosomal RNA caused by the toxins. *J Biol Chem* 1987; 262: 5908-12.
- Lampe KF. *Common poisonous and injurious plants*. Washington, DC: US Department of Health, and Human Services, 1981. HHS Publication No (FDA) 81-7006.
- Kopferschmitt J, Flesch F, Lugnier A, Sauder P, Jaeger A, Mantz JM. Acute voluntary intoxication by ricin. *Hum Toxicol* 1983; 2: 239-42.
- Karmali MA. Infection by verocytotoxin-producing *Escherichia coli*. *Clin Microbiol Rev* 1989; 2: 15-38.
- Knight B. Ricin - a potent homicidal poison. *BMJ* 1979; 1: 350-1.
- Gannon VPJ, King RK, Kim JY, Thomas EJG. Rapid and sensitive method for detection of Shiga-like toxin-producing *Escherichia coli* in ground beef using the polymerase chain reaction. *Appl Environ Microbiol* 1992; 58: 3809-15.

Reply

EDITOR,—We appreciate the comments from Dr Heyworth. Further details about toxin structure: function relations may be found in reference 21¹ in the leading article (*Gut* 1994; 35: 872-4). We have been proponents of the critical role of Shiga family toxins in the pathogenesis of bloody diarrhoea for over two decades, a concept that has become more ten-

able since the discovery of the 'non-invasive' toxin producing enterohaemorrhagic *E coli*, but which is still far from conclusively proved in human clinical disease.

While we pointed out the importance of rapid detection of this group of organisms in our leading article, and urged that a search for enterohaemorrhagic *E coli* becomes routine when any stool is examined for causes of bloody diarrhoea (an idea recently endorsed and recommended by a Consensus Conference of the American Gastroenterological Association that took place in July 1994), we do not believe that polymerase chain reaction is the best way at present to proceed, as Dr Heyworth suggests. This is because as of today, and for the foreseeable future, most clinical laboratories will not be doing polymerase chain reaction diagnosis at all. Our laboratory has developed monoclonal antibodies and an EIA toxin detection system for direct evaluation of free toxin in patient stools, which has been substantially increased in its sensitivity by Meridian Diagnostics of Cincinnati, Ohio. The preliminary clinical evaluation of this test, which is not restricted to the detection of *E coli* O157:H7 but detects all serotypes of toxin producing organisms, has been highly favourable. Meridian is carrying out further testing and evaluation to obtain approval by the US Food and Drug Administration. It would seem at this moment that a clinically useful early diagnostic test for enterohaemorrhagic *E coli* infection, independent of sorbitol fermentation patterns and the potential impact of antimicrobials on the viability of the micro-organisms, is imminent and could become available within a year's time. This test, in contrast with polymerase chain reaction, could be immediately used in all clinical laboratories that conduct routine microbiological, immunological, and serological assays for the diagnosis of infectious enteritis.

D W K ACHESON
G T KEUSCH
New England Medical Center and Tufts University
School of Medicine,
Boston 02111,
Massachusetts, USA

- Acheson DWK, Donohue-Rolfe A, Keusch GT. The family of Shiga and Shiga-like toxins. In: Alouf JE, Freer JH, eds. *Sourcebook of bacterial protein toxins*. London: Academic Press, 1991: 415-33.

Unethical research relating to *Helicobacter pylori*?

EDITOR,—We are writing concerning the paper by Vincent *et al* (*Gut* 1994; 35: 313-6) regarding the prevalence of *Helicobacter pylori* infection in cohabiting children. In that study, the authors examined the prevalence of *H pylori* infection among mentally retarded children residing in an institution. *H pylori* infection was confirmed by gastroscopy. 'Informed consent' was obtained from the families.

We believe that this study raises important ethical questions and concerns. Research involving children always raises the 'ethical' flag and the use of invasive procedures requires a very critical look at the risk versus benefit obtained. In most instances invasive procedures will be deemed to be unjustified in asymptomatic children. The problem is somewhat easier if you are dealing with a population of symptomatic children, but even in that group the motivation and clinical

practices of a group reporting the results of invasive procedures for diagnosis must be questioned.

This study describes endoscopic investigations in asymptomatic mentally retarded children. It is extremely difficult to imagine any benefit that the children could have gained from participation in the study. We are hard pressed to come up with any ethical justification for doing the study. Hammerschmidt and Gross¹ note that when a journal accepts a paper for publication after peer review that 'acceptance constitutes at least a *nihil obstat* [no objection] if not an *imprimatur* [official licence to print]'. Authority of the journal has been placed behind the paper announcing that it has been subjected to scrutiny and has been found to pass muster in terms of basic quality, novelty and potential importance.' There are two questions: firstly, should the study have been done at all and secondly should it have been accepted for publication?

We recognise that these concerns are serious, but they must be considered and considered now as the role of *H pylori* in the paediatric population is coming under increasing scrutiny.

D Y GRAHAM
M F GO
Department of Medicine,
VA Medical Center and Baylor College of Medicine,
Houston,
Texas 77030, USA
Correspondence to:
Dr Y Graham,
VA Medical Center (111D),
2002 Holcombe Blvd,
Houston, TX 77030, USA

- Hammerschmidt DE, Gross AG. Withdrawal of aegis? So what's that? *J Lab Clin Med* 1994; 123: 792-4.

Reply

EDITOR,—Ethical aspects regarding our paper (*Gut* 1994; 35: 313-6) were asked by one of the reviewers, and the necessary answers were given. Study protocol was much more detailed in the first version of our paper, but was shortened afterwards according to the reviewers' recommendations. Finally, only epidemiological and microbiological aspects were described in the published version.

We agree that such a study should not be done only for epidemiological purposes. Our study took place in a clinical context with therapeutic concerns as in the first instance, three cases of *H pylori* endoscopic gastritis were found in this institution and then treated with clinical ameliorations. These three children had endoscopy because of suspected oesophagitis and macroscopic nodules were seen in the antrum. This led us to perform biopsies, which showed the presence of *H pylori* gastritis. Specific *H pylori* treatment was given because we could not distinguish the potential role of the infection in the upper gastrointestinal complaints of these patients.

In severely neurologically impaired children, gastro-oesophageal reflux is common.¹ In our experience, 20-5% of these children suffer from gastro-oesophageal reflux. If not controlled in time, the disease can evolve and lead to severe complications such as oesophageal stricture or Barrett's oesophagus. In these children, gastro-oesophageal reflux is often associated with other disorders (nutritional, metabolic, respiratory, urological, or digestive diseases). It is widely admitted that gastro-oesophageal reflux is difficult to recognise on clinical grounds in mentally retarded