Our recent paper⁴ focused primarily on describing the histological features of microscopic colitis syndrome in our first 30 well defined cases. We have previously, however, studied different aetiological aspects in both lymphocytic⁵ and collagenous colitis.⁶ The importance of factor(s) in the faecal stream has also been discussed by others.⁷

B VERESS
R LÖFBERG
L BERGMAN
Department of Pathology,
Malmö University Hospital,
S-205 02 Malmö, Sweden

- 1 Lindström CG. 'Collagenous colitis' with watery diarrhoea. A new entity? *Pathol Eur* 1976; 11: 87-9.
- 2 Veress B, Bergman L. Collagenous colitis - a new disease. [Abstract]. *Svensk Kirurgi* 1983; 41: 82.
- 3 Veress B, Löfberg R, Bergman L. Lymfocytär och kollagen kolit - ytterligheter på samma spektrum? [Abstract]. *Svenska Läkaresällskapet Riksstämman* 1991; 100: 288.
- 4 Veress B, Löfberg R, Bergman L. Microscopic colitis syndrome. *Gut* 1995; 36: 880-6.
- 5 Einarsson K, Eusufzai S, Johansson U, Löfberg R, Theodorsson E, Veress B. Villous atrophy of distal ileum and lymphocytic colitis in a woman with bile acid malabsorption. *European Journal of Gastroenterology and Hepatology* 1992; 4: 585-90.
- 6 Eusufzai S, Löfberg R, Veress B, Einarsson K, Angelin B. Studies on bile acid metabolism in collagenous colitis: no evidence of bile acid malabsorption as determined by the SeHCAT test. *European Journal of Gastroenterology and Hepatology* 1992; 4: 317-21.
- 7 Riddell RH, Tanaka M, Mazzoleni G. Non-steroidal anti-inflammatory drugs as a possible cause of collagenous colitis: a case-control study. *Gut* 1992; 33: 683-6.

Helicobacter pylori reinfection

EDITOR,—We read with interest the article by Schütze *et al* (*Gut* 1995; 36: 831-3). It is difficult to agree with their statement 'conjugal transmission of reinfection has been shown to occur for the first time with near certainty'.

Besides transmission from the spouse, there are at least three possibilities of reinfection of their patients. (1) *H. pylori* has been demonstrated in the dental plaque of patients from developed¹⁻⁴ and developing^{5,6} countries. Furthermore, *H. pylori* in the dental plaque persists despite its clearance from the gastric mucosa after treatment.⁶ Hence, dental plaque is considered a potential source of reinfection by the same strain of *H. pylori*.^{7,8} (2) It is possible that both the patient and the spouse had acquired infection of the same strain of *H. pylori* from a common source. Contaminated water supply has been previously shown to act as common reservoir of infection.⁹ The fact that the prevalence of serum antibodies to *H. pylori* and hepatitis A virus in different age groups of subjects, both in developing (50% at 5 years) and developed (50% at 50 years) countries is comparable, shows faeco/oral mode of transmission of *H. pylori* and hepatitis A virus.¹⁰ (3) Intra-familial spread from another member of the family has been reported.¹¹ The study does not provide any data of other family members or the exact mode of transmission (kissing?) from the spouse.

A Y PHADKE
H G DESAI
Jaslok Hospital and Research Centre,
Bombay 400 026, India

- 1 Nguyen AMH, Engstrand L, Genta RM, Graham DY, El-Zaatari FAK. Detection of *Helicobacter pylori* in dental plaque by reverse transcription-polymerase chain reaction. *J Clin Microbiol* 1993; 31: 783-7.
- 2 Mapstone NP, Lynch DAF, Lewis FA, Axon ATR, Topkins DS, Dixon MF, *et al*.

Identification of *Helicobacter pylori* DNA in the mouths and stomachs of patients with gastritis using PCR. *J Clin Pathol* 1993; 46: 540-3.

- 3 Olsson K, Wadstrom T, Tsykiewicz T. *H. pylori* in dental plaques. *Lancet* 1993; 341: 956-7.
- 4 Banatwala N, Lopez CR, Owen R, Abdi Y, Davies G, Hardie J, *et al*. *Helicobacter pylori* in dental plaque. *Lancet* 1993; 341: 380.
- 5 Majumdar P, Shah SM, Dhunjibhoy KR, Desai HG. Isolation of *Helicobacter pylori* from dental plaques in healthy volunteers. *Ind J Gastroenterol* 1990; 9: 270-2.
- 6 Desai HG, Gill HH, Shankaran K, Mehta PR, Prabhu SR. Dental plaque: a permanent reservoir of *Helicobacter pylori*. *Scand J Gastroenterol* 1991; 26: 689-95.
- 7 Shames B, Krajden S, Fuksa M, Babida C, Panter JL. Evidence for the occurrence of the same strain of campylobacter *pylori* in the stomach and dental plaque. *J Clin Microbiol* 1989; 27: 2849-50.
- 8 Khandekar K, Palmer KR, Eastwood MA, Scott AC, Desai M, Owen RJ. DNA fingerprinting of *Helicobacter pylori* from mouth and antrum of patients with chronic ulcer dyspepsia. *Lancet* 1991; 342: 751.
- 9 Klein PD, Graham DY, Gailloyr A, Opekun AR, Smith EO. Gastrointestinal Physiology Working Group. *Lancet* 1991; 337: 1503-6.
- 10 Gill HH, Desai HG. *Helicobacter pylori* and gastroduodenal disorders in India: lessons from epidemiology. [Editorial]. *J Clin Gastroenterol* 1993; 16: 6-9.
- 11 Langenberg W, Rauws EAJ, Widjijojusume A, Tytgat GNJ, Zanen HC. Identification of *Campylobacter pyloridis* isolates by restriction endonuclease DNA analysis. *J Clin Microbiol* 1986; 24: 414.

Reply

EDITOR,—In our publication an autoreinfection has been discussed as a possibility but simultaneously considered as highly unlikely as the reinfections occurred as late as after 14 and 43 months, respectively.

The possibility of a common exogenous source cannot be ruled out but it seems rather unlikely and has also been discussed in our publication. The possibility of a contaminated water supply is extremely unlikely as water supplies in Vienna originate from Alpine sources and are renowned for their excellent quality.

The patients studied in our project were married couples without any other family members.

K SCHÜTZE
Medical Department 1,
Hannusch Hospital,
Heinrich Collinstr 30,
A-1140 Vienna,
Austria

Gastric emptying in patients with insulin dependent diabetes mellitus

EDITOR,—The paper by Nowak *et al* (*Gut* 1995; 37: 23-9) reports on the highly variable gastric emptying rates in patients with diabetes mellitus. The finding of accelerated (apart from the classic delayed) gastric emptying in humans with insulin dependent diabetes mellitus is consistent with previous findings.¹ The authors investigated the possible correlation between gastric emptying and chronic renal failure caused by diabetes. As the values obtained in this study have a very wide distribution, they concluded that the influence of chronic renal failure on gastric emptying cannot be estimated from their own results. However, chronic renal failure is associated with autonomic neuropathy² like diabetes mellitus and it would have been important to know the authors' comments on the role of the autonomic nervous system in their patients with chronic renal failure resulting from diabetes.

In another study,³ small groups of patients

with chronic renal failure of non-diabetic aetiology given longterm dialysis showed different patterns of gastric emptying depending on the involvement of the autonomic nervous system. In subjects with no autonomic neuropathy gastric emptying was faster than normal. In subjects with only parasympathetic neuropathy gastric emptying was similar to controls but in subjects with both parasympathetic and sympathetic autonomy, it was delayed. In this study gastric emptying was measured from the sonographic registration of antral area variation, according to Bolondi.⁴ This method has a good correlation with the radioisotopic method.⁵ The spectrum of the gastric emptying curves of the patients studied by Nowak *et al* seems to be influenced by autonomic neuropathy in the same manner. We believe that a similar approach focused only on chronic renal failure resulting from diabetes mellitus will bring similar results. Indeed, as Nowak *et al* show, autonomic neuropathy is the main factor that changes the emptying of the stomach.

D L DUMITRASCU
University of Medicine and Pharmacy,
Third Medical Dept,
str Dragalina 2-5,
3400 Cluj, Romania

- 1 Oliveira M, Troncon L. Accelerated gastric emptying in diabetes mellitus. [Abstract]. *J Gastrointest Motil* 1992; 4: 118.
- 2 Ewing DJ, Winny R. Autonomic function in patients with renal failure on intermittent hemodialysis. *Nephron* 1975; 15: 424-9.
- 3 Dumitrascu DL, Barnert J, Kirschner T, Wienbeck M. Antral emptying of semisolid meal measured by real-time ultrasonography in chronic renal failure. *Dig Dis Sci* 1995; 40: 636-44.
- 4 Bolondi L, Bortolotti M, Santi V, Calletti T, Gaiani S, Labo G. Measurement of gastric emptying time by real time ultrasonography. *Gastroenterology* 1985; 89: 752-9.
- 5 Wedmann B, Schaffner J, Wegener M, Schmidt G, Coenen C, Ricken D. Sonographische Erfassung der Magenentleerung. *Z Gastroenterol* 1990; 28: 448-52.

Reply

EDITOR,—The study performed by Dr D L Dumitrascu *et al*¹ is indeed relevant to our article, which describes highly variable gastric emptying in patients with diabetes mellitus and we appreciate his comments. Our study showed that some diabetic patients have rapid gastric emptying, some have a normal rate of gastric emptying, while others have what is appreciated to be delayed gastric emptying—that is, 'diabetic gastroparesis.' The precise mechanism responsible for the variability in gastric emptying in diabetic patients is unknown although autonomic neuropathy has been incriminated. In our study nearly all of the patients showed evidence of parasympathetic autonomic neuropathy evidenced by an abnormal respiratory variation in heart rate (E:1 ratio). Those patients who had autonomic sympathetic neuropathy, evidenced by orthostatic hypotension, showed a significantly longer half time of gastric emptying than non-diabetic control subjects. Likewise, those patients who did not have evidence of sympathetic autonomic neuropathy showed a significantly faster gastric emptying half time (accelerated gastric emptying) than non-diabetic controls. Finally, our study showed that the duration of diabetes was significantly correlated with the gastric emptying shape coefficient, implying that a longer duration of diabetes is associated with a slower rate of gastric emptying. Our findings lead us to speculate that short-term diabetes mellitus is