

Leading article

Age related changes in gut physiology and nutritional status

With an aging population, the gastroenterologist is seeing increasing numbers of elderly patients with unexplained gastrointestinal disorders or malnutrition. If the presentation can be ascribed to normal aging or if no treatment is judged to be available, the patient may not be investigated. To make such an evaluation, however, the doctor must possess a thorough understanding of how gastrointestinal function and nutritional parameters change in the elderly. This review shows that in the light of current knowledge, active treatment will probably be seen as the most appropriate course to pursue in most cases.

Malnutrition in the elderly

The scale of the problem

Protein-energy malnutrition is common in the elderly. Approximately 10% of acute geriatric patients are malnourished on admission.¹ The prevalence is 50% in elderly people living in institutional care and even in free living people aged over 65 years, it is approximately 3% although most of these people have coexisting disease.^{2,3} Unless the possibility is considered, however, malnutrition is rarely diagnosed, as even in grossly undernourished patients, haematological and biochemical parameters can be normal.⁴ Even serum transferrin, transthyretin (prealbumin), and retinol binding protein, which are considered the best biochemical markers of acute malnutrition, can be normal in the chronic state.⁵

Does nutrient intake fall in old age?

In healthy people, earlier cross sectional studies suggested a fall in energy intake with aging, but this no longer seems to be the case. There was no difference in food intake in old compared with young socially privileged women in the USA, although old men did eat less,⁶ and in Finland and the USA, longitudinal data suggest that nutrient and energy intake does not change in healthy people in the eighth or ninth decade.^{7,8} Food intake is severely diminished in people in institutions, most of whom are physically disabled,⁹ and older studies had already suggested that the age related fall in energy intake is associated with decreasing physical ability.¹⁰ Recently, physically active elderly women were shown to eat more than sedentary women, but a moderate exercise training programme did not increase energy intake.¹¹ Psychological stresses such as bereavement are important causes of decreased appetite as is the inability to chew food because of poor dentition or a dry mouth, factors often neglected by physicians.^{12,13}

Vitamin and trace element deficiencies are well recognised in hospitalised patients. Of more concern are the widespread deficiencies in the 'healthy' elderly. In an American survey, large numbers of healthy people aged over 65 years had very low intakes of vitamins A, E, calcium, and zinc. Not surprisingly, subclinical deficiencies of all vitamins have been widely shown. Indeed, the recommended daily intake for many vitamins, which are

based on studies in younger people are probably too low for the elderly.¹⁴

Changes in anthropometric parameters in old age

Advancing years are accompanied by loss of height. The proportion of body fat increases and lean body mass decreases as shown by measurements of total body potassium and total body water, or by densitometry or anthropometry; there may also be a shift of fat to non-subcutaneous sites with ageing. Anthropometric data for healthy people over 65 are available to allow useful comparisons to be made in this age group. Measurements of triceps skinfold thickness and mid-arm circumference with the derived arm muscle circumference in healthy normal subjects in Europe and North America show no pronounced change in these indicators of nutrition until at least 75 years of age. There is also no age related fall in plasma haemoglobin or plasma protein concentration in healthy people.

Changes in gut function with age

Mouth

Some 40% of healthy elderly people complain of dry mouth, but whereas unstimulated salivary flow probably decreases, in response to stimulation, salivation is unchanged in both healthy and edentulous people.¹⁵⁻¹⁷ Medication, diabetes, arthritis, and physical dependence are all associated with mouth dryness and those affected are more likely to report difficulty eating and communicating. However, taste and smell sensitivity do decrease with age.^{18,19} It is presumed that this decreased sensitivity results in foods tasting similar, with elderly people consequently eating more bland diets. The manufacturing industry is tackling this problem by developing foodstuffs with sharper tastes, which might be preferred by older people.

Healthy elderly subjects open their mouths less widely and chew with less power than young people, a finding that parallels the loss of muscle bulk with age, and which is greater in edentulous people.^{20,21} It has been suggested that some of the changes noted are not just a normal part of aging but represent preclinical manifestations of neurodegenerative disorders such as Parkinson's disease.¹⁷

Oesophagus

Classic radiographical studies showed pharyngeal hypotonicity with incomplete opening of cricopharyngeus muscle in 22% of asymptomatic subjects over age 65.^{22,23} Manometric studies have confirmed that the upper oesophageal sphincter resting pressure is lower in old age with delayed relaxation on swallowing accompanied by increased pharyngeal contraction pressures.^{24,25} In a radiological study of 56 asymptomatic patients in their 80s, two thirds had difficulties in the oral phase of swallowing, one

quarter had dysfunction of the pharyngeal phase, and oesophageal abnormalities were described in almost 40%.²³ The prevalence of silent acid reflux does not change with age.²⁶

Stomach

Most studies suggest that a large proportion of 'normal' asymptomatic subjects above 60 years of age have atrophic gastritis. Bird *et al* found only 50 histologically normal gastric biopsy specimens in 201 asymptomatic subjects aged 65 to 90 years, but there was no relation between age and the degree of gastric atrophy in this series.²⁷ Furthermore, in a study to evaluate the development of gastritis, no subject aged over 60 with initially normal mucosa or superficial gastritis went on to develop atrophic gastritis during a 10 year follow up.²⁸ Taken together, these findings suggest that gastric atrophy is not a normal part of aging, rather that it is due to specific abnormal mechanisms (see below).

Baron's classic study suggested that both basal and peak gastric acid output decrease with increasing age.²⁹ This change is probably due to the increased prevalence of atrophic gastritis, however, with most healthy old people actually maintaining gastric acid secretion.³⁰ Recently, serum gastrin concentration has been shown to be increased only in *Helicobacter pylori* infected subjects but not in elderly uninfected subjects.^{30 31} Pepsin secretion does not change with age either.³²

The importance of *H pylori* infection in the pathogenesis of gastric atrophy and hypochlorhydria is now well recognised and previous or current *H pylori* infection is found in most patients with atrophic gastritis. More surprisingly, *H pylori* might also have a role in the development of pernicious anaemia as in a recent study, 83% of patients with clinical pernicious anaemia also demonstrated antibodies to *H pylori*.^{33 34} Whether these findings are of pathogenetic significance is not yet clear. The histological and functional changes in the stomach, however, previously attributed to aging can probably all be explained by the presence of this pathogenic agent, the prevalence of which also rises with age.

Isotope studies have shown a considerable prolongation of gastric emptying for liquids in healthy elderly subjects compared with younger controls. However, gastric emptying for solids is unchanged in old age and the gastric electrical rhythm is maintained in the elderly.³⁵⁻³⁷

Small intestine

There is no age related change in small intestinal anatomy and enterocyte height and intraepithelial lymphocyte counts are unchanged.^{38 39} Increased cell proliferation in response to injury has been shown in a rodent model which, if also true in humans, would present a possible mechanism for the increased susceptibility to gastrointestinal cancers in the elderly.⁴⁰

The healthy small intestine also maintains its absorptive function for carbohydrates with no change in duodenal brush border membrane enzyme activity of glucose transport; and for fats, although adaptive reserve is more limited.⁴¹⁻⁴⁵ Vitamin B12 absorption is also maintained in the healthy elderly.^{44 45} 'Malabsorption in the elderly' does not seem to be either a non-specific occurrence or to be due to a specific age related process, but should be explicable in terms of recognised abnormal mechanisms. Bacterial overgrowth may be a comparatively common cause reflecting the high incidence of gastric hypochlorhydria and jejunal diverticulosis,^{41 46 47} and although tests of absorptive capacity are normal in most cases, empirical

antibiotic treatment has been shown to have a profound impact on nutritional state.^{48 49}

Zinc absorption is also decreased from a mixed meal in the elderly, but excretion is similarly diminished so that zinc balance remains.⁵⁰ Calcium absorption probably also declines over 60 years of age,⁵¹ but is also lessened in the presence of gastric atrophy. Recent work suggests a fall in vitamin D receptor concentration in the small bowel may decrease intestinal responsiveness to vitamin D activity.⁵² Parathyroid hormone secretion and 1,25 dihydroxyvitamin D production may then need to rise to maintain serum calcium homeostasis with the consequence of increased bone loss. The implications of these changes are unclear and further study into their effect on health maintenance and chronic disease in the elderly are required.⁵³

Pancreas

There is a steady increase in the calibre of the main pancreatic duct with age, and the other branches show areas of focal dilatation or stenosis not associated with any other abnormality.⁵⁴ *x* Rays obtained during endoscopic retrograde pancreatography should be interpreted with caution, but this finding is otherwise of no obvious physiological relevance.

Aging changes pancreatic secretion through a decrease in flow rate, bicarbonate and enzyme secretion while calcium secretion is increased. Pancreatic secretion has also been shown to fall significantly upon repeated stimulation.⁵⁵ Thus, rare cases of pancreatic exocrine insufficiency could be explained on the basis of aging alone without malnutrition.

Splanchnic blood flow

Splanchnic blood flow declines with increasing age both absolutely and as a fraction of cardiac output.^{56 57} The splanchnic circulation is also susceptible to hypoxia associated with cardiac or respiratory insufficiency and to hypovolaemia or systemic hypotension.⁵⁸ Very few cases of painless occult malabsorption resulting from vascular insufficiency have been described in geriatric patients. The classical syndromes associated with vascular insufficiency of both small and large intestine, which are almost always painful are usually seen in very elderly people.

Gastrointestinal motility

Constipation is a common symptom in the elderly. Orocaecal transit time is not generally changed in elderly volunteers⁵⁹ although it does seem to be exquisitely sensitive to thyroid hormone status and can be considerably prolonged even in subclinical hypothyroidism.⁶⁰ Colonic transit may slow with aging, but is highly variable⁶¹ and interestingly, strength training shortens bowel transit, with the effect being limited entirely to the colon.⁶²

Rectum

Mechanical changes in the rectum probably cause most of the problems old people suffer in evacuating as well as the increasing prevalence of faecal incontinence with increasing age. A reduction in rectal wall elasticity means that tonic activity of the external sphincter is lost at a smaller volume.⁶³ An age dependent increase in rectal pressure threshold to produce initial sensation of rectal filling has been found⁶⁴ and structural changes also occur in the internal anal sphincter, which has been shown by endosonography to be thicker and hyperechoic in older subjects.⁶⁵ Maximum resting anal pressure and maximum

squeeze pressure have been shown to decline with age, particularly in postmenopausal women. The decrease in squeeze anal pressure has been shown in one study to be accompanied by increased mean pudendal nerve terminal motor latency in women in the fifth decade, indicating damage to this nerve. Compensatory reinnervation of this muscle does, however, seem to develop after the menopause.⁶⁶ The difference between maximum resting anal pressure and rectal pressure also falls in old age.⁶⁷ Taken together, these findings may explain the increased prevalence of faecal incontinence in the elderly and suggest that at least in women, oestrogen treatment may be valuable. This hypothesis remains to be tested.

Conclusion

Few gastrointestinal functions decline to an important extent as a result of old age alone and there is little clinical evidence that significant malnutrition occurs in any normal elderly person as a result of the aging process itself. Nevertheless, decreased gastrointestinal reserve makes older people highly sensitive to minor insults and decompensation can rapidly occur. Drugs appreciably affect taste sensation, which is already blunted and psychological as well as physical disability can have a major impact on appetite. Malabsorption can be caused by gastric hypochlorhydria with small bowel bacterial overgrowth and while gastrointestinal dysmotility can be caused by subclinical hypothyroidism, it can improve in response to physical exercise. Evidence is now mounting that thorough investigation of gastrointestinal disturbances in elderly patients coupled with intensive nutritional support can make a very real impact on their outcome. Gastroenterologists should therefore seek out and actively treat gastrointestinal disorders in the elderly and not just ascribe them to old age.

L B LOVAT

Department of Medicine,
Hammersmith Hospital,
Du Cane Road,
London W12 0NN

- Kemm JR, Allcock J. The distribution of supposed indicators of nutritional status in elderly patients. *Age Ageing* 1984; **13**: 21-8.
- Nutrition and health in old age. Report on health and social subjects No 16 DHSS, 1979.
- Morgan DB, Newton HM, Schorah CJ, Jewitt MA, Hancock MR, Hullin RP. Abnormal indices of nutrition in the elderly: a study of different clinical groups. *Age Ageing* 1986; **15**: 65-76.
- Lipski PS, Torrance A, Kelly PJ, James OF. A study of nutritional deficits of long-stay geriatric patients. *Age Ageing* 1993; **22**: 244-55.
- Lemonnier D, Acher S, Boukaiba N, Flament C, Doucet C, Piau A, et al. Discrepancy between anthropometry and biochemistry in the assessment of the nutritional status of the elderly. *Eur J Clin Nutr* 1991; **45**: 281-6.
- Munro HN, McGandy RB, Hartz SC, Russell RM, Jacob RA, Otradovec CL. Protein nutrition of a group of free-living elderly. *Am J Clin Nutr* 1987; **46**: 586-92.
- Rasanen L, Mutanen M, Pekkanen J, Laitinen S, Koski K, Halonen S, et al. Dietary intake of 70- to 89-year-old men in eastern and western Finland. *J Intern Med* 1992; **232**: 305-12.
- Vellas BJ, Albarede JL, Garry PJ. Diseases and aging: patterns of morbidity with age; relationship between aging and age-associated diseases. *Am J Clin Nutr* 1992; **55**: 1225S-30S.
- Lowik MR, Schneijder P, Hulshof KF, Kistemaker C, Sleutel L, van Houten P. Institutionalized elderly women have lower food intake than do those living more independently (Dutch Nutrition Surveillance System). *J Am Coll Nutr* 1992; **11**: 432-40.
- Stanton BR, Exton-Smith AN. *A longitudinal study of the dietary intake of elderly women*. London: King Edward's Hospital Fund for London, 1970.
- Butterworth DE, Nieman DC, Perkins R, Warren BJ, Dotson RG. Exercise training and nutrient intake in elderly women. *J Am Diet Assoc* 1993; **93**: 653-7.
- Rosenbloom CA, Whittington FJ. The effects of bereavement on eating behaviors and nutrient intakes in elderly widowed persons. *J Gerontol* 1993; **48**: S223-S9.
- Laurin D, Brodeur JM, Bourdages J, Vallee R, Lachapelle D. Fibre intake in elderly individuals with poor masticatory performance. *J Can Dent Assoc* 1994; **60**: 443-6.
- Suter PM, Russell RM. Vitamin requirements of the elderly. *Am J Clin Nutr* 1987; **45**: 501-12.
- Sherm RJ, Fox PC, Le SH. Influence of age on the secretory rates of the human minor salivary glands and whole saliva. *Arch Oral Biol* 1993; **38**: 755-61.
- Gilbert GH, Heft MW, Duncan RP. Mouth dryness as reported by older Floridians. *Community Dent Oral Epidemiol* 1993; **21**: 390-7.
- Baum BJ, Bodner L. Aging and oral motor function: evidence for altered performance among older persons. *J Dent Res* 1983; **62**: 2-6.
- Doty RL, Shaman P, Applebaum SL, Giberson R, Sikorski L, Rosenberg L. Smell identification ability: changes with age. *Science* 1984; **226**: 1441-3.
- Weiffenbach JM, Baum BJ, Burghauer R. Taste thresholds: quality specific variation with human aging. *J Gerontol* 1982; **37**: 372-7.
- Karlsson S, Persson M, Carlsson GE. Mandibular movement and velocity in relation to state of dentition and age. *J Oral Rehabil* 1991; **18**: 1-8.
- Newton JP, Yemm R, Abel RW, Menhinick S. Changes in human jaw muscles with age and dental state. *Gerodontology* 1993; **10**: 16-22.
- Piaget F, Fouillet L. Le pharynx et l'oesophage senile: etude clinique, radiologique et radio-cinematographique. *J Med Lyon* 1959; **40**: 961-5.
- Ekberg O, Feinberg MJ. Altered swallowing function in elderly patients without dysphagia: radiologic findings in 56 cases. *Am J Roentgenol* 1991; **156**: 1181-4.
- Fulp SR, Dalton CB, Castell JA, Castell DO. Aging-related alterations in human upper esophageal sphincter function. *Am J Gastroenterol* 1990; **85**: 1569-72.
- Wilson JA, Pryde A, Macintyre CC, Maran AG, Heading RC. The effects of age, sex, and smoking on normal pharyngo-oesophageal motility. *Am J Gastroenterol* 1990; **85**: 686-91.
- Fass R, Sampliner RE, Mackel C, McGee D, Rappaport W. Age- and gender-related differences in 24-hour esophageal pH monitoring of normal subjects. *Dig Dis Sci* 1993; **38**: 1926-8.
- Bird T, Hall MR, Schade RO. Gastric histology and its relation to anaemia in the elderly. *Gerontology* 1977; **23**: 309-21.
- Ihamaki T, Kekki M, Sipponen P, Siurala M. The sequelae and course of chronic gastritis during a 30- to 34-year bioptic follow-up study. *Scand J Gastroenterol* 1985; **20**: 485-91.
- Baron JH. Studies of basal and peak acid output with an augmented histamine meal. *Gut* 1963; **4**: 136-44.
- Katelaris PH, Seow F, Lin BP, Napoli J, Ngu MC, Jones DB. Effect of age, Helicobacter pylori infection, and gastritis with atrophy on serum gastrin and gastrin acid secretion in healthy men. *Gut* 1993; **34**: 1032-7.
- Goldschmied M, Barnett CC, Schwarz BE, Karnes WE, Redfern JS, Feldman M. Effect of age on gastric acid secretion and serum gastrin concentrations in healthy men and women. *Gastroenterology* 1991; **101**: 977-90.
- McCloy RF, Arnold R, Bardhan KD, Cattani D, Klinkenberg KE, Maton PN, et al. Pathophysiological effects of long-term acid suppression in man. *Dig Dis Sci* 1995; **40** (Suppl): 96-120S.
- Lechago J, Correa P. Prolonged achlorhydria and gastric neoplasia: is there a causal relationship? *Gastroenterology* 1993; **104**: 1554-7.
- Wyatt JJ, Shallcross TM, Crabtree JE, Heatley RV. Helicobacter pylori, gastritis, and peptic ulceration in the elderly. *J Clin Pathol* 1992; **45**: 1070-4.
- Riezzo G, Pezzolla F, Giorgio I. Effects of age and obesity on fasting gastric electrical activity in man: a cutaneous electrogastrographic study. *Digestion* 1991; **50**: 176-81.
- Kao CH, Lai TL, Wang SJ, Chen GH, Yeh SH. Influence of age on gastric emptying in healthy Chinese. *Clin Nucl Med* 1994; **19**: 401-4.
- Moore JG, Tweedy C, Christian PE, Datz FL. Effect of age on gastric emptying of liquid-solid meals in man. *Dig Dis Sci* 1983; **28**: 340-4.
- Corazza GR, Frazzoni M, Gatto MR, Gasbarrini G. Ageing and small-bowel mucosa: a morphometric study. *Gerontology* 1986; **32**: 60-5.
- Warren PM, Pepperman MA, Montgomery RD. Age changes in small-intestinal mucosa. *Lancet* 1978; **ii**: 849-50.
- Atilasoy E, Holt PR. Gastrointestinal proliferation and aging. *J Gerontol* 1993; **48**: B43-9.
- Wallis JL, Lipski PS, Mathers JC, James OF, Hirst BH. Duodenal brush-border mucosal glucose transport and enzyme activities in aging man and effect of bacterial contamination of the small intestine. *Dig Dis Sci* 1993; **38**: 403-9.
- Haeney MR, Culank LS, Montgomery RD, Sammons HG. Evaluation of xylose absorption as measured in blood and urine: a one-hour blood xylose screening test in malabsorption. *Gastroenterology* 1978; **75**: 393-400.
- Holt PR, Balint JA. Effects of aging on intestinal lipid absorption. *Am J Physiol* 1993; **264**: G1-6.
- Arora S, Kassarjian Z, Krasinski SD, Croffey B, Kaplan MM, Russell RM. Effect of age on tests of intestinal and hepatic function in healthy humans. *Gastroenterology* 1989; **96**: 1560-5.
- Webster SG, Wilkinson EM, Gowland E. A comparison of fat absorption in young and old subjects. *Age Ageing* 1977; **6**: 113-7.
- Roberts SH, James O, Jarvis EH. Bacterial overgrowth syndrome without 'blind loop': A cause for malnutrition in the elderly. *Lancet* 1977; **ii**: 1193-6.
- Husebye E, Skar V, Hoverstad T, Melby K. Fasting hypochlorhydria with gram positive gastric flora is highly prevalent in healthy old people. *Gut* 1992; **33**: 1331-7.
- Haboubi NY, Montgomery RD. Small-bowel bacterial overgrowth in elderly people: clinical significance and response to treatment. *Age Ageing* 1992; **21**: 13-9.
- Saltzman JR, Kowdley KV, Pedrosa MC, Sepe T, Golner B, Perrone G, et al. Bacterial overgrowth without clinical malabsorption in elderly hypochlorhydric subjects. *Gastroenterology* 1994; **106**: 615-23.
- Turmlund JR, Durkin N, Costa F, Margen S. Stable isotope studies of zinc absorption and retention in young and elderly men. *J Nutr* 1986; **116**: 1239-47.
- Bullamore JR, Wilkinson R, Gallagher JC, Nordin BE, Marshall DH. Effect of age on calcium absorption. *Lancet* 1970; **ii**: 535-7.
- Ebeling PR, Sandgren ME, DiMaggio EP, Lane AW, DeLuca HF, Riggs BL. Evidence of an age-related decrease in intestinal responsiveness to vitamin D: relationship between serum 1,25-dihydroxyvitamin D3 and intestinal vitamin D receptor concentrations in normal women. *J Clin Endocrinol Metab* 1992; **75**: 176-82.
- Russell RM. Changes in gastrointestinal function attributed to aging. *Am J Clin Nutr* 1992; **55**: 1203-7S.
- Sahel J, Cros RC, Lombard C, Sarles H. Morphometrique de la pancreatographie endoscopique normale du sujet age. *Gastroenterologie Hepatologie* 1979; **15**: 574-7.
- Gullo L, Ventrucci M, Naldoni P, Pezzilli R. Aging and exocrine pancreatic function. *J Am Geriatr Soc* 1986; **34**: 790-2.

- 56 Zoli M, Iervese T, Abbati S, Bianchi GP, Marchesini G, Pisi E. Portal blood velocity and flow in aging man. *Gerontology* 1989; 35: 61–5.
- 57 Bender AD. The effect of increasing age on the distribution of peripheral blood flow in man. *J Am Geriatr Soc* 1965; 13: 192–201.
- 58 Almy TP. Factors leading to digestive disorders in the elderly. *Bull NY Acad Med* 1981; 57: 709–17.
- 59 Husebye E, Engedal K. The patterns of motility are maintained in the human small intestine throughout the process of aging. *Scand J Gastroenterol* 1992; 27: 397–404.
- 60 Rahman Q, Haboubi NY, Hudson PR, Lee GS, Shah IU. The effect of thyroxine on small intestinal motility in the elderly. *Clin Endocrinol (Oxf)* 1991; 35: 443–6.
- 61 Madsen JL. Effects of gender, age, and body mass index on gastrointestinal transit times. *Dig Dis Sci* 1992; 37: 1548–53.
- 62 Koffler KH, Menkes A, Redmond RA, Whitehead WE, Pratley RE, Hurley BF. Strength training accelerates gastrointestinal transit in middle-aged and older men. *Med Sci Sports Exerc* 1992; 24: 415–9.
- 63 Ihre T. Studies on anal function in continent and incontinent patients. *Scand J Gastroenterol* 1974; 25: 1–64.
- 64 Akervall S, Nordgren S, Fasth S, Oresland T, Pettersson K, Hulten L. The effects of age, gender, and parity on rectoanal functions in adults. *Scand J Gastroenterol* 1990; 25: 1247–56.
- 65 Burnett SJ, Bartram CI. Endosonographic variations in the normal internal anal sphincter. *Int J Colorectal Dis* 1991; 6: 2–4.
- 66 Laurberg S, Swash M. Effects of aging on the anorectal sphincters and their innervation. *Dis Colon Rectum* 1989; 32: 737–42.
- 67 Haadem K, Dahlstrom JA, Ling L. Anal sphincter competence in healthy women: clinical implications of age and other factors. *Obstet Gynecol* 1991; 78: 823–7.