

F253

**OCTREOTIDE DECREASES GASTRIC MUCOSAL BLOOD FLOW: A CONTROLLED ASSESSMENT BY LASER DOPPLER FLOMETERY.**

**A K KUBBA**, G H HAYDON, P C HAYES, K R PALMER WESTERN GENERAL HOSPITAL & THE ROYAL INFIRMARY OF EDINBURGH.

The somatostatin analogue octreotide has been used to stop peptic ulcer haemorrhage, although its mechanism of action is unclear. It may exert an effect by reducing gastroduodenal mucosal blood flow.

Mucosal blood flow was measured using a Laser Doppler Flowmeter. The Doppler probe was positioned in the upper gastrointestinal tract (either endoscopically or via a small gastrotomy) of seven rabbits. Preliminary experiments showed the method to be reproducible, non invasive and simple.

Mucosal blood flow was decreased in a dose dependent manner in the body (193.5U to 71.25U) and antrum (154.8U to 34.7U) of the stomach and duodenum (187.5U to 67.5U) by doses of octreotide ranging from 10-50 microgram/kg of body weight. The change was not associated with systemic haemodynamic changes. Vasopressin (1 microgram/kg) but not normal saline also reduced gastroduodenal mucosal blood flow, but this was associated with profound bradycardia and a fall in non gastrointestinal tissue perfusion.

Gastroduodenal mucosal blood flow is reduced in a dose dependent manner by octreotide suggesting that this agent may be helpful in the management of bleeding from gastroduodenal ulcer disease and gastritis.

F255

**Experimental studies of injection therapy for ulcer haemorrhage in rabbits**

**Kubba A K**, Palmer K R  
Gastro-intestinal Unit, Western General Hospital, Edinburgh, Scotland.

Endoscopic injection therapy is a well established method of controlling peptic ulcer haemorrhage but the optimum solution and the mechanism involved in inducing haemostasis are unknown. Efficacy and tissue effects of various therapeutic agents in the control of gastric mucosal haemorrhage were studied in 10 rabbits. Forty bleeding mucosal ulcers created at gastrotomy (Blood loss > 1.5 ml/min) were studied. Adrenaline (1:100,000), thrombin, fibrin (thrombin plus fibrinogen), 5% ethanolamine and 50% dextrose were injected into the bleeding ulcers.

Sclerosants were found to be least effective in the control of bleeding and were associated with significant tissue necrosis. Although all the other solutions significantly decreased blood loss within 30 minutes of injection therapy (median blood loss = 0.25ml/min), only a mixture of adrenaline plus thrombin and sprayed fibrin achieved complete haemostasis within 2 minutes of treatment and with no recurrence of bleeding. Neither modality caused significant tissue damage.

Histological examination showed that no injected solution caused arterial thrombosis when injected next to a major ear artery.

We conclude that a combination of dilute adrenaline plus human thrombin may represent optimal haemostatic therapy for peptic ulcer haemorrhage.

**Stomach and bleeding F254-F258**

F254

**ANTIBIOTIC TRANSFER ACROSS THE GASTRIC MUCOSA - AN IN VITRO MODEL.** AF Goddard & RC Spiller. Division of Gastroenterology, University Hospital, Nottingham. NG7 2UH. UK

**Introduction** The mechanisms underlying transfer of antimicrobial agents for the treatment of *H. pylori* across the gastric mucosal barrier are poorly understood. An in vitro model was designed and validated to investigate these mechanisms.

**Methods** Gastric corpus mucosa was obtained from male Wistar rats (300-350g), stripped of muscle layers, mounted in Ussing chambers and bathed in oxygenated Krebs solution. Metronidazole or amoxycillin were added on either side of the mucosa and timed samples taken from the opposite side. Antibiotic concentrations were measured using HPLC. Metronidazole flux from serosa to mucosa ( $J_{S \rightarrow M}$ ) was measured (n=8 for each experiment): 1) under basal conditions; 2) compared with mucosal to serosal flux ( $J_{M \rightarrow S}$ ); 3) in chambers of different cross-section; 4) with stimulation by 100 mM histamine and carbachol; 5) in comparison with amoxycillin flux under basal conditions.  $J_{M \rightarrow S}$  for metronidazole was measured with mucosal pH at 2.2, 2.7, 3.2 and 7.4.

**Results** Reservoir concentrations varied between 0.2 and 0.4 mM between but not within each experiment and were correlated ( $P < 0.001$ ) with metronidazole flux. Flux results are expressed as mean (SEM) in  $\mu\text{mol} \cdot \text{cm}^{-2} \cdot \text{hr}^{-1}$ : 1) Metronidazole  $J_{S \rightarrow M}$  was 48.9 (8.7) under basal conditions; 2)  $J_{S \rightarrow M}$  was 79.2 (13.5) compared with  $J_{M \rightarrow S}$  of 98.3 (37.3) ( $P = \text{NS}$ ); 3)  $J_{S \rightarrow M}$  with a chamber diameter of 1.2 cm was 39.8 (2.4) and 40.8 (5.8) with a 0.9 cm diameter ( $P = \text{NS}$ ); 4)  $J_{S \rightarrow M}$  was 68.2 (6.2) with stimulation and 73.4 (4.6) without ( $P = \text{NS}$ ); 5) metronidazole  $J_{S \rightarrow M}$  was 39.8 (2.4) compared to amoxycillin  $J_{S \rightarrow M}$  of 22.4 (2.4) ( $P < 0.01$ ). When mucosal pH was changed,  $J_{M \rightarrow S}$  was proportional to the unionised concentration on the mucosal side ( $P < 0.001$ ).

**Conclusions** Metronidazole transfer across the gastric mucosa occurs by simple diffusion and pH partition occurs on the mucosal side. This model shows selectivity for different antibiotics and will be useful for the study of other therapeutic agents in the treatment of *H. pylori*.

AFG is in receipt of an Astra Research Fellowship.

F256

**ROLE OF TIPSS IN THE MANAGEMENT OF GASTRIC VARICEAL HAEMORRHAGE.**

K.I. Bzeizi, A.J. Stanley, R. Jalan, D. Redhead, K. Simpson, P.C. Hayes, Centre of Liver and Digestive Diseases, Royal Infirmary of Edinburgh

Bleeding from gastric varices is associated with a high risk of early rebleeding and mortality. We assessed the impact of transjugular intrahepatic portosystemic shunt (TIPSS) on the management of bleeding from gastric varices and compared this to the treatment prior to introduction of TIPSS. A total of 42 patients were referred with gastric variceal bleed during the period 1989-1995. 39 were considered for analysis. 30 patients (12 males, 18 females, mean age = 49.1 years) with Child B cirrhosis had TIPSS inserted after their initial bleed (group I). Group II patients (6 males, 3 females) referred before TIPSS was available, were treated either endoscopically (6 patients) or by surgery (1 spleno-renal, and 2 had oesophageal transection with splenectomy).

Both groups were matched for age and Child score. Mean portal pressure gradient (PPG) was reduced from  $13.1 \pm 1.42$  to  $6.3 \pm .82$  mmHg ( $p < 0.01$ ) post TIPSS. Over a mean follow up of 18.1 months, rebleeding rate in group I was 13.3% compared with 66.6% in group II ( $p < 0.01$ ) who had a mean follow up of 13.4 months. Mortality 6 weeks following the index haemorrhage was 23.3% in group I with no bleeding related deaths (sepsis in 3, and 4 with hepato-renal failure). This was not significant when compared with the number of deaths in group II (33.3%), although this group included 2 deaths due to rebleeding. Encephalopathy occurred in 5 patients post TIPSS (16.6%).

**CONCLUSION:** Rebleeding from gastric varices is significantly lower in patients treated with TIPSS as compared with those treated differently. TIPSS can be performed successfully in this group of patients who frequently are not fit to undergo surgery. We currently consider TIPSS as the treatment of choice for bleeding gastric varices.

### Audit (upper GI tract) F259-F264

F257

#### DIRECT MEASUREMENT OF POST-PRANDIAL PORTAL HAEMODYNAMICS.

Stanley A.J., Forrest E.H., #Redhead D.N., Hayes P.C.  
 Departments of Medicine and #Radiology, Royal Infirmary of Edinburgh, EH3 9YW.

Post-prandial changes in portal haemodynamics have been reported using derived methods or doppler ultrasonography. The presence of TIPSS allows direct assessment of portal venous response to a protein meal.

**Methods:** We assessed the portal and systemic haemodynamic response to a protein meal in 10 cirrhotic patients with TIPSS in situ (mean age 60.2 (3.7) years, mean Childs-Pugh score 7.8 (0.6)). Portal venous pressure (PVP) and flow (PVF) (measured by direct reverse thermodilution method with double thermistor catheter inserted in portal vein), right atrial pressure (RAP) and mean arterial pressure (MAP) were recorded whilst fasting and every 15 minutes for one hour following a 505 kcal. protein based meal. Pre- and post-prandial portal venous blood gas measurements were obtained.

**Results:** The maximal change in haemodynamic parameters from fasting levels are shown below. All measurements returned to normal 60 minutes following the meal and there were no significant changes between pre- and post-prandial portal venous blood gas analyses.

	MAP	RAP	PVP	PVF
Fasting:	79.5 (2.6)	1.9 (0.8)	9.9 (1.2)	1137.7 (162.1)
Post-prandial:	78.8 (1.9)	2.6 (0.7)	13.0 (1.3)	1642.2 (256.6)
p. value:	NS	NS	p<0.001	p<0.01

Results are expressed as mean (standard error). Pressures are shown in mmHg and flow in ml/min.

**Conclusion:** After a protein meal, PVP and PVF increase significantly whilst MAP and RAP remain unchanged. This technique allows direct assessment of therapeutic strategies to reduce post-prandial hyperaemia.

F259

#### INDICATIONS FOR MAINTENANCE ACID SUPPRESSION AT THE GENERAL PRACTICE/HOSPITAL INTERFACE

P Connor<sup>1</sup>, A McDonald<sup>\*</sup>, D N Lynch<sup>\*</sup>, P P Garratt<sup>\*</sup>, C D Gillen<sup>1</sup>  
 Gastroenterology Department, RNH Haslar<sup>1</sup> and Stoke Road Surgery<sup>\*</sup>, Gosport, Hampshire PO12 1PA

**Introduction:** Repeat prescription of H2 receptor antagonists (H2RA) and proton pump inhibitors (PPI) in gastroesophageal reflux disease (GORD) and peptic ulcer is commonplace and expensive.

**Aim:** We sought to determine the indication for maintenance acid suppressive therapy at the general practice/ hospital interface.

**Method:** We examined the GP case records of 312 patients on long term acid suppression representing 4% of the total practice list (8,185) and consuming 11% (£69,000) of the total annual drug budget (£640,000).

**Results:** 246 patients (79%) had GORD, 46 patients duodenal ulcer (15%) and 20 patients gastric ulcer (6%). Thirty-one of the 66 peptic ulcer patients (47%) had been tested for H. pylori, (13 positive).

The diagnosis was established by gastroscopy in 218 patients (70%) and barium meal in 13 (4%). Neither investigation had been undertaken in the remaining 81 patients (26%).

The median age at the start of treatment was 62yrs (range 6-97yrs) and 44 patients were over 80 yrs. 203 patients used H2RA's (65%) and 109 patients PPI's (35%). The median duration of treatment was 4 years (range 1-20yrs).

**Conclusion:** Scrutiny of GP records is likely to identify examples of the over-prescription of maintenance acid suppressive therapy and should be a prerequisite to assessing the impact of alternative therapeutic strategies in the management of GORD and peptic ulcer.

F258

#### ARTIFICIAL NEURAL NETWORK VS MULTIVARIATE LOGISTIC REGRESSION ANALYSIS FOR THE PREDICTION OF EARLY MORTALITY AFTER TIPSS FOR VARICEAL HAEMORRHAGE

<sup>1</sup>Jalan R, <sup>1</sup>Ala-Korpela M, <sup>2</sup>Hiltunen Y, <sup>3</sup>Redhead DN, <sup>4</sup>Stanley AJ, <sup>5</sup>Elton RA, <sup>6</sup>Bell JD, <sup>7</sup>Hayes PC. <sup>1</sup>NMR Unit, Hammersmith Hospital, London, <sup>2</sup>Raabe Institute of Computer Engineering, Finland; <sup>3</sup>Department of Radiology and <sup>4</sup>Medicine, and <sup>5</sup>Statistics Unit, Royal Infirmary Edinburgh, UK.

**Background and Aims:** TIPSS for variceal haemorrhage is followed by deterioration in liver function tests and early mortality in about 20-30% patients. The purpose of this study was to compare the performance of a trained Artificial neural network (ANN) with the model based upon multiple logistic regression (MLR).

**Methods:** Over a four-year-period 82 consecutive patients undergoing TIPSS for variceal haemorrhage were studied. They were divided into two groups. Group I (66 patients) comprised the patients that were used to train the ANNs and establish the model based upon the results MLR. Group II (16 patients) comprised the patients that were used in a blinded manner to assess the trained neural network and also the MLR.

**ANN:** A feed-forward fully-connected ANN with 10 hidden neurons (DynaMind™) was trained with the 25 clinical variables related to clinical or biological data obtained from patients in Group I (input) to predict early mortality (output).

**MLR:** Significant independent predictors (sodium; p < 0.001 and Pugh score p < 0.001) were combined using the formula:  $p = e^x / 1 + e^x$ , where p is the predicted probability of survival and  $x = 13.42 - 0.1429 X [\text{sodium}] + 0.445 X [\text{Pugh score}]$ .

This network and the MLR model were then applied to predict early mortality of patients in Group II.

**Results:** Sensitivity and specificity for predicting early mortality were 100% and 87.5% for the ANN and 25% and 93.8% for the MLR, respectively.

**Conclusions:** This study illustrates that ANN analysis can be useful in the prediction of early mortality before TIPSS is inserted, using routine clinical and biochemical parameters. Moving from the assessment of outcome by current methods towards ANN analysis will require similar comparisons, prospective evaluation and an open mind.

F260

#### H. PYLORI SEROLOGY IN THE INVESTIGATION OF DYSPEPSIA: A GENERAL PRACTICE BASED FOLLOW-UP STUDY. T.G.Reilly, D Stone, V Poxon, TSJ Elliott, RP Walt. University Department of Medicine, Queen Elizabeth Hospital Birmingham.

**Introduction:** H. pylori serology has been proposed as a screening tool to focus endoscopy resources in young dyspeptics. We offer ELISA Helicobacter serology to GPs prior to referral for endoscopy.

**Methods:** Patients identified from a serology database were followed by examination of GP records for data on investigation, diagnosis, and H. pylori eradication therapy (ET). Prescriptions for antisecretory treatment and number of GP consultations were the outcome measures.

**Results:** From 1262 patients whose blood was tested for H. pylori over 14 months we examined the GP records of 958. 220 patients whose serology was performed after investigation are excluded from this analysis. 738 had either no investigation (406) or investigation following serology (332):

Diagnosis n	Hp serology n (P=pos)	Had eradication therapy n (±SD)	
		Yes	No
406 none (no tests)	157 P	113 c=0.6 (±1.0) r=42 (±77)	44 c=0.42 (±0.63) r=39 (±66)
	249 N	5 c=0.96 (±0.65) r=33 (±20)	244 c=0.59 (±0.87) r=47 (±79)
269 non ulcers (after tests)	189 P	113 c=1.27 (±1.44) r=58 (±76)	76 c=0.83 (±0.94) r=60 (±92)
	80 N	1	79 c=1.09 (±1.18) r=69 (±97)
63 ulcers	60 P	56 c=0.50 (±0.82) r=46 (±88)	4
	3 N	1	2

Legend: c=mean consultations/pt/year, r=mean days prescribed treatment/pt/yr.

Of 406 seropositives 282 received ET, 118(29%) without further investigation. 124(43%) of those receiving ET had a further test to confirm success. In the 332 undergoing investigation there were 63 ulcers (3 seronegative, 2 <50 yrs.) and 2 gastric cancers (>50). Neither the differences between non-ulcer patients (NUP) who did and did not have ET nor those between seronegative and seropositive NUP were significant for either outcome measure. Seropositives who had no investigation had fewer prescriptions (p=0.03, Anova) and consultations (p<0.0001) than either seropositive or seronegative NUP but not ulcer patients. **Conclusions:** 29% of dyspeptics seropositive for Helicobacter received ET with no investigation. From our data, withholding endoscopy from seronegative patients under 50 would have missed 3.3% of ulcers. Treatment with ET was not beneficial in terms of lower consultation or prescription rates, in the absence of proven ulcer.