

Gut

An International Journal of
Gastroenterology and Hepatology

SEPTEMBER 2000 VOLUME 47 NUMBER 3

Leading articles

- 317 Pancreas divisum: opinio divisa *L Quest, M Lombard*
320 COX-1 and COX-2 products in the gut: therapeutic impact of COX-2 inhibitors *B J R Whittle*

Commentaries

- 326 Distension in non-consulting dyspeptics: a swell idea *J Tack*
327 Suicide and the not-so-innocent bystander *R J Aspinall, N R Lemoine*
328 Dissecting out cryptogenic liver disease *J G C Kingham*

Science @lert

- 330 CagA: a role at last *J C Atherton*

Motility and visceral sensation

- 332 Differences in gastric mechanosensory function after repeated ramp distensions in non-consulters with dyspepsia and healthy controls *G Holtmann, J Gschossmann, J Neufang-Hüber, G Gerken, N J Talley*
337 Functional assessment of β adrenoceptor subtypes in human colonic circular and longitudinal (taenia coli) smooth muscle *L Manara, T Croci, G Aureggi, F Guagnini, J-P Maffrand, G Le Fur, S Mukenge, G Ferla*

Cancer

- 343 Combined suicide and cytokine gene therapy for peritoneal carcinomatosis *C Lechanteur, P Delvenne, F Princen, M Lopez, G Fillet, J Gielen, M-P Merville, V Bours*
349 Role of fucosyltransferases in the association between apomucin and Lewis antigen expression in normal and malignant gastric epithelium *A López-Ferrer, C de Bolós, C Barranco, M Garrido, J Isern, I Carlstedt, C A Reis, J Torrado, F X Real*
357 Detection of Ki-ras gene point mutations in bile specimens for the differential diagnosis of malignant and benign biliary strictures *J-C Saurin, M-O Joly-Pharaboz, P Pernas, L Henry, T Ponchon, J-J Madjar*
362 Elevated levels of the pro-carcinogenic adduct, O⁶-methylguanine, in normal DNA from the cancer prone regions of the large bowel *A C Povey, C N Hall, A F Badawi, D P Cooper, P J O'Connor*

Small intestine

- 366 IgG₁ antiendomysium and IgG antitissue transglutaminase (anti-tTG) antibodies in coeliac patients with selective IgA deficiency *F Cataldo, D Lio, V Marino, A Picarelli, A Ventura, G R Corazza, the Working Groups on Celiac Disease of SIGEP and Club del Tenue*
370 Elevated plasma glucagon-like peptide 1 and 2 concentrations in ileum resected short bowel patients with a preserved colon *P B Jeppesen, B Hartmann, J Thulesen, B S Hansen, J J Holst, S S Poulsen, P B Mortensen*

Intestinal infection

- 377 Enterohaemorrhagic *Escherichia coli* O157:H7 target Peyer's patches in humans and cause attaching/effacing lesions in both human and bovine intestine *A D Phillips, S Navabpour, S Hicks, G Dougan, T Wallis, G Frankel*
- 382 Effect of vasoactive intestinal polypeptide (VIP) antagonism on rat jejunal fluid and electrolyte secretion induced by cholera and *Escherichia coli* enterotoxins *F H Mourad, C F Nassar*

Inflammatory bowel disease

- 387 Neurokinin-1 receptor expression in inflammatory bowel disease: molecular quantitation and localisation *T Goode, J O'Connell, P Anton, H Wong, J Reeve, G C O'Sullivan, J K Collins, F Shanahan*
- 397 Butyrate inhibits inflammatory responses through NF κ B inhibition: implications for Crohn's disease *J-P Segain, D Raingeard de la Bl ti re, A Bourreille, V Leray, N Gervois, C Rosales, L Ferrier, C Bonnet, H M Blottiere, J-P Galmiche*
- 404 A reproducible grading scale for histological assessment of inflammation in ulcerative colitis *K Geboes, R Riddell, A  st, B Jensfelt, T Persson, R L fberg*
- 410 Mucosal 5-aminosalicylic acid concentration inversely correlates with severity of colonic inflammation in patients with ulcerative colitis *G Frieri, R Giacomelli, M Pimpo, G Palumbo, A Passacantando, G Pantaleoni, R Caprilli*
- 415 Comparable expression of matrix metalloproteinases 1 and 2 in pouchitis and ulcerative colitis *A Stallmach, C C Chan, K-W Ecker, G Feifel, H Herbst, D Schuppan, M Zeitz*

Pancreatitis

- 423 Expression of interleukin 8 (IL-8) and substance P in human chronic pancreatitis *P Di Sebastiano, F F di Mola, C Di Febbo, G Baccante, E Porreca, P Innocenti, H Friess, M W B chler*

Liver disease

- C** 429 Pathological and virological findings in patients with persistent hypertransaminasaemia of unknown aetiology *C Berasain, M Bet s, A Panizo, J Ruiz, J I Herrero, M-P Civeira, J Prieto*

Case report

- 436 Topical tacrolimus may be effective in the treatment of oral and perineal Crohn's disease *D H Casson, M Eltumi, S Tomlin, J A Walker-Smith, S H Murch*

Occasional viewpoint

- 441 Effect on outcome of the lengthening waiting list for liver transplantation *R Ransford, B Gunson, D Mayer, J Neuberger, E Christensen*

Review

- 444 Quality of life measurement in gastrointestinal and liver disorders *M R Bargaonkar, E J Irvine*

Letters to the editor

- 455 Liver biopsy under ultrasound control *R J Aspinall*
- 455 Informed consent *J R Bennett*
- 456 Increased prevalence of methylenetetrahydrofolate reductase C677T variant in patients with IBD *J N Nielsen, T B Larsen, L Fredholm, I Brandslund, P Munkholm, H Hey*
- 457 **Book reviews**

Cover illustration: Scanning electron microscopy of human intestine in vitro organ culture sample showing presence of dome shaped lymphoid follicle. (From the paper by Phillips *et al*, pp 377-81.)