Prolonged measurement of lower oesophageal sphincter function in patients with intestinal metaplasia at the oesophagogastric junction

C Wolf, R Timmer, R Breumelhof, C A Seldenrijk, A J P M Smout

Abstract

Background and aims—It has been shown that gastro-oesophageal reflux plays a role in the pathogenesis of intestinal metaplasia (IM) limited to the oesophagogastric junction (OGJ), similar to the pathogenesis of IM in long segments of columnar lined oesophagus. The aim of this study was to examine lower oesophageal sphincter (LOS) function by means of prolonged recording in patients with IM limited to a normal appearing OGJ.

Patients and methods—Eighteen patients with IM at the OGJ (five females, 13 males; mean age 55.4 years) and 22 patients without IM (nine females, 13 males; mean age 53.9 years) underwent conventional stationary oesophageal manometry. Thereafter, seven hour water perfused manometry with simultaneous pH measurement (probe 5 cm proximal to the LOS) was performed. Swallowing was monitored with a pharyngeal sidehole and LOS pressure was recorded with a Dent sleeve. Patients were studied in the fasted state (three hours) and after a standardised meal (four hours). LOS pressure was analysed using customised software, and the incidence of reflux episodes (pH <4 for at least five seconds) and transient LOS relaxations (TLOSRS) were examined. TLOSRS were judged to be accompanied by reflux if a decrease of 1 pH unit occurred during relaxation.

Results—Patients with IM at the OGJ had a higher prevalence of postprandial acid reflux compared with patients without IM. No differences were observed in LOS pressure (pre- and postprandially) or in the prevalence of TLOSRS. However, in the postprandial phase, the rate of TLOSRS accompanied by acid reflux was increased in patients with IM.

Conclusion—Patients with IM at the OGJ have a higher prevalence of postprandial acid reflux. This is not associated with a higher prevalence of TLOSRS or a decreased LOS pressure but with a higher rate of TLOSRS accompanied by reflux.

(Gut 2001;49:354-358)

Keywords: intestinal metaplasia; gastro-oesophageal reflux; transient lower oesophageal sphincter relaxations; oesophagogastric junction

While many studies have examined the pathogenesis of traditional columnar lined oesophagus with intestinal metaplasia (IM) (Barrett's

oesophagus with a columnar lining >3 cm and biopsy proved IM), there are only a few studies which have focused on the pathogenesis of IM in short segments of columnar lining or IM confined to the area of the oesophagogastric junction (OGJ). IM harbours the risk of malignant transformation and adenocarcinomas have also been observed in short columnar lined segments.^{1,2} Research interest has focused on IM at or close to the OGJ because of the increasing incidence of adenocarcinomas in this area.^{3,4} It is important to know more about the pathogenesis of IM in this region as it might be possible to treat the underlying mechanism in order to prevent the neoplastic transformation or even to achieve regression of IM. In patients with short segments of columnar lining with IM, a higher incidence of acid gastro-oesophageal reflux has been observed.⁵⁻⁷ Even IM limited to an endoscopically normal appearing OGJ has been found to be associated with gastro-oesophageal reflux.⁸ There are no clear data concerning the mechanism underlying reflux in this patient group. A reduction in lower oesophageal sphincter (LOS) pressure in patients with IM has been described.⁸ However, this was measured with conventional stationary manometry, a technique that does not allow prolonged monitoring of LOS function in different digestive states. One important mechanism for gastro-oesophageal reflux is transient lower oesophageal sphincter relaxations (TLOSRS), characterised by an abrupt decrease in LOS pressure that is not associated with swallowing.⁹⁻¹¹ The increased prevalence of reflux in patients with IM at the OGJ could be due to an increased number of TLOSRS. Therefore, we wished to study the prevalence of TLOSRS and to monitor LOS function over a longer period of time, including the postprandial phase, in patients with IM at an endoscopically normal OGJ compared with patients without IM.

Materials and methods

STUDY GROUP

The study population consisted of two groups: patients with IM at the OGJ and patients without IM. Patients were recruited from a group

Abbreviations used in this paper: IM, intestinal metaplasia; OGJ, oesophagogastric junction; SCJ, squamocolumnar junction; LOS, lower oesophageal sphincter; TLOSRS, transient lower oesophageal sphincter relaxation; SAPLOSRS, swallow associated prolonged lower oesophageal sphincter relaxation; MSAPLOSRS, multiple swallow associated prolonged lower oesophageal sphincter relaxation; SANLOSRS, swallow associated normal lower oesophageal sphincter relaxation; OR, odds ratio.

Department of
Gastroenterology, St
Antonius Hospital
Nieuwegein, the
Netherlands
C Wolf
R Timmer

Department of
Gastroenterology,
Diaconessenhuis
Utrecht, the
Netherlands
R Breumelhof

Department of
Pathology, St Antonius
Hospital Nieuwegein,
the Netherlands
C A Seldenrijk

Gastrointestinal
Research Unit,
Departments of
Gastroenterology and
Surgery, University
Medical Centre,
Utrecht, the
Netherlands
A J P M Smout

Correspondence to:
C Wolf, Department of
Gastroenterology, St
Antonius Hospital,
Koekoekslaan 1, NL-3435
CM Nieuwegein, the
Netherlands. c.wolf@
cerberus.dimes.tudelft.nl

Accepted for publication
26 February 2001

who participated in a previous biopsy study performed by our group. In that study all patients had been endoscoped within the past six months and were found to have a normal appearing OGJ. During that endoscopy two forceps biopsies were obtained from immediately distal to the squamocolumnar junction (SCJ)—that is, within 0–5 mm of the SCJ. Additional biopsies were taken from the distal stomach, antrum, corpus, and “cardia”, the latter being defined 2 cm distal to the proximal stop of the gastric folds. All biopsies were stained with haematoxylin and eosin, alcian blue, and modified Giemsa. The diagnosis of IM was made according to histological findings. Patients were classified as IM positive if goblet cells intensively stained with alcian blue were identified. The IM positive group consisted of 18 patients (five female, 13 male) with a mean age of 55.4 years (SD 10.4); six had a hiatal hernia. The IM negative group consisted of 22 patients (nine female, 13 male; mean age 53.9 (14.9) years; six patients had a hiatal hernia). There were no significant differences between the two groups concerning the use of acid reducing medication, reflux symptoms, smoking habits, and prevalence of *Helicobacter pylori* infection.

MANOMETRIC AND pH RECORDING SETUP

After an overnight fast all patients first underwent a conventional stationary manometry. A polyvinyl catheter with three sideholes located 5 cm apart was introduced nasally. The catheter was connected to a low compliance pneumohydraulic pump and perfused with distilled water at a rate of 0.5 ml/min. A station pull through was performed to locate the LOS and then the oesophageal response to 10 wet swallows (5 ml) was tested. Thereafter the catheter was removed.

A micromanometric silicone catheter was introduced (Dentsleeve Pty Ltd, Parkside, Australia). This 11 lumen catheter had a total length of 160 cm, its external diameter was 3.5 mm, and the internal diameter of the individual channels was 0.4 mm, except for one centrally located infusion channel (1.9 mm). Four pharyngeal sideholes (located 24, 26, 28, and 30 cm above the proximal border of the sleeve) were used to monitor swallowing and three sideholes were located in the oesophageal body (4, 9, and 14 cm above the sleeve). A reverse perfused 6 cm long Dent sleeve was used for recording LOS pressure and a sidehole at the proximal border of the sleeve was used for its positioning. Gastric pressure was recorded from a sidehole located 2 cm below the distal border of the sleeve. The catheter was connected to external pressure transducers (Abbott, Chicago, Illinois, USA) which were mounted on a belt.¹² A customised water pump (water reservoir 360 ml, kept at a pressure of 1 bar) developed by the Department of Biomedical Engineering of the University Medical Centre Utrecht, was used to perfuse the resistors via a silicone manifold (Dentsleeve Pty Ltd). The resistors reduced the water flow to a perfusion rate of 0.08 ml/min.¹³ The pressure transducers were connected to a portable

data logger (MMS, Enschede, the Netherlands) that digitised the data at a rate of 4 or 8 Hz (oesophageal and pharyngeal recordings, respectively). Patients wore the belt around their waist so that the transducers were located directly distal to the lower end of the ribcage. After correct positioning of the manometry catheter a glass pH probe was introduced through the same nostril and placed with its tip located 5 cm above the oral border of the LOS. This probe was also connected to the data logger.

STUDY PROTOCOL

Patients were studied in a semi recumbent position. After a short accommodation period, data recording was started. Subjects were studied preprandially for three hours and for another four hours after a standardised solid meal (Menu Nasi, Iglo, the Netherlands; total energy content 1575 kJ, 45 g carbohydrate, 12.5 g fat, 22.5 g protein). The study was approved by the ethics committee of St Antonius Hospital.

ANALYSIS OF PRESSURE RECORDINGS

Data from the portable logger were downloaded to a PC and visualised with commercially available software (MMS, Enschede, the Netherlands). LOS pressure was evaluated using a computer program that was developed by one of the investigators (AJPMs). This program constructs the end expiratory LOS pressure curve and calculates mean end expiratory LOS pressure values (with respect to gastric baseline pressure) for the individual study periods. It has been validated by comparing its results with manual analysis by several experienced investigators (MA van Herwaarden, unpublished data).

The prevalence of TLOSRS was analysed for the pre- and postprandial periods. TLOSRS were defined using established criteria: abrupt decrease (>0.4 kPa/3 s) in end expiratory LOS pressure (nadir pressure of <0.4 kPa) not initiated by a swallow (no signal in pharyngeal channels four seconds before or two seconds after the onset of relaxation) with a relaxation duration of at least five seconds.^{14 15} During this analysis the pH channel was hidden from the observer. The prevalence of reflux episodes, defined as an abrupt decrease in pH <4 for at least five seconds, was examined for the pre- and postprandial study periods. The number of reflux episodes and number of TLOSRS were calculated per hour. The underlying mechanism of reflux events was classified as being one of the following: (1) TLOSRS, (2) normal swallow associated relaxation (SANLOSRS, pharyngeal peak within four seconds before or two seconds after the onset of relaxation, duration of relaxation shorter than 10 s), (3) swallow associated prolonged LOS relaxation (SAPLOSRS, pharyngeal peak within four seconds before or two seconds after the onset of relaxation, duration of relaxation equal to or longer than 10 s), (4) multiple swallow associated LOS relaxation (MSAPLOSRS, multiple pharyngeal peaks within four seconds before or during relaxation), (5) low LOS pressure

Figure 1 Prevalence of reflux episodes in patients with (+) and without (-) intestinal metaplasia (IM) in the pre- and postprandial periods (mean (SEM)). * $p=0.042$, IM positive versus IM negative group.

(undetectable LOS pressure for at least 30 s before reflux), or (6) straining (sharp and brief positive elevations in gastric and oesophageal pressure prior to pH decrease). Reflux was judged to have accompanied a TLOSRS if a decrease in oesophageal pH greater than 1 pH unit was observed during the relaxation.

STATISTICAL ANALYSIS

Data are expressed as mean (SEM) or percentages. Statistical analysis was performed with SPSS software. The χ^2 test, Mann-Whitney U test, and t test were used where appropriate.

Results

As shown in fig 1, the prevalence of reflux episodes was higher in the postprandial than in the fasting period (0.7/hour and 1.8/hour, respectively; odds ratio (OR) 2.08; 95% confidence interval (CI) 1.33–3.26). Patients with IM at the OGJ had more reflux episodes per hour in the postprandial state compared with patients without IM (2.25/hour *v* 1.35/hour, respectively; OR 1.65; 95% CI 1.01–2.74). Similar findings were noted for the percentage of time with pH below 4 in the postprandial phase (1.9% *v* 4.38%; OR 1.42; 95% CI 1.03–1.95) (table 1). There was no difference between the two groups in the preprandial phase. End expiratory LOS pressures did not differ between the two groups neither preprandially or in the postprandial phase (table 1). If both groups were analysed together a trend towards a lower LOS pressure in the postprandial state was observed (1.12 and 0.96 kPa; ns). For the entire group, the prevalence of TLOSRS (calculated per hour) was higher in the postprandial period than in the preprandial period (1.8/hour and 2.9/hour, respectively; OR 2.4; 95% CI 1.48–3.88). The prevalence of TLOSRS pre- and postprandially was not different between IM positive and IM negative

Figure 2 Prevalence of transient lower oesophageal sphincter relaxations (TLOSRS) and TLOSRS associated with reflux in patients with (+) and without (-) intestinal metaplasia (IM) (mean (SEM)). * $p=0.05$, TLOSRS associated with reflux, IM positive versus IM negative group.

patients. However, a higher rate of TLOSRS accompanied by reflux (postprandially) was observed in patients with IM at the OGJ. In these patients 66.9% of TLOSRS were associated with reflux compared with 46.5% in patients without IM (OR 7.41; 95% CI 1.02–60.58) (fig 2). Pre- and postprandially there was no difference in the prevalence of SAPLOSRS or MSAPLOSRS, or in the rate of SAPLOSRS or MSAPLOSRS associated with reflux between patients with or without IM.

Preprandially, patients with IM had a total number of 49 reflux episodes compared with 34 reflux episodes in IM negative patients. As shown in fig 3, the mechanisms through which these reflux episodes occurred were not different between groups (IM+ 57.1% TLOSRS, 4.1% SAPLOSRS, 4.1% MSAPLOSRS, 22.5% SANLOSRS, 0% low LOS pressure, 12.2% straining; IM- 58.8% TLOSRS, 2.9% SAPLOSRS, 0% MSAPLOSRS, 23.5% SANLOSRS, 2.9% low LOS pressure, 11.7% straining). In the postprandial period a total of 135 reflux episodes occurred in the IM positive patients (57.1% TLOSRS, 5.9% SAPLOSRS, 5.2% MSAPLOSRS, 13.3% SANLOSRS, 6.7% low LOS pressure, 11.9% straining) and a total of 97 episodes in the IM negative patients (70.1% TLOSRS, 6.2% SAPLOSRS, 4.1% MSAPLOSRS, 12.4% SANLOSRS, 2.1% low LOS pressure, 5.2% straining).

Discussion

Barrett's oesophagus is characterised by the presence of columnar lining with IM in the distal oesophagus. It is known that this condition is the consequence of longstanding gastro-oesophageal reflux disease caused by disturbances in oesophageal motility and LOS dysfunction.^{16–18} Patients with Barrett's oesophagus have a 30–40-fold increased risk of developing oesophageal malignancies.¹⁹ IM in the columnar lining of Barrett's oesophagus represents a precursor lesion for oesophageal adenocarcinoma.² As the prevalence of carcinomas in the area of the OGJ has been increasing over the last 10 years, IM in that region and its pathogenesis have gained research interest.^{3,4} In an earlier study (unpublished data) we confirmed the results of Öberg *et al* in a similar patient population.⁸ The results of our and

Table 1 Reflux parameters and lower oesophageal sphincter (LOS) characteristics in patients with and without intestinal metaplasia (IM) at the oesophagogastric junction

	IM + (n=18)	IM - (n=22)	OR (95% CI)
% time pH <4			
Preprandial period (%)	1.75 (0.46)	1.15 (0.22)	1.33 (0.84–2.10)
Postprandial period (%)	4.38 (0.99)	1.90 (0.34)	1.42 (1.03–1.95)
End expiratory LOS pressure			
Preprandial period (kPa)	1.10 (0.09)	1.18 (0.14)	0.76 (0.24–2.48)
Postprandial period (kPa)	0.94 (0.09)	1.09 (0.13)	0.55 (0.14–2.10)

Data are mean (SEM) with odds ratios (OR) and 95% confidence interval (CI).

Figure 3 Distribution of reflux mechanisms (mean (SEM), expressed as percentage of reflux episodes) in patients with (+) and without (-) intestinal metaplasia (IM), pre- and postprandially. No significant differences were found. LOS, lower oesophageal sphincter; TLOS, transient lower oesophageal sphincter relaxation; SAPLOS, swallow associated prolonged lower oesophageal sphincter relaxation; MSAPLOS, multiple swallow associated prolonged lower oesophageal sphincter relaxation; SANLOS, swallow associated normal lower oesophageal sphincter relaxation.

their studies suggest a causative role for gastro-oesophageal reflux disease, similar to long segment traditional Barrett's oesophagus. However, in contrast with Öberg *et al*, we could not find a clear reduction in LOS competence, as measured with short term conventional manometry. In the interpretation of results obtained from stationary oesophageal manometry, one has to bear in mind that the LOS is not a constant high pressure zone but a dynamic structure that undergoes physiological changes in the circadian rhythm in healthy subjects as well as in patients with gastro-oesophageal reflux disease.^{9 10 20} Therefore, LOS measurements that monitor LOS function over a longer period of time provide more information than short term recordings. Prolonged recordings of LOS function can be obtained with a miniature Dent sleeve assembly. Oesophageal body and sphincter measurements with this device have been shown to be reproducible and stable.²¹ The feasibility of the recording system that we used (transducers on belt, portable perfusion pump) has been demonstrated previously.^{12 13}

To our knowledge this is the first study that has measured LOS function over a prolonged period of time in patients with IM at the OGJ and in a comparable group of patients without IM. This method not only allows measurement of LOS pressure but it is also possible to assess LOS relaxation. Low LOS pressure and TLOSs have been shown to be important

factors in the pathogenesis of gastro-oesophageal reflux. Simultaneous pH recording in the distal oesophagus permits identification of the underlying mechanisms of observed reflux episodes.

Results obtained with this new method support the earlier findings that gastro-oesophageal reflux plays a role, as we found a higher prevalence of reflux episodes in patients with IM compared with patients without IM. This higher prevalence was observed in the postprandial state only. We did not find a statistically significant difference in the prevalence of LOS pressure or TLOSs between the two groups, either in the preprandial or postprandial state. However, we observed that in the postprandial period the proportion of TLOSs associated with reflux was higher in patients with IM. These findings indicate that the increased oesophageal acid exposure in IM positive patients is caused by factors other than the "traditional" reflux mechanisms such as low LOS pressure and a high prevalence of TLOSs. The prevalence of a hiatal hernia was not different between the two groups and therefore it seems unlikely that this played a role in the current study. Apparently, some other factor is involved in patients with IM at the OGJ which causes relaxations of the LOS to lead more readily to reflux than in patients without IM. The most likely explanation for the observed phenomenon is that in patients with IM, the volume, acidity, or distribution of

gastric contents favours reflux more than in IM negative patients. This could be due to a slower rate of gastric emptying, a higher rate of gastric acid secretion, altered intragastric content distribution, or differences in postprandial gastric motility.²²⁻²⁶

In conclusion, using a method for prolonged monitoring of LOS function with combined pH measurement, this study provided evidence that IM at an endoscopically normal OGJ was associated with acid gastro-oesophageal reflux. LOS pressure was not lower in patients with IM compared with patients without IM. Furthermore, a higher prevalence of TLOSRS was not observed in the IM positive patient group. However, these patients had a higher prevalence of postprandial acid gastro-oesophageal reflux; the percentage of TLOSRS associated with acid reflux was increased. Elucidation of mechanisms underlying this requires further studies.

We would like to thank the endoscopy and pathology departments for their support in the study. This study was supported by Aventis BV, the Netherlands.

- Schnell TG, Sontag SJ, Chejfec G. Adenocarcinomas arising in tongues or short segments of Barrett's esophagus. *Dig Dis Sci* 1992;37:137-43.
- Haggitt RC. Barrett's esophagus, dysplasia, and adenocarcinoma. *Hum Pathol* 1994;25:982-93.
- Blot WJ, Devesa SS, Kneller RW, et al. Rising incidence of adenocarcinoma of the esophagus and gastric cardia. *JAMA* 1991;265:1287-9.
- Pera M, Cameron AJ, Trastek VF, et al. Increasing incidence of adenocarcinoma of the esophagus and esophagogastric junction. *Gastroenterology* 1993;104:510-13.
- Loughney T, Maydonovitch CL, Wong RKH. Esophageal manometry and ambulatory 24-hour pH monitoring in patients with short and long segment Barrett's esophagus. *Am J Gastroenterol* 1998;93:916-19.
- Clark GWB, Ireland AP, Peters JH, et al. Short-segment Barrett's esophagus: a prevalent complication of gastro-oesophageal reflux disease with malignant potential. *J Gastrointest Surg* 1997;1:113-22.
- Öberg S, Ritter MP, Crookes PF, et al. Gastroesophageal reflux disease and mucosal injury with emphasis on short-segment Barrett's esophagus and duodenogastroesophageal reflux. *J Gastrointest Surg* 1998;2:547-53.
- Öberg S, Peters JH, DeMeester TR, et al. Inflammation and specialized intestinal metaplasia of cardiac mucosa is a manifestation of gastroesophageal reflux disease. *Ann Surg* 1997;226:522-32.
- Dent J, Dodds WJ, Friedman RH, et al. Mechanism of gastroesophageal reflux in recumbent asymptomatic human subjects. *J Clin Invest* 1980;65:256-67.
- Dodds WJ, Dent J, Hogan WJ, et al. Mechanisms of gastroesophageal reflux in patients with reflux esophagitis. *N Engl J Med* 1982;307:1547-52.
- Mittal RK, Holloway RH, Penagini R, et al. Transient lower esophageal sphincter relaxation. *Gastroenterology* 1995;109:601-10.
- Samsom M, Smout AJ, Hebbard G, et al. A novel portable perfused manometric system for recording of small bowel motility. *Neurogastroenterol Motil* 1998;10:139-48.
- van Herwaarden MA, Samsom M, Smout AJPM. 24-h ambulatory LOS pressure recording using a portable water-perfused manometric system and fully automated signal analysis. *Neurogastroenterol Motil* 1998;10:A72.
- Holloway RH, Kocyan P, Dent P. Provocation of transient lower esophageal sphincter relaxations by meals in patients with symptomatic gastroesophageal reflux. *Dig Dis Sci* 1991;36:1034-9.
- Holloway RF, Penagini R, Ireland AC. Criteria for objective definition of transient lower esophageal sphincter relaxation. *Am J Physiol* 1995;268:G128-33.
- Stein JH, Hoeft S, DeMeester TR. Functional foregut abnormalities in Barrett's esophagus. *J Thorac Cardiovasc Surg* 1993;105:107-11.
- Singh P, Taylor RH, Colin-Jones DG. Esophageal motor dysfunction and acid exposure in reflux esophagitis are more severe if Barrett's metaplasia is present. *Am J Gastroenterol* 1994;89:349-56.
- Coenraad M, Masclee AAM, Straathof JWA, et al. Is Barrett's esophagus characterized by more pronounced acid reflux than severe esophagitis? *Am J Gastroenterol* 1998;93:1068-72.
- Reid BJ. Barrett's esophagus and esophageal adenocarcinoma. *Gastroenterol Clin North Am* 1991;20:817-34.
- Schoeman MN, Tippet MD, Akkermans LMA, et al. Mechanisms of gastroesophageal reflux in ambulant healthy human subjects. *Gastroenterology* 1995;108:83-91.
- Crowell MD, Schettler-Duncan VA, Yu SS. Manometric evaluation of esophageal function utilizing a miniature Dent sleeve: within and between session reproducibility in healthy controls. *Gastroenterology* 1999;115:G4257.
- McCallum RW, Berkowitz DM, Lerner E. Gastric emptying in patients with gastroesophageal reflux. *Gastroenterology* 1981;80:285-91.
- Cadiot G, Bruhat A, Rigaud D, et al. Multivariate analysis of pathophysiological factors in reflux oesophagitis. *Gut* 1997;40:167-74.
- Vaezi MF, Richter JE. Role of acid and duodenogastric esophageal reflux in GER. *Gastroenterology* 1996;111:1192-9.
- Shay SS, Conwell DL, Mehindru V, et al. The effect of posture on gastroesophageal reflux event frequency and composition during fasting. *Am J Gastroenterol* 1996;91:54-60.
- Penagini R, Hebbard G, Horowitz M, et al. Motor function of the proximal stomach and visceral perception in gastro-oesophageal reflux disease. *Gut* 1998;42:251-7.