

IRRITABLE BOWEL SYNDROME

Morphological characterisation of Crohn's disease fistulae

F Bataille, F Klebl, P Rümmele, J Schroeder, S Farkas, P-J Wild, A Fürst, F Hofstädter, J Schölmerich, H Herfarth, G Rogler

Gut 2004;53:1314-1321. doi: 10.1136/gut.2003.038208

See end of article for authors' affiliations

Correspondence to: Dr F Bataille, Institute of Pathology, University of Regensburg, 93042 Regensburg, Germany; frauke.bataille@klinik.uni-regensburg.de

Revised version received 10 March 2004
Accepted for publication 16 March 2004

Background: Fistulae are a common complication in up to 35% of all patients with Crohn's disease. Their therapy is difficult and frequently unsatisfactory. To date, no histological comparison of Crohn's disease fistulae with non-inflammatory bowel disease fistulae has been performed. In addition, Crohn's disease fistulae have not been well characterised morphologically.

Methods: Eighty four fistulae from Crohn's disease patients were compared with 13 fistulae from controls. Haematoxylin-eosin staining, electron microscopy, and immunohistochemistry for panCytokeratin (epithelial cells), CD20 (B cells), CD45R0 (T cells), and CD68 (macrophages) were performed according to standard techniques. In addition, histopathological findings were compared with clinical and laboratory data.

Results: In 31.0% of controls and 27.4% of Crohn's disease specimens, fistulae had a lining of flattened intestinal epithelium without goblet cells or, in the case of cutaneous/perianal disease, narrow squamous epithelium. Non-epithelialised fistulae were covered by a thin layer of (myo)fibroblasts, focally forming a new basement membrane, as demonstrated by electron microscopy. All fistulae were surrounded by granulation tissue. Crohn's disease fistulae presented with central infiltration by CD45R0⁺ T cells, followed by a small band of CD68⁺ macrophages and dense accumulation of CD20⁺ B cells. In contrast, in controls, there was dense infiltration by CD68⁺ macrophages with only few CD20⁺ B cells and CD45R0⁺ T lymphocytes.

Conclusions: Fistulae in Crohn's disease differ markedly from non-Crohn's disease fistulae with regard to their cellular composition. The presence of an epithelial lining in a subgroup of fistulae may be important for the therapeutic approach and healing process.

Fistulae are a frequent problem in patients with Crohn's disease (CD). The reported incidence ranges from 17% up to 50%.¹⁻⁵ This may be due to a centre bias as fistulising CD is seen more frequently in centres for inflammatory bowel disease (IBD). However, population based studies also show a high incidence of fistulae in CD.⁶⁻⁷

In the perianal area, formation of fistulae can be easily identified clinically and are sometimes associated with abscesses.⁸ Enterointestinal fistulae on the other hand are difficult to diagnose and can be the cause of malabsorption in CD patients but they are also less frequent.⁸ In a recent population based study, fistulae were reported in up to 35% of patients with CD, with perianal fistulae occurring in approximately 20% of cases.⁶⁻⁹ The cumulative incidence of fistulising CD in the study was 33% after 10 years and increased up to 50% after 20 years, making it a major clinical problem of disease management. Two thirds of those patients however had only a single fistula episode: 34% of these patients had recurrent fistulae. Recurrent fistulae occurred less frequently in patients who received maintenance therapy with an immunosuppressive agent.

In approximately 10% of patients suffering from CD, perianal fistulisation is the initial presentation of disease.¹ Fistulae may precede the onset of intestinal disease by several years. Patients with colonic disease, especially with involvement of the rectum, have a higher incidence of perianal fistulae. Internal fistulae have been classified clinically into two types: those which form an internal connection between two bowel layers or segments, and those that occur between the intestine and other organs, such as enterovesical or abdominal wall fistulae.¹⁰⁻¹¹

The pathogenesis of fistulae formation is still unknown. It is clear that fistulae in the perianal area do not arise from the

small intestine but develop locally.¹¹⁻¹² However, their histological features have never been investigated in detail.

Fistulae are an important problem in the management of CD. A number of patients will develop an abscess, which either drains spontaneously or must be drained surgically. Surgical excision of fistulae has not proved to be very successful.¹³⁻¹⁵ Seton drainage seems to be more helpful but the frequency of complete healing of fistulae is low.¹⁶⁻¹⁷ Medical therapy is based on treatment with antibiotics or immunosuppressants such as azathioprine, cyclosporin, tacrolimus, or infliximab.^{1-3 8 18 19} Permanent closure of fistulae can only be achieved in approximately 20-30% of patients. Therefore, close cooperation between surgeons and gastroenterologists is usually recommended.

To investigate the pathophysiology and develop better therapeutic approaches for fistulising CD, it is necessary to understand the histological features of Crohn's fistulae. However, a detailed histological investigation of Crohn's fistulae has not been published previously. Therefore, in this study, histological differences between CD and non-CD fistulae were characterised.

MATERIALS AND METHODS

Patients

In this retrospective study, 97 fistula specimens from 78 patients were examined. The series was obtained from unselected cases at the Institute of Pathology, University of Regensburg. Specimens had been surgically removed between August 1993 and May 2003. Eighty four fistula

Abbreviations: CD, Crohn's disease; IBD, inflammatory bowel disease; CRP, C reactive protein; H&E, haematoxylin-eosin; DAB, diaminobenzidine chromogen

Table 1 Clinical data on patients in the control group

No	Sex	Age at surgery (y)	Fistula location	Clinical diagnosis
1	M	49	Skin, cervical	Recurring cervical fistulae after squamous cell carcinoma of the tongue
2	M	40	Perianal	Recurring perianal fistulae, exclusion of CD
3	F	79	Skin, abdominal	Fistula of the skin, abdominal
4	M	34	Perianal	Sinus peronidialis with recurring perianal fistulae
5	F	51	Stomach	Fistula of the stomach after ulcus ventriculi
6	M	63	Oral cavity	Dental abscess with recurring fistulae of the oral cavity
7	F	42	Perianal	Fistula after periproctitic abscess, since 1994 until 2003 no CD, 2003 breast cancer
8	M	42	Perianal	Fistula after thoracal abscess
		47	Skin, thoracal	
		47	Skin, thoracal	
9	F	73	Vaginal	Vaginal fistula after radiation of vaginal squamous cell carcinoma
10	M	41	Perianal	Sinus pilonidalis with recurring perianal fistulae
11	F	21	Perianal	Sinus pilonidalis with perianal fistula

specimens were derived from 67 CD patients and 13 fistulae from 11 controls with no IBD. The diagnosis of CD was based on established clinical, endoscopic, and radiological parameters.^{20, 21} In addition to fistula characterisation, histopathological, electron microscopic, and immunohistochemical characteristics were compared between both groups of patients.

Patients with CD comprised 33 males and 34 females; mean age was 36.9 years (range 13–75). Six controls were male and five were female; mean age was 48.4 years (range 21–79). Fifty two fistulae (53.6%) were diagnosed in males and 45 (46.4%) in females. Clinical data are presented in tables 1–3. Most CD fistulae were located in the small or large intestine whereas most control fistulae were perianal or cutaneous fistulae. Eleven patients with CD and two control patients had more than one fistula. Details on fistula distribution are shown in tables 1–3.

Specimen preparation

Tissue specimens were fixed in 4% buffered formalin for at least 24 hours and embedded in paraffin. Sections of approximately 2–3 μ m thickness were cut from tissue blocks and stained with haematoxylin and eosin, according to standard protocols.

Histology

Two gastrointestinal pathologists (FB, PR) performed the histopathological examinations without knowledge of the clinical data. Specimens were examined in random order. The configuration of inflammation (acute/chronic), granulation tissue, capillarity, deepness of fistula, formation of abscesses, and presence of lining epithelium were evaluated separately.

Electron microscopy

Small tissue (primarily fixed in buffered 4% formalin for LM) samples were deparaffinised in xylol, hydrated in descending ethanol, additionally fixed in 0.1 M cacodylate buffered 4% glutaraldehyde (overnight), postfixed in 1% osmium tetroxide (two hours) at pH 7.3, dehydrated in graded ethanol, and embedded in the EmBed-812 epoxy resin (all reagents from Science Services, Munich, Germany). After 48 hours of heat polymerisation at 60°C, semithin (0.8 μ m) sections were cut, stained with toluidine blue, and after selection of appropriate areas of interest the Epon block was trimmed for ultrathin sectioning. Ultrathin (80 nm) sections were cut with a diamond knife on a Reichert Ultracut-S ultramicrotome and double contrasted with aqueous 2% uranyl acetate and lead citrate solutions for 10 minutes each. Sections were

examined in a LEO912AB electron microscope operating at 80 kV. Details of the investigated fistulae are shown in table 4.

Immunohistochemistry

Immunohistochemical studies for expression of CD20, CD45RO, CD68, and panCytokeratin were performed on 21 fistulae of patients with CD and 13 controls using an avidin-biotin peroxidase method with diaminobenzidine (DAB) chromogen. After antigen retrieval (microwave treatment of formalin fixed paraffin embedded 2–3 μ m tissue sections for 40 minutes at 240 W in citrate buffer, pH 6.0) immunohistochemistry was carried out in a NEXES immunostainer (Ventana Medical System, Tucson, Arizona, USA) according to the manufacturer's instructions. As primary antibody, mouse monoclonal antibodies were used at a dilution of 1:50 for panCytokeratin (panCK; Coulter-Immunotech Diagnostics, Hamburg, Germany), 1:100 for CD20 (clone L26, DakoCytomation, Hamburg, Germany) and CD45RO (clone UCHL T cell; Dako), and at a dilution of 1:200 for CD68 (clone KPI; Dako). After incubation for 24 minutes at 37°C, slides were rinsed in phosphate buffered saline and incubated with the secondary antibody (rabbit-antimouse, 1:500 dilution in phosphate buffered saline; Ventana Medical System) for two hours at room temperature. Antibody binding was visualised with 0.05% DAB (Ventana Medical System) and 0.01% hydrogen peroxide. The material was rinsed in phosphate buffered saline and counterstained with haematoxylin. Cells were regarded as expressing panCytokeratin, CD20, CD45RO, and CD68 when there was cytoplasmic staining.

Clinical and serological parameters

Clinical and laboratory data were retrieved by chart review. Data collected included duration between diagnosis and resection, medication at time of surgery, previous medical therapy, last C reactive protein (CRP) level before surgery, and white blood cell count before resection.

Statistics

Associations between histopathological features, clinicopathological parameters, clinical data, and patient sex and age were analysed statistically in univariate fashion by χ^2 testing, Mann-Whitney U test, or Fisher's exact test, where appropriate. Statistics were computed using SPSS software (SPSS, Inc.; Chicago, Illinois, USA). All statistical tests were performed two sided with a p value <0.05 considered statistically significant.

Table 2 Clinical data on Crohn's disease fistulae. In Crohn's disease, every fistula is counted as a single event

No	Year of birth	Sex	Year of surgery	Location of fistula	Year of diagnosis of CD
1	1959	M	1994	Sigma	1982
			1994	Sigma	
			1994	Sigma	
			1997	Colon	
2	1973	F	1998	Colon	1997
			1997	Perianal	
3	1964	F	1994	Rectum	1993
4	1970	F	2003	Ileum	1996
			2003	Ileum	
5	1967	M	2000	Ileum	1998
6	1948	M	1999	Colon	1982
7	1963	F	1998	Ileum	1998
8	1972	M	1994	Ileum	1989
9	1967	M	1997	Ileum	1987
10	1954	M	2003	Perianal	Not known
11	1975	M	1997	Ileum	1997
12	1951	M	1999	Ileum	1988
13	1939	M	1994	Ileum	1994
14	1940	F	2003	Ileum	Not known
15	1953	F	2000	Ileum	1985
16	1974	M	1995	Ileum	1995
			1998	Ileum	
17	1970	F	1998	Colon	1990
			2002	Colon	
18	1964	M	2002	Sigma	2002
19	1972	F	1993	Ileum	1982
20	1956	F	1993	Ileum	1993
			1993	Perianal	
21	1925	M	2000	Ileum	2000
22	1953	F	1995	Rectum	1980
23	1954	M	2001	Ileum	1992
24	1953	M	1997	Ileum	1978
25	1950	F	1997	Colon	1973
26	1969	M	2000	Sigma	1999
27	1947	F	2000	Perianal	1973
28	1984	M	2002	Colon	1998
29	1985	M	2002	Colon	2001
30	1962	F	1999	Perianal	1993
31	1972	F	2000	Ileum	1988
32	1957	F	1994	Ileum	1986
			1996	Ileum	
33	1971	M	1993	Ileum	1988
34	1954	F	1998	Colon	1997
			2003	Ileum	
35	1969	M	2002	Ileum	1995
36	1934	F	2002	Ileum	2002
37	1952	F	1998	Colon	1986
38	1951	F	1994	Ileum	1993
39	1937	F	1997	Colon	1994
40	1961	F	1998	Ileum	1998
41	1943	F	2001	Ileum	2001
42	1975	F	2002	Colon	1994
43	1968	M	1994	Ileum	1991
			1994	Colon	
44	1963	M	2003	Colon	1991
			1999	Ileum	
45	1934	M	1995	Ileum	1994
			2002	Ileum	
46	1968	M	2001	Ileum	1995
			2001	Colon	
47	1969	M	2000	Ileum	2000
48	1988	M	2001	Ileum	2001
49	1963	M	1994	Ileum	1994
50	1959	M	1999	Perianal	1991
51	1971	M	1996	Ileum	1989
52	1973	F	1999	Ileum	1999
53	1975	F	2002	Ileum	2001
54	1961	M	2001	Ileum	1997
55	1949	F	1993	Perianal	1993
56	1976	F	2002	Perianal	1996
57	1956	M	1999	Ileum	1999
58	1978	M	2003	Ileum	2003
59	1964	F	2001	Ileum	1987
60	1938	M	2001	Ileum	2001
61	1969	F	1993	Ileum	1993
62	1957	F	1992	Colon	1992
63	1982	M	1997	Colon	1996
			1999	Ileum	
64	1967	M	1998	Ileum	1997
65	1963	F	2000	Ileum	1996
			2000	Colon	
66	1967	F	2000	Ileum	1997
67	1946	F	2002	Colon	1985
			2002	Ileum	

Table 3 Clinical data on Crohn's disease patients

Patients	
Sex (M/F)	34/33
Age at surgery (y)	36.7 (4.0)
Time between diagnosis of fistula and surgery (months)	4.0 (37.9)
Prior medical treatment for fistular disease	
Antibiotics	10 (11.9%)
Azathioprine	5 (6%)
Oral 5-ASA derivatives	6 (7.1%)
Systemic glucocorticoids	12 (14.3%)
Topical 5-ASA derivatives	1 (1.2%)
Cyclophosphamide	1 (1.2%)
Cyclosporin	1 (1.2%)
Infliximab	1 (1.2%)
Mycophenolat mofetile	1 (1.2%)
Present medication	
Antibiotics	29 (34.5%)
Azathioprine	10 (11.9%)
Oral 5-ASA derivatives	27 (32.1%)
Systemic glucocorticoids	42 (50%)
Topical 5-ASA derivatives	2 (2.4%)
Topical glucocorticoids	0 (0%)
Cyclosporin	0 (0%)
<i>E coli</i> Nissle	1 (1.2%)
Infliximab	0 (0%)
Methotrexate	1 (1.2%)
Location of fistula	
Ileum	50 (59.5%)
Colon	26 (31%)
Perianal	8 (9.5%)
Concomitant complication	
None	27 (32%)
Abscess	26 (31%)
Conglomerate tumour	24 (28.6%)
Peritonitis	7 (8.3%)

RESULTS

Histology

Histologically, all studied fistulae had a central fissure in common, which branched and penetrated through lamina propria and muscularis mucosae deep into the underlying tissue. All fistulae were lined by granulation tissue with conspicuous pale plump histiocytes and a dense network of tender capillaries. According to the duration of the process, the lumen was filled up by nuclear debris, sometimes erythrocytes, and non-specific acute (neutrophils; n = 83) or chronic (lymphocytes; n = 14) inflammation (fig 1). In 2.0% of fistula specimens there was no acute inflammation, in 12.4% there was mild, in 28.9% moderate, and in 56.7% severe acute inflammation. Independent of the inflammatory infiltrate, 31.0% of control fistulae and 27.4% of CD fistulae had a lining epithelium central in the fistulising inflammation (fig 2). Depending on the location, the lining epithelium consisted of flattened epithelium of the small intestine or colon without goblet cells, or narrow squamous epithelium in cutaneous or perianal fistulae (fig 2). There was no association between epithelialisation of fistulae and deepness of infiltration or extent of inflammation.




Figure 1 Histopathological picture of colonic Crohn's disease with a fistula. (A) At low magnification, the mucosa is destroyed, ulcerated, and replaced by granulation tissue, including a dense capillary network. The surface is covered by fibrin and neutrophils. In the right half, there is a deep fistula which infiltrates the muscularis propria (arrows) (haematoxylin-eosin (H&E), ×16). (B) The central fissure (arrowheads) is branching deep into the underlying tissue, lined by neutrophils, granulation tissue with numerous capillaries (arrows), histiocytes, and lymphocytes (H&E, ×50). (C) At the surface of the central fissure, there is a small layer of histiocytes and neutrophils (arrows). Lining epithelium is not detectable (H&E, ×400).

For patients with more than one fistula, six CD patients (55%) and two patients in the control group had fistulae with different histological compositions. CD patients had

Table 4 Epithelialised and non-epithelialised fistulae of Crohn's disease (CD) patients and controls by electron microscopy

Sex	Age at surgery (y)	Location of fistula	Epithelialised/ non-epithelialised	Disease
Male	31	Sigma	Epithelialised	CD
Male	33	Terminal ileum	Epithelialised	CD
Male	71	Perianal	Non-epithelialised	CD
Female	42	Perianal	Epithelialised	Non-CD
Female	73	Sigma	Epithelialised	Non-CD
Male	34	Perianal	Non-epithelialised	Non-CD




Figure 2 Representative histological pictures of epithelialised and non-epithelialised fistulae in Crohn's disease (CD) (A, C) and controls (B, D). Depending on the location, the lining epithelium was similar to epithelium of the small intestine, colon, or squamous epithelium. (A) Colonic fistula (arrows) without lining epithelium in a CD patient (haematoxylin-eosin (H&E), $\times 16$ (inlay $\times 100$)). (B) Perianal fistula (arrows) without lining epithelium in a control patient (H&E, $\times 16$ (inlay $\times 50$)). (C) Epithelialised fistula (arrows) of the colon in a CD patient (H&E, $\times 16$ (inlay $\times 100$)). (D) Epithelialised fistula (arrows) of the small intestine in a control patient (H&E, $\times 16$ (inlay $\times 200$)).

synchronous ($n = 7$) or metachronic ($n = 10$) fistulae in the small intestine and colon. In four of six patients with CD, only fistulae of the small intestine had a lining epithelium. Synchronous fistulae in the control group were located perianally or cutaneously, each in a single patient. In both patients, only one of the two fistulae was epithelialised.

Electron microscopy

Fistulae were located in the small intestine, colon, or perianal region (table 4). Independent of the underlying disease, the lining epithelium consisted of flattened epithelium of the small or large intestine, or narrowed squamous cells. Cells showed tight junctions and a basement membrane. Non-epithelialised CD fistulae were lined in sections with myofibroblasts, with gap junctions to each other. In some areas, a new basement membrane, localised between the myofibroblasts and the underlying granulation tissue, was visible. Myofibroblasts were connected by fibronexus to the newly generated basement membrane. Adjacent sections with disordered myofibroblasts were visible. They showed no gap junctions and the underlining basement membrane was fragmented. In the control group, the basement membrane was more continuous, not only in sections, as shown in CD fistulae. Details are shown in fig 3.

Immunohistochemistry

Immunohistochemically, there was a different cellular composition within the groups. In patients with CD, the interior wall of the fistulae was infiltrated by CD45RO positive T cells, followed by a small band of CD68 positive macrophages (in 19/21 fistulae (90.5%)). At the outer fistulae wall, there was a dense infiltrate of CD20 positive B cells (in 16/21 fistulae (76.2%)) (fig 4A, C, E, G). In fistulae of the control group, there was intense infiltration by CD68 positive macrophages throughout the whole wall (in 10/13 fistulae (76.9%)). Only a few CD20 positive B lymphocytes could be

identified (in 8/13 fistulae (61.5%)). CD45RO positive T lymphocytes were present in the outer two thirds of fistulae wall (fig 4B, D, F, H). This distribution was independent of fistulae localisation and deepness of infiltration. Some (4/13), and mostly perianal fistulae of the control group, had an isolated infiltrate of only monocytes, surrounded by fibrosis, thereby resembling advanced chronified fistulae.

Clinical and serological parameters

There was no significant association between histopathological findings and sex ($p = 0.650$), age ($p = 0.942$), fistulae location ($p = 0.902$), medical treatment at the time of surgery ($p = 0.738$), CRP level ($p = 0.407$), or white blood cell count ($p = 0.918$) prior to surgical intervention.

DISCUSSION

Fistulae are a complication of CD which is feared by patients and physicians because of their associated morbidity and difficult therapy. The specific pathogenesis of fistulae in CD, and also in non-CD, is not known. For perianal fistulae, two alternative hypotheses have been developed. The first suggests that fistulae begin as deep penetrating ulcers in the anus or rectum.²² Over time, faeces accumulate in these ulcers, and the intraluminal pressure during defecation forces the faecal matter into subcutaneous tissue, extending the ulcer and creating a fistula. The second differs in that the fistula results from an anal gland abscess that serves as the point of origin for a fistula tract.²²

The natural history of CD fistulae is one of exacerbations and long episodes of active drainage.¹⁻⁷ Despite the obvious clinical importance of this complication, little is known about its pathogenesis and histopathological composition.

Epithelialisation of fistulae was found in approximately one third of CD and control patients. Oberhuber *et al* described herniated mucosa within the muscularis propria or the subserosal fatty tissue in seven of 27 cases of fistulising CD.²³




Figure 3 Electron micrographs of epithelialised and non-epithelialised fistulae of Crohn's disease patients and controls. (A–D) Re-epithelialised fistulae of the small intestine (A: semithin section, toluidine blue, $\times 200$; B: electron micrograph) and colon (C: semithin section, toluidine blue, $\times 200$; D: electron micrograph). Histomorphologically, there is no similarity to the original lining epithelium of the small intestine (A: arrow) or colon (C: arrow) but by electron microscopy there are typical microvilli (B: white arrows) and goblet cells (D: arrows). (E–G) Central fissure of a fistula lined by myofibroblasts (E: arrow). There is the typical appearance of myofibroblasts (F: asterisks): a cell membrane with multiple caveolae (F: white arrows), a rough endoplasmic reticulum and Golgi complex, and the cytoplasm containing bundles of microfilaments with focal densities (G: arrowhead). Myofibroblasts are connected to each other by adherens and gap junctions (F: black arrows) and are in contact with a newly formed thin basement membrane by fibronexus (G: white arrows). (H–J) Central fissure of a fistula lined by epithelium (H: asterisks) and myofibroblasts (H: arrow). The myofibroblasts rest irregularly on the surface of the fissure of the fistula (I: arrows), there are no generated adherens or gap junctions, and the underlining newly formed basement membrane (J: white arrows) is not continuous.

In our series, there was no herniated mucosa but the epithelium was similar to that of the surface, most likely mimicking a process of regeneration. Dysplasias were not detectable. Whether fistulae trigger malignant transformation and cancer is controversial. Early onset of disease, disease duration, chronic (pan-)colitis with high inflammatory activity, and the persistence of chronic fistulae and stenosis have been reported as risk factors for the development of malignancy.²⁴ In contrast, Wong *et al* suggested that fistula associated adenocarcinomas represent anal gland carcinomas.²⁵ Whether epithelialisation of fistula tracks with introduction of epithelial cells into deeper layers of the intestinal wall is associated with a higher risk of cancer is an intriguing question which cannot yet be answered. In addition, we do not know whether epithelialisation occurs in every fistula at the same stage or if epithelialised and non-epithelialised fistulae need different therapeutic approaches. It may be speculated that epithelialised fistulae represent a form of failed but stable restitution which is difficult to treat solely with medical therapy. Furthermore, if epithelialisation implies an increased cancer risk and there is a sequence from non-epithelialised to epithelialised fistulae, early aggressive medical therapy aimed at healing the fistula (for example, use of antibiotics, azathioprine,²⁶ or infliximab²⁷) may reduce this risk. Certainly, this is speculative at present and more data are needed.

Whereas on histology, fistulae of both patient groups shared some similarities, they differed markedly immunohistochemically. The most conspicuous finding was accumulation of CD45R0 positive T cells in the inner layer and dense infiltration of CD20 positive B cells surrounding the fistulae in patients with CD, as opposed to predominance of CD68 positive histiocytes in the granulation tissue of control fistulae. Similar to inflammation in CD itself, the role of various cell types is likely to differ in the development and evolution of fistulae. Activation of T cells and monocytes/macrophages is regarded as an important factor in the pathogenesis of IBD.^{28–31} Memory T cells play a role in initiating or perpetuating the inflammatory process. They may be primed to foreign micro-organisms or targeted to self antigens.^{32–33} Under normal circumstances, regulatory mucosal T cells maintain a state of controlled inflammation and suppress pathological inflammation.³⁴

Treatment of CD has focused on neutralisation of pro-inflammatory mediators or administration of anti-inflammatory molecules but also on reduction of proinflammatory cells within the intestine.³⁵ Medical treatment of CD reduces the mucosal content of proinflammatory cytokine producing T cells. All of our CD patients had received medication. This may have influenced the cellular composition of CD fistulae and may explain why there were less histiocytic cells compared with control fistulae. Nevertheless, we did not




Figure 4 Immunohistochemical picture of Crohn's disease (CD) (A, C, E, G) and control (B, D, F, H) fistula. Presented are representative pictures of the distribution of CD20 positive B cells (A and B: arrows), CD68 positive macrophages (C and D: arrows), CD45RO positive T cells (E and F: arrows), and panCytokeratin positive epithelial cells (G: arrows) (overview $\times 16$, inlay $\times 400$).

observe any association between the type of medication and the histological picture of CD fistulae. Therefore, it seems possible that the mechanisms involved in the pathogenesis of CD and non-CD fistulae are different and that the various cell types contribute differently.

It is likely that epithelial restitution in vivo involves a complex interaction between epithelial cells and the underlying lamina propria cells, which occurs via the basement membrane or pores within it.^{36, 37} Intestinal myofibroblasts are located adjacent to the basement membrane and in close proximity to the basal surface of epithelial cells.^{38, 39} In this position, myofibroblasts are capable of regulating a number

of epithelial functions such as epithelial restitution,⁴⁰ barrier function,⁴¹ and electrolyte transport.⁴² In contrast with the control group, CD fistulae had sections with disordered myofibroblasts and fragmented underlying basement membrane.

Myofibroblasts appear to be key cells in the events of tissue repair but CD myofibroblasts seem to differ from control myofibroblasts⁴³ and this may account for induction of some dissimilarities observed in CD and control fistulae.

In the case of deep ulcers, subepithelial tissue containing interstitial substance, blood vessels, nerves, and mesenchymal cells must be reconstituted in addition to the epithelium.

If the basement membrane has been destroyed by noxious stimuli (for example, inflammation), epithelial cells and mesenchymal elements form a new basement membrane.⁴⁴ Our electron microscopy findings suggest that myofibroblasts also play a role in this repair process, being able to form a basement membrane even in the absence of epithelial cells.

Rapid migration of epithelial cells over a denuded basement membrane is an important response to injury. Whether this migration also occurs with time in non-epithelialised fistulae or whether these represent a different type of fistula cannot be determined from our data. However, based on our findings here, we hypothesise that the development and maturation of fistulae follows several successive steps: (1) initially, a stimulus in combination with a proinflammatory response leads to deep tissue destruction which may be perpetuated in the base of the fistula by intruding luminal antigens/bacteria; (2) myofibroblasts migrate or divide in the vicinity of the lesion; (3) they become organised by development of gap junctions and adherens and, trying to protect the body from the contents within the intestinal lumen, form a basement membrane; and (4) epithelial cells migrate from the surrounding of the inner lumen of the fistula onto the freshly formed basement membrane, therefore leading to an epithelialised fistula. According to this hypothesis, fistula formation would be caused by failure to close a tissue defect by a linear basement membrane. Perpetuation of the inflammation at the base of the fistula may prevent the migration and arrangement of myofibroblasts which might be a necessary step to achieve closure of the fistula. Differences in the cellular constitution of the inflammatory infiltrate between CD and normal fistulae suggest that either the response to a proinflammatory stimulus or the stimulus itself is different. However, it is necessary to further understand the molecular mechanisms and cellular interplay which are responsible for the formation of fistulae and differences between CD and non-CD fistulae.

ACKNOWLEDGEMENTS

This work was supported by the BMBF Competence network "Inflammatory Bowel Disease". We thank Sabine Troppmann and Doris Gaag for outstanding technical assistance.

Authors' affiliations

F Bataille, P Rümmele, J Schroeder, P-J Wild, F Hofstädter, Institute of Pathology, University of Regensburg, Regensburg, Germany
F Klebl, J Schölmerich, H Herfarth, G Rogler, Department of Internal Medicine I, University of Regensburg, Regensburg, Germany
S Farkas, A Fürst, Department of Surgery, University of Regensburg, Regensburg, Germany

REFERENCES

- Hellers G, Bergstrand O, Ewerth S, et al. Occurrence and outcome after primary treatment of anal fistulae in Crohn's disease. *Gut* 1980;**21**:525-7.
- Michelassi F, Stella M, Balestracci T, et al. Incidence, diagnosis, and treatment of enteric and colorectal fistulae in patients with Crohn's disease. *Ann Surg* 1993;**218**:660-6.
- Solomon MJ. Fistulae and abscesses in symptomatic perianal Crohn's disease. *Int J Colorectal Dis* 1996;**11**:222-6.
- Allan A, Keighley MR. Management of perianal Crohn's disease. *World J Surg* 1988;**12**:198-202.
- Bell SJ, Williams AB, Wiesel P, et al. The clinical course of fistulating Crohn's disease. *Aliment Pharmacol Ther* 2003;**17**:1145-51.
- Loftus EV jr, Schoenfeld P, Sandborn WJ. The epidemiology and natural history of Crohn's disease in population-based patient cohorts from North America: a systematic review. *Aliment Pharmacol Ther* 2002;**16**:51-60.
- Loftus EV jr, Silverstein MD, Sandborn WJ, et al. Crohn's disease in Olmsted County, Minnesota, 1940-1993: incidence, prevalence, and survival. *Gastroenterology* 1998;**114**:1161-8.
- Givel JC, Hawker P, Allan R, et al. Entero-enteric fistula complicating Crohn's disease. *J Clin Gastroenterol* 1983;**5**:321-3.
- Schwartz DA, Loftus EV jr, Tremaine WJ, et al. The natural history of fistulizing Crohn's disease in Olmsted County, Minnesota. *Gastroenterology* 2002;**122**:875-80.
- D'Haens G. Medical management of major internal fistulae in Crohn's disease. *Inflamm Bowel Dis* 2000;**6**:244-5.
- Present DH. Urinary tract fistulas in Crohn's disease: surgery versus medical therapy. *Am J Gastroenterol* 2002;**97**:2165-7.
- Present DH. Crohn's fistula: current concepts in management. *Gastroenterology* 2003;**124**:1629-35.
- O'Hanlon DM, O'Connell PR. Complex fistulae in Crohn's disease. *J Am Coll Surg* 2002;**194**:87.
- Hoffmann JC, Zeitz M. Treatment of Crohn's disease. *Hepatogastroenterology* 2000;**47**:90-100.
- Francois Y, Vignal J, Descos L. Outcome of perianal fistulae in Crohn's disease—value of Hughes' pathogenic classification. *Int J Colorectal Dis* 1993;**8**:39-41.
- Takesue Y, Ohge H, Yokoyama T, et al. Long-term results of seton drainage on complex anal fistulae in patients with Crohn's disease. *J Gastroenterol* 2002;**37**:912-15.
- Williams JG, Macleod CA, Rothenberger DA, et al. Seton treatment of high anal fistulae. *Br J Surg* 1991;**78**:1159-61.
- Bell SJ, Kamm MA. Review article: the clinical role of anti-TNFalpha antibody treatment in Crohn's disease. *Aliment Pharmacol Ther* 2000;**14**:501-14.
- Present DH. Review article: the efficacy of infliximab in Crohn's disease— healing of fistulae. *Aliment Pharmacol Ther* 1999;**13**(suppl 4):23-8.
- Gasche C, Scholmerich J, Brynskov J, et al. A simple classification of Crohn's disease: report of the Working Party for the World Congresses of Gastroenterology, Vienna 1998. *Inflamm Bowel Dis* 2000;**6**:8-15.
- Modigliani R, Mary JY, Simon JF, et al. Clinical, biological, and endoscopic picture of attacks of Crohn's disease. Evolution on prednisolone. Groupe d'Etude Therapeutique des Affections Inflammatoires Digestives. *Gastroenterology* 1990;**98**:811-18.
- Hughes LE. Surgical pathology and management of anorectal Crohn's disease. *J R Soc Med* 1978;**71**:644-51.
- Oberhuber G, Stangl PC, Vogelsang H, et al. Significant association of strictures and internal fistula formation in Crohn's disease. *Virchows Arch* 2000;**437**:293-7.
- Winkler R, Wittmer A, Heusermann U. Cancer and Crohn's disease. *Z Gastroenterol* 2002;**40**:569-76.
- Wong NA, Shirazi T, Hamer-Hodges DW, et al. Adenocarcinoma arising within a Crohn's-related anorectal fistula: a form of anal gland carcinoma? *Histopathology* 2002;**40**:302-4.
- Dejaco C, Harrer M, Waldhoer T, et al. Antibiotics and azathioprine for the treatment of perianal fistulas in Crohn's disease. *Aliment Pharmacol Ther* 2003;**18**:1113-20.
- Present DH, Rutgeerts P, Targan S, et al. Infliximab for the treatment of fistulas in patients with Crohn's disease. *N Engl J Med* 1999;**340**:1398-405.
- Saubermann LJ, Probert CS, Christ AD, et al. Evidence of T cell receptor beta-chain patterns in inflammatory and noninflammatory bowel disease states. *Am J Physiol* 1999;**276**:G613-21.
- Schreiber S. Monocytes or T cells in Crohn's disease: does IL-16 allow both to play at that game? *Gut* 2001;**49**:747-8.
- Zareie M, Singh PK, Irvine EJ, et al. Monocyte/macrophage activation by normal bacteria and bacterial products: implications for altered epithelial function in Crohn's disease. *Am J Pathol* 2001;**158**:1101-9.
- Boirivant M, Marini M, Di Felice G, et al. Lamina propria T cells in Crohn's disease and other gastrointestinal inflammation show defective CD2 pathway-induced apoptosis. *Gastroenterology* 1999;**116**:557-65.
- Picarella D, Hurlbut P, Rottman J, et al. Monoclonal antibodies specific for beta 7 integrin and mucosal addressin cell adhesion molecule-1 (MAdCAM-1) reduce inflammation in the colon of scid mice reconstituted with CD45Rbhigh CD4+ T cells. *J Immunol* 1997;**158**:2099-106.
- Ludviksson BR, Strober W, Nishikomori R, et al. Administration of mAb against alpha E beta 7 prevents and ameliorates immunization-induced colitis in IL-2-/- mice. *J Immunol* 1999;**162**:4975-82.
- Shanahan F. Crohn's disease. *Lancet* 2002;**359**:62-9.
- Caprilli R, Viscido A, Guagnozzi D. Review article: biological agents in the treatment of Crohn's disease. *Aliment Pharmacol Ther* 2002;**16**:1579-90.
- Mahida YR, Galvin AM, Gray T, et al. Migration of human intestinal lamina propria lymphocytes, macrophages and eosinophils following the loss of surface epithelial cells. *Clin Exp Immunol* 1997;**109**:377-86.
- McAlindon ME, Gray T, Galvin A, et al. Differential lamina propria cell migration via basement membrane pores of inflammatory bowel disease mucosa. *Gastroenterology* 1998;**115**:841-8.
- Kaye GI, Pascal RR, Lane N. The colonic pericyptal fibroblast sheath: replication, migration, and cytodifferentiation of a mesenchymal cell system in adult tissue. 3. Replication and differentiation in human hyperplastic and adenomatous polyps. *Gastroenterology* 1971;**60**:515-36.
- Powell DW, Miffin RC, Valentich JD, et al. Myofibroblasts. II. Intestinal subepithelial myofibroblasts. *Am J Physiol* 1999;**277**:C183-201.
- McKaig BC, Makh SS, Hawkey CJ, et al. Normal human colonic subepithelial myofibroblasts enhance epithelial migration (restitution) via TGF-beta3. *Am J Physiol* 1999;**276**:G1087-93.
- Beltinger J, McKaig BC, Makh S, et al. Human colonic subepithelial myofibroblasts modulate transepithelial resistance and secretory response. *Am J Physiol* 1999;**277**:C271-9.
- Berschneider HM, Powell DW. Fibroblasts modulate intestinal secretory responses to inflammatory mediators. *J Clin Invest* 1992;**89**:484-9.
- McKaig BC, Hughes K, Tighe PJ, et al. Differential expression of TGF-beta isoforms by normal and inflammatory bowel disease intestinal myofibroblasts. *Am J Physiol Cell Physiol* 2002;**282**:C172-82.
- Wallace JL, Granger DN. The cellular and molecular basis of gastric mucosal defense. *Faseb J* 1996;**10**:731-40.