

INFLAMMATORY BOWEL DISEASE

Patients with active inflammatory bowel disease lack immature peripheral blood plasmacytoid and myeloid dendritic cells

D C Baumgart, D Metzke, J Schmitz, A Scheffold, A Sturm, B Wiedenmann, A U Dignass

Gut 2005;54:228–236. doi: 10.1136/gut.2004.040360

See end of article for authors' affiliations

Correspondence to:
Dr D C Baumgart, Charité
Medical Centre-Virchow
Hospital, Medical School
of the Humboldt-
University, Department of
Medicine, Division of
Hepatology &
Gastroenterology,
D-13344 Berlin, Germany;
daniel.baumgart@
charite.de

Revision received
21 June 2004
Accepted for publication
22 June 2004

Background: Breakdown of tolerance against the commensal microflora is believed to be a major factor in the pathogenesis of inflammatory bowel disease (IBD). Dendritic cells (DC) have been implicated in this process in various animal models, but data on human DC in IBD are very limited.

Aim: To characterise plasmacytoid DC (PDC) and myeloid DC (MDC) in patients with active versus inactive IBD and healthy controls.

Patients and Methods: Peripheral blood was obtained from 106 patients (Crohn's disease (CD) n = 49, ulcerative colitis (UC) n = 57) and healthy controls (n = 19). Disease activity was scored using the modified Truelove Witts (MTWSI) for UC and the Harvey Bradshaw severity indices (HBSI) for CD. Four colour flow cytometric analysis was used to identify, enumerate, and phenotype DC. DC from patients with acute flare ups and healthy controls were cultured and stimulated with CpG ODN 2006 or lipopolysaccharide (LPS).

Results: IBD patients in remission (PDC UC, 0.39%; CD, 0.35%; MDC-1 UC, 0.23%; CD, 0.22% of PBMC) have slightly lower numbers of circulating DC compared with healthy controls (PDC 0.41%, MDC-1 0.25% of PBMC). In acute flare ups IBD patients experience a significant drop of DC (PDC UC, 0.04%; CD, 0.11%; MDC-1 UC, 0.11%; CD, 0.14% of PBMC) that correlates with disease activity (correlation coefficients: PDC MTWSI, 0.93; HBSI, 0.79; MDC-1 MTWSI, 0.75; HBSI, 0.81). Moreover, both express $\alpha 4\beta 7$ integrin and display an immature phenotype. Freshly isolated PDC and MDC-1 from untreated flaring IBD patients express higher baseline levels of CD86 which increases further in culture and upon stimulation compared with healthy controls.

Conclusion: IBD patients lack immature blood DC during flare ups which possibly migrate to the gut. An aberrant response to microbial surrogate stimuli suggests a disturbed interaction with commensals.

Dendritic cells (DC) are the longest known professional antigen presenting cells and were first described by Langerhans in 1868 at Charité in Berlin.¹ They are located at mucosal interfaces with the outside world, and migrate and patrol the organism ubiquitously via the blood stream. Dendritic cells control several aspects of the immune response including type (Th1, Th2, Th3, or Tr) and homing of antigen specific effector cells. Dendritic cells are thought to be involved in maintenance of oral tolerance depending on their location, number, and maturational state.^{2–3}

The expression of molecular pattern recognition receptors such as toll-like receptors (TLRs) on dendritic cells suggests a close interaction with microbes and microbial products, which in turn determines their influence on the respective immune response—that is, immunity versus tolerance.^{4–5} Several lines of evidence, particularly from animal and in vitro studies, suggest that dendritic cell function is critical for intestinal immune regulation—for example, colonic lamina propria dendritic cells display an activated phenotype; T cell proliferation in mucosal dendritic cell aggregates suggests direct involvement in T cell activation/silencing; the number of activated (CD134L positive) dendritic cells dramatically increases in mesenteric lymph nodes of a T cell transfer model of colitis and can be abrogated by CD134L antibody treatment; dendritic cells were shown to directly sample bacterial through the epithelial layer in an in vitro model; and dendritic cells from Peyer's patches induce high levels of $\alpha 4\beta 7$ integrin in T cells, thereby likely mediating gut tropism of T cells.^{6–21} Thus, dendritic cells are likely to be involved in the balance between immunity and non-responsiveness against the commensal microflora. Breakdown of oral

tolerance against the commensal microflora is believed to be a key factor in the pathogenesis of inflammatory bowel disease (IBD).^{22–24}

To date, the data on human DC in IBD are very limited. Due to their low frequency and the absence of a specific antibody it was very difficult to identify, isolate, and enrich them. Modern flow cytometry allowing staining with multiple fluorochromes simultaneously, employment of complex gating strategies, and the discovery of new dendritic cell specific antibodies such as BDCA-1 to 4 provide new tools to study this cell population.^{25–29}

Here we present the first comprehensive analysis of all three peripheral blood DC populations known to date in patients with IBD and healthy controls.

MATERIALS AND METHODS

Patients and controls

Human peripheral blood was obtained from 106 patients with IBD (Crohn's disease (CD) n = 49, ulcerative colitis (UC) n = 57) seen at our IBD clinic and 19 controls (total n = 125). Table 1 summarises the demographic data of patients and healthy controls. Samples were preferably analysed from patients off immunomodulators or prior to adding them to control flare ups; however some patients were receiving IBD

Abbreviations: CD, Crohn's disease; DC, dendritic cell(s); HBSI, Harvey Bradshaw severity index; IBD, inflammatory bowel disease; LPS, lipopolysaccharide; MDC, myeloid dendritic cells; MTWSI, modified Truelove Witts severity index; PDC, plasmacytoid dendritic cells; PBDC, peripheral blood dendritic cells; PBMC, peripheral blood mononuclear cells; PBS, phosphate buffered saline; UC, ulcerative colitis




Figure 1 Inflammatory bowel disease medications summary. Bars show percentages.

related drugs. Figure 1 summarises the medication data. All patients gave informed consent to the study. The study protocol was approved by the institutional review board.

Scoring of disease activity

Two widely accepted scores, the modified Truelove Witts severity index (MTWSI) for UC and the Harvey Bradshaw severity index (HBSI) for CD were used to quantitate disease activity.^{30–32} Both scores and modifications of it have been validated in numerous clinical trials (tables 2 and 3).^{33–34} UC patients who scored ≥ 10 on the MTWSI and CD patients who scored ≥ 7 on the HBSI were classified to have active disease (flare up). The average MTWSI for flaring UC patients was 16 (range 10–21) and the average HBSI for flaring CD patients 14 (range 11–17), respectively.

Preparation of peripheral blood mononuclear cells

Peripheral blood mononuclear cells (PBMC) were isolated using a protocol published previously.³⁵ Briefly, 10 ml of freshly drawn blood treated with NH_4 heparin was diluted in phosphate buffered saline (PBS) 1:2 and well mixed. 20 ml of diluted blood were then carefully layered over 18 ml of Ficoll Paque (Pharmacia, Freiburg, Germany). The tubes were centrifuged without brakes at 2200 rpm for 20 minutes at 20°C. The interphase containing mononuclear cells was transferred into a new Falcon tube. Cells were washed twice in $10\times$ their volume PBS and spun for six minutes at 1700 rpm at 4°C. The final pellet was resuspended in FACS

staining buffer (PBS with 20% FCS and 1% NaN_3 , pH 7.4) and counted using a Neubauer chamber. Only vital cells were counted using Trypan blue exclusion. Viability was greater than 98%. 1×10^5 – 1×10^6 cells were resuspended in 100 μl FACS staining buffer prior to staining with the respective fluorochrome labelled antibodies on ice.

FACS staining

Three colour or four colour flow cytometric analysis was used to identify and enumerate different populations of peripheral blood dendritic cells (PBDC) as described previously.³⁶ Briefly, before staining all cell preparations were first incubated with human Ig (Beriglobin, Behring, Marburg, Germany). 100 μl of each sample were pipetted in 96 V

Table 1 Demographic data of patients and controls

	(n)	%	Mean (SD)	Min	Max
Number	125	100			
Ethnicity					
White	120	96			
Arabic	5	4			
Crohn's disease	49	39			
Male	12				
Female	37				
Age (years)			40 (11)	19	63
Disease duration (years)			14 (11)	2	45
Age at diagnosis (years)			27 (10)	10	50
Ulcerative colitis	57	46			
Male	26				
Female	31				
Age (years)			40 (16)	21	74
Disease duration (years)			10 (6)	2	28
Age at diagnosis (years)			33 (14)	14	55
Healthy controls	19	15			
Male	8				
Female	11				
Age (years)			37 (13)	28	46

Table 2 Calculation of the modified Truelove Witts severity index (MTWSI) for ulcerative colitis

Category	Score
A Diarrhoea	0 = 1–2 BM/day 1 = 3–4 BM/day 2 = 5–6 BM/day 3 = 7–9 BM/day 4 = 10 or more BM/day
B Nocturnal diarrhoea/ Early AM awakening for BM	0 = No 1 = Yes
C Bloody stools	0 = None 1 = Occasionally with BM 2 = 50% of BM 3 = With every BM
D Fecal incontinence/ soiling	0 = No 1 = Yes
E Abdominal pain/cramping	0 = None 1 = Mild; aware, but tolerable 2 = Moderate; interferes with usual activities 3 = Severe; incapacitating 0 = Excellent
F General wellbeing	1 = Very Good 2 = Good 3 = Fair 4 = Poor 5 = Terrible
G Antidiarrhoeals/ narcotics	0 = No 1 = Yes
H Abdominal tenderness	0 = None 1 = Mild to moderate; localised 2 = Mild to moderate; diffuse 3 = Severe or rebound tenderness

BM, bowel movement.

Table 3 Calculation of the Harvey Bradshaw severity index (HBSI) for Crohn's disease

Category	Score
A General wellbeing	0 = Very well 1 = Slightly below par 2 = Poor 3 = Very poor 4 = Terrible
B Abdominal pain	0 = None 1 = Mild 2 = Moderate 3 = Severe
C Number of liquid stools per day	Number of stools
D Abdominal mass	0 = None 1 = Dubious 3 = Definite 4 = Definite and tender
E Complications	Score 1 for each of: Arthralgia Uveitis Erythema nodosum Pyoderma gangrenosum Aphthous ulcers Anal fissure New fistula Abscess

bottom well plates (Falcon, BD Biosciences, Heidelberg, Germany) and spun at 2000 rpm for two minutes at 4°C. Cells were resuspended in 50 µl FACS buffer and placed on ice. The appropriate amount of antibody was added and the cells were incubated for 30 minutes on ice in the dark. After washing them three times in FACS buffer they were analysed immediately on a FACS Calibur (BD Biosciences) flow cytometer.

Antibodies

Antibodies were obtained from commercial sources as fluorochrome and/or biotin conjugates. Streptavidin FITC, streptavidin PE, and streptavidin APC were purchased from Caltag, San Diego, CA, USA and used to detect reactivity of biotinylated primary antibodies. Light (forward and side) scatter gates were set to include all viable cells. Fluorochrome conjugated isotype matched control antibodies against mouse IgM, IgG1, and IgG2b (BD Biosciences, Heidelberg, Germany) as well as mouse IgG2a (Caltag, San Diego, CA, USA) were used to control for non-specific reactivity and anti-CD3 and anti-CD19 as fluorochrome compensation controls, respectively.

Identification of plasmacytoid dendritic cells

Plasmacytoid dendritic cells (PDC) were identified using double staining for both BDCA-2 (Miltenyi Biotec, Bergisch Gladbach, Germany) and CD123 (BD Biosciences, Heidelberg, Germany).

Identification of myeloid dendritic cells (MDC-1 and MDC-2)

As there is some cross reactivity with monocytes, granulocytes, and B cells when using the BDCA-1 and CD-11c antibodies to identify MDC-1 myeloid dendritic cells, these populations were removed from the analysis by using exclusion gates on anti-CD14 and anti-CD19 positive cells (BD Biosciences) prior to double staining with CD11c (BD Biosciences) and BDCA-1 (Miltenyi Biotec). Identification of MDC-2 required only the exclusion of granulocytes prior to staining with the respective antibody. MDC-2 were identified using BDCA-3 (Miltenyi Biotec) and HLA-DR (BD

Biosciences) counterstaining. For some experiments a newly developed dendritic cell enumeration kit using BDCA-1, 2, and 3 antibodies manufactured by Miltenyi Biotec was used to assess the numbers of dendritic cells in blood. Cells were prepared according to a protocol supplied by the manufacturer. Side by side experiments demonstrated similar, reproducible results when compared with individual staining (data not shown).

Assessment of maturation and activation markers

Anti-CD1a, anti-CD40, anti-CD83, anti-CD80, and anti-CD86 as known markers of activated/mature DC were purchased from BD Biosciences. Staining was performed after gating on PDC, MDC-1, or MDC-2, respectively.

Assessment of migration markers

Anti-CD62L, anti-CD49d ($\alpha 4$ integrin), and anti- $\beta 7$ integrin were purchased from BD Biosciences. Staining was performed after gating on PDC, MDC-1, or MDC-2, respectively.

Isolation of plasmacytoid and myeloid dendritic cells

Plasmacytoid dendritic cells (PDC) were isolated using magnetic cell separation using the BDCA-4 and myeloid dendritic cells (MDC-1) using BDCA-1 (CD1c) antibodies and microbeads (Miltenyi Biotec) from PBMC as described previously.³⁷ As a subset of peripheral blood B cells also expresses CD1c, cells were depleted using CD19 microbeads prior to enrichment of CD1c myeloid DC. The purity of the isolated DC population was checked with FACS and samples with less than 90% PDC or MDC-1 were not used for culture and stimulation experiments.

Culture and stimulation of plasmacytoid and myeloid dendritic cells

PDC were cultured and stimulated as described previously.^{38–40} Briefly, 15 000 freshly isolated PDC were resuspended in 150 µl RPMI plus L-glutamine, medium with 1% penicillin/streptomycin, 1% sodium pyruvate, and 10% FCS (all from Invitrogen, Karlsruhe, Germany) and with IL-3 at 10 ng/ml (R&D Systems, Wiesbaden, Germany) (to prevent apoptosis) and incubated for 24 hours at 37°C and 5% CO₂. MDC-1 were cultured in the same manner except for the IL-3 supplementation.

The sequence of the CpG B (ODN 2006) used to stimulate PDC at a concentration of 2 µg/ml in the present study was 5'-TCG TCG TTT TGT CGT TTT GTC GTT-3'. Phosphorothioate ODN 2006 was synthesised at TIB Molbiol, Berlin, Germany. MDC-1 were stimulated with 5 µg/ml lipopolysaccharide (LPS) (Sigma, Taufkirchen, Germany).

Flow cytometry

Before running the samples on a flow cytometer, propidium iodide (PI) (1:100 of a 1 mg/ml stock, Sigma, USA) was added to label dead cells. Flow cytometry was performed on a FACS Calibur (BD Biosciences) four colour flow cytometer. Data were analysed using Cell Quest (BD Biosciences) software. Advanced gating strategies were used to exclude dead (PI positive) cells and debris.

Statistical analysis

Statistical analysis including linear regression to assess correlation between dendritic cell counts, and established disease activity indices was performed using SPSS Science version 11.5 (SPSS Inc, Chicago, IL, USA) software. Curve fit was expressed as R-Square with 95% mean and individual prediction intervals. The correlation coefficient (CC) was calculated by extracting the square root from r^2 .

RESULTS

IBD patients in remission have slightly lower numbers of circulating PBDC (MDC < PDC) compared with healthy controls. In acute flare ups IBD patients experience a significant drop of both PDC and MDC.

The fraction of PDC in UC patients in remission was calculated to be on average 0.39% of vital PBMC and dropped to 0.04% in acute flare ups (figs 2A, B, and 3A). The MDC-1 fraction in UC patients in remission was calculated to be 0.23% of vital PBMC and dropped to 0.11% in acute flare ups (figs 2C, D, and 4A).

For CD patients in remission a mean PDC fraction of 0.35% of vital PBMC was calculated that dropped to 0.11% in acute flare ups (figs 2E, 2F, and 3A). The mean MDC-1 fraction of CD patients in remission was calculated to be 0.22% of vital PBMC and dropped to 0.14% in acute flare ups (figs 2G, H, and 4A). The same trend was observed for the MDC-2 fraction (data not shown).

Healthy controls were found to have a mean PDC fraction of 0.41% and a mean MDC-1 fraction of 0.25% of vital PBMC, respectively (figs 3A and 4A).

The fraction of circulating PBDC correlates with established IBD disease activity indices

A nearly perfect correlation between disease activity and the fraction of circulating PBDC was found for PDC in UC patients at linear regression analysis expressed as quality of fit (r^2) and correlation coefficient (CC)—PDC and MTWSI (r^2 , 0.86; CC, 0.93) (fig 3B). The other DC fractions were found to have a very high correlation with the respective indices as well—PDC and HBSI (r^2 , 0.63; CC, 0.79) (fig 3C); MDC-1 and MTWSI (r^2 , 0.57; CC, 0.75) (fig 4B); MDC-1 and HBSI (r^2 , 0.65; CC, 0.81) (fig 4C). A similar trend was observed in MDC-2; however due to the low cell numbers no meaningful statistical analysis could be performed.

Circulating PBDC in IBD patients display an immature phenotype which does not change in active disease

Having demonstrated disease activity related changes of PDC and MDC numbers in UC and CD patients, we were interested in the phenotype of the respective cell populations to assess their maturational status. We therefore decided to look at commonly used DC maturation markers.

Virtually all PDC and MDC express MHC class II (HLA-DR) molecules. Low level expression of the co-stimulatory molecule CD86 and the PBDC marker CD62L were detected as well. Other co-stimulators, such as CD80 migration and CD40 and activation markers such as CD83 and CD1a were essentially not expressed at all. Maturation state of PDC and MDC-1 does not change in acute flare ups of IBD—that is, the percentage of activation marker positive cells is not increased (fig 5). No difference between UC and CD was detected. The phenotype of MDC-2 was similar to MDC-1 (data not shown). Taken together, these data suggest that patients with IBD lack immature dendritic cells during acute flare ups.

PBDC express gut homing marker

The observed decrease of all three blood dendritic cell types in active IBD suggests a migration to secondary lymphatic organs. As the gut is an obvious site for such a migratory process in IBD patients, we assessed the expression of a well accepted gut homing marker— $\alpha 4\beta 7$ integrin—which has been associated with intestinal homing of other immune cell populations.^{41 42}

PDC, MDC-1, and MDC-2 express $\alpha 4\beta 7$ integrin. The strongest expression was seen in myeloid DC (fig 6). This correlates with preliminary data from our group on intestinal DC, where a rise of the myeloid subpopulations in active IBD




Figure 2 FACS plots from representative experiments illustrating the significant drop of peripheral blood dendritic plasmacytoid (PDC = PI-(vital), CD123+, BDCA-2+ cells) and myeloid dendritic (MDC-1 = PI-(vital), CD14-, CD19-, CD11c+, BDCA-1) cells in patients with active ulcerative colitis and Crohn's disease. The same trend was seen with MDC-2 (data not shown). Ulcerative colitis PDC: (A) remission v (B) flare up: 0.66% PDC v 0.06% PDC. Ulcerative colitis MDC-1: (C) remission v (D) flare up: 0.25% MDC-1 v 0.06% MDC-1. Crohn's disease PDC: (E) remission v (F) flare up: 0.31% PDC v 0.12% PDC. Crohn's disease MDC-1: (G) remission v (H) flare up: 0.23% MDC-1 v 0.06% MDC-1.

was seen (DCB, unpublished data) and the work from other groups.⁴³ No difference between active and inactive UC and CD was detected (data not shown).

Upon culture, the mean CD86 expression went up in DC from both healthy controls and untreated patients with active CD and UC. However, cultured PDC and MDC-1 from flaring IBD patients on average had a more than twofold higher CD86 expression compared with healthy controls (fig 7, table 4).

PBDC of patients with active IBD respond differently to microbial surrogate stimuli compared with controls

As we hypothesised that PBDC migrate to the gut in patients with active IBD, we were curious to investigate their response to microbial stimuli they might encounter there. We picked




Figure 3 Assessment of circulating plasmacytoid dendritic cells (PDC) in IBD. (A) IBD patients in remission have slightly lower numbers of circulating peripheral blood plasmacytoid dendritic cells compared with healthy controls. In IBD flare-ups PDC drop significantly. Data is expressed as mean percentage of vital PBMC. (B) The fraction of circulating PDC correlates (CC: 0.93) with the modified Truelove Witts severity index (MTWSI) in patients with ulcerative colitis. (C) The fraction of circulating PDC shows a high correlation (CC: 0.80) with the Harvey Bradshaw severity index (HBSI) in patients with Crohn's disease.

two microbial surrogate stimuli, LPS (a bacterial wall compound known to act on MDC) and CpG-ODN 2006 (a bacterial DNA motif known to act uniquely on PDC). Both have been demonstrated to induce maturation, activation, and survival in DC in previous studies.^{40-44, 45} As our phenotypic analysis of circulating PBDC identified low levels




Figure 4 Assessment of circulating myeloid dendritic cells (MDC) in IBD. (A) IBD patients in remission have slightly lower numbers of circulating peripheral blood myeloid (MDC-1) dendritic cells compared with healthy controls. In IBD flare ups MDC drop significantly. Data are expressed as mean percentage of vital PBMC. (B) The fraction of circulating MDC correlates (CC: 0.75) with the modified Truelove Witts severity index (MTWSI) in patients with ulcerative colitis. (C) The fraction of circulating MDC shows a high correlation (CC: 0.81) with the Harvey Bradshaw severity index (HBSI) in patients with Crohn's disease.

of CD86 on both PDC and MDC-1, while all other currently known maturation markers were negative, we decided to further study this co-stimulatory molecule in the following stimulation experiments.

On average, freshly isolated blood DC from healthy controls and untreated patients with active CD and UC




Figure 5 FACS analysis of maturation markers on PDC and MDC in heparinised whole blood. (Stars and asterisks denote extremes and outliers, respectively.) (A) PDC and (B) MDC-1 display an immature phenotype that does not change in acute flare ups.

display higher base line levels of the co-stimulatory molecule CD86 than their respective circulating counterparts analysed in whole blood (figs 5 and 7).

Of note, freshly isolated PDC from both flaring CD and UC patients had a twofold higher mean expression of CD86 compared with healthy controls (fig 7, table 4). Freshly isolated MDC-1 from both healthy controls as well as flaring IBD patients were found to have an almost tenfold higher baseline mean CD86 expression compared with PDC.

Table 4 Expression of the maturation marker CD86

Disease type	DC population	CD86 Expression		
		Freshly isolated (% of PBMC)	Cultured (% of PBMC)	Stimulated (% of PBMC)
Healthy controls	PDC	8.0	27.1	93.2
	MDC-1	71.1	77.4	85.3
Crohn's disease	PDC	15.6	42.7	99.3
	MDC-1	75.7	92.1	95.7
Ulcerative colitis	PDC	14.3	37.3	81.7
	MDC-1	67.3	91.1	90.2




Figure 6 PDC, MDC-1, and MDC-2 express the gut homing marker $\alpha 4\beta 7$ integrin. Example of a patient with Crohn's disease (expression in ulcerative colitis is similar (data not shown)). (A) 93.6% of all vital PDCs are double positive for $\alpha 4\beta 7$ integrin. (B) 67.1% of all vital PDCs are double positive for $\alpha 4\beta 7$ integrin. (C) 98.2% of all vital MDC-2 are double positive for $\alpha 4\beta 7$ integrin.

Differences between healthy controls and patients with active IBD were not as striking as seen with PDC, but CD patients showed a slightly higher CD86 expression (fig 7, table 4).

Following CpG ODN 2006 or LPS stimulation of PDC or MDC-1, the mean expression of CD86 went up even higher in both healthy controls and untreated flaring IBD patients. Most interestingly though, stimulated PDC and MDC-1 from flaring CD patients reached the highest CD86 expression




Figure 7 Histograms summarizing the regulation of the co-stimulatory molecule CD86, a DC maturation marker, in healthy controls (dotted line), and IBD patient PDC and MDC-1, freshly isolated, 24 hour cultured, or stimulated (dotted line) as well as isotype matched controls (bold line). Plots are representative for five independent experiments. Cultured PDC and MDC-1 from untreated IBD patients with acute flare ups express higher levels of the co-stimulatory molecule CD86 compared with healthy controls. CD86 is upregulated faster and to a higher level upon stimulation with microbial surrogate stimuli such as CpG ODN 2006 or LPS, respectively.

compared with healthy controls and flaring UC patients (fig 7, table 4).

In summary, freshly isolated PDC and MDC-1 from untreated flaring IBD patients express higher baseline levels of the co-stimulatory molecule CD86 compared with healthy controls. The expression increases further in culture and upon stimulation with microbial surrogate stimuli such as CpG ODN 2006 and LPS to an overall higher level compared with healthy controls.

DISCUSSION

Dendritic cells are thought to be involved in the maintenance of tolerance depending on their location, number, and

maturation state.²⁻⁴⁶ Despite the effectiveness of central (thymic) tolerance, self-reactive T cells can escape deletion and editing. In addition, the human body is constantly exposed to environmental antigens including food proteins and the intestinal commensal microflora. These limitations require additional mechanisms to effectively silence (auto) reactive T cells.⁴⁷ It is assumed that blood DC in healthy individuals display an immature phenotype and induce T cell unresponsiveness.⁴⁸⁻⁴⁹ Little is known about circulating DC in IBD patients.

Our data demonstrate that patients with acute flare ups of Crohn's disease and ulcerative colitis experience a significant drop in their peripheral PDC and MDC dendritic cell

populations. The observed changes in the studied DC populations may not be IBD specific, but the demonstrated striking correlation with established disease activity indices underscore their importance in the disease process. Certainly, other, acute inflammatory bowel conditions such as infectious colitis or diverticulitis may share some of the observed phenomena.

Similar changes in the circulating DC population have been reported in patients with other immune disorders—for example, systemic lupus (SLE) and HIV, where the number of DC dropped peripherally and—for SLE patients—at the same time increased in skin lesions.^{50–52} This suggests a migration of peripheral DC to other sites, possibly to the gut in IBD patients. We provide evidence for this hypothesis by demonstrating the expression of a widely accepted gut homing marker $\alpha 4\beta 7$ integrin on PBDC (fig 6). Our view is further substantiated by the work of other groups that demonstrated functional chemokine receptors required for migration on both plasmacytoid and myeloid dendritic cells.^{53–57}

A well accepted concept for the pathogenesis of IBD is an inappropriate T cell response towards luminal antigens of microbial origin.^{23–24} The inappropriate inflammatory response may not occur primarily or exclusively due to T cell dysfunction, but is probably also promoted by a disturbed intestinal barrier and/or an aberrant antigen presenting cell—that is, DC function.^{58–59} The resulting imbalance between proinflammatory (that is, TNF- α , IFN- γ) and anti-inflammatory cytokines (that is, TGF- β , IL10), is caused by a differential activation of Th1, Th2 and Tr, Th3 cells, respectively, which is in part disease specific (Crohn's disease, Th1; ulcerative colitis, Th2).^{60–61} We provide preliminary evidence for DC dysfunction by demonstrating a different regulation of the co-stimulatory molecule CD86, a well accepted DC maturation marker, in flaring IBD patients. The observed differences between PDC and MDC as well as CD and UC likely reflect the long known differences in the pathophysiology of these two disorders.

Due to their dual role as tolerogenic or activating antigen presenting cells, depending on their maturational state, DC may induce the differentiation of distinct T cell subsets in IBD patients compared with healthy individuals. In other words it is conceivable that circulating, immature DC normally maintain tolerance by inducing Th3 and Tr cells and, once migrated to the (inflamed) intestine and matured, promote inflammation by activating Th1 cells in IBD patients.

Our study provides preliminary evidence for this hypothesis by demonstrating a more vigorous response of circulating PDC and MDC towards microbial surrogate stimuli such as LPS and CpG ODN 2006. The postulated migration and aberrant activation process may subsequently induce the lack of circulating DC reported in our study. Although this response may occur in other inflammatory bowel conditions, such as infectious colitis and diverticulitis (not included in this analysis) our findings further support the hypothesis of their potential involvement in the breakdown of tolerance in IBD.

We are aware that medications *may* influence the phenotype of circulating DC, although a recent meta-analysis concluded that immunomodulators do *not* influence DC maturation.⁶² Certainly, it would have been ideal to study exclusively cells from patients off all drugs. It is unrealistic and unethical though to study a large cohort of untreated IBD patients when most medications are widely available and in use. However, as demonstrated in figure 1 we have preferably analysed samples from patients off immune modulators or prior to adding them to control flare ups. Several patients in remission were receiving IBD related drugs. Interestingly though, the reported drop of circulating

DC occurred in the group that received the least drugs—patients with acute flare ups *before* therapy. The patients in remission had PDC and MDC fractions closer to those of controls, which in our opinion supports the view that the DC counts and phenotypes in our study are drug independent.

Two other groups found that PBDC from patients with UC showed significantly increased stimulatory capacities and produced significantly higher levels of nitric oxide compared with controls with infectious or ischaemic colitis. In contrast to our study, UC patients' blood DC displayed a mature, activated phenotype in one study.^{63–64} However, due to the lack of specific antibodies these studies did not distinguish between PDC or MDC and the unequivocal identification of DC versus monocytes/macrophages must be questioned because of the lack of specific antibodies at that time.

In conclusion, we have demonstrated a lack of immature, possibly tolerogenic peripheral DC in patients with active IBD. Circulating DC express the gut homing marker, $\alpha 4\beta 7$ integrin. Moreover, preliminary evidence suggests a different response of IBD patient DC to microbial surrogate stimuli. Taken together these data suggest a migratory process of blood DC in IBD patients to secondary lymphatic organs such as the intestine, where they potentially mature, become activated, and contribute to gut inflammation and tissue damage. Our data support the hypothesis that dendritic cells are involved in the breakdown of tolerance in IBD. Future studies will need to address functional properties of blood DC in IBD in detail and elucidate the interaction of peripheral and intestinal subsets.

ACKNOWLEDGEMENTS

This work was supported by research grants of The Eli & Edythe L Broad Foundation, Los Angeles, CA, USA and Deutsche Forschungsgemeinschaft, Bonn, Germany as well as a Humboldt-University Bonus Research Grant to DCB. The authors thank the patients, medical, laboratory, and nursing staff of the Charité Inflammatory Bowel Disease Clinic for generously donating their blood for this study.

Authors' affiliations

D C Baumgart, D Metzke, A Sturm, B Wiedenmann, A U Dignass, Department of Medicine, Division of Hepatology & Gastroenterology, Charité Medical Centre-Virchow Hospital, Medical School of the Humboldt-University of Berlin, Germany
J Schmitz, A Scheffold, Immunocytometry Group, German Rheumatism Research Centre (DRFZ), Berlin, Germany

REFERENCES

- 1 Langerhans P. Über die Nerven der menschlichen Haut. *Virchows Arch Pathol Anat Physiol Klin Med* 1868;**44**:325.
- 2 Banchereau J, Briere F, Caux C, et al. Immunobiology of dendritic cells. *Annu Rev Immunol* 2000;**18**:767–811.
- 3 Banchereau J, Steinman RM. Dendritic cells and the control of immunity. *Nature* 1998;**392**:245–52.
- 4 Hornung V, Rothenfusser S, Britsch S, et al. Quantitative expression of toll-like receptor 1–10 mRNA in cellular subsets of human peripheral blood mononuclear cells and sensitivity to CpG oligodeoxynucleotides. *J Immunol* 2002;**168**:4531–7.
- 5 Krug A, Towarowski A, Britsch S, et al. Toll-like receptor expression reveals CpG DNA as a unique microbial stimulus for plasmacytoid dendritic cells which synergizes with CD40 ligand to induce high amounts of IL-12. *Eur J Immunol* 2001;**31**:3026–37.
- 6 Stagg AJ, Kamm MA, Knight SC. Intestinal dendritic cells increase T cell expression of alpha4beta7 integrin. *Eur J Immunol* 2002;**32**:1445–54.
- 7 Leithauser F, Trobonjaca Z, Moller P, et al. Clustering of colonic lamina propria CD4(+) T cells to subepithelial dendritic cell aggregates precedes the development of colitis in a murine adoptive transfer model. *Lab Invest* 2001;**81**:1339–49.
- 8 Mora JR, Bono MR, Manjunath N, et al. Selective imprinting of gut-homing T cells by Peyer's patch dendritic cells. *Nature* 2003;**424**:88–93.
- 9 Iwasaki A, Kelsall BL. Mucosal immunity and inflammation. I. Mucosal dendritic cells: their specialized role in initiating T cell responses. *Am J Physiol* 1999;**276**:G1074–G1078.
- 10 Singh B, Read S, Asseman C, et al. Control of intestinal inflammation by regulatory T cells. *Immunol Rev* 2001;**182**:190–200.

- 11 Rescigno M, Urbano M, Valzasina B, et al. Dendritic cells express tight junction proteins and penetrate gut epithelial monolayers to sample bacteria. *Nat Immunol* 2001;**2**:361-7.
- 12 Iwasaki A, Kelsall BL. Unique functions of CD11b+, CD8 alpha+, and double-negative Peyer's patch dendritic cells. *J Immunol* 2001;**166**:4884-90.
- 13 Iwasaki A, Kelsall BL. Freshly isolated Peyer's patch, but not spleen, dendritic cells produce interleukin 10 and induce the differentiation of T helper type 2 cells. *J Exp Med* 1999;**190**:229-39.
- 14 Iwasaki A, Kelsall BL. Localization of distinct Peyer's patch dendritic cell subsets and their recruitment by chemokines macrophage inflammatory protein (MIP)-3alpha, MIP-3beta, and secondary lymphoid organ chemokine. *J Exp Med* 2000;**191**:1381-94.
- 15 Singh B, Powrie F, Mortensen NJ. Immune therapy in inflammatory bowel disease and models of colitis. *Br J Surg* 2001;**88**:1558-69.
- 16 Maraskovsky E, Brasel K, Teepe M, et al. Dramatic increase in the numbers of functionally mature dendritic cells in Flt3 ligand-treated mice: multiple dendritic cell subpopulations identified. *J Exp Med* 1996;**184**:1953-62.
- 17 Henri S, Vremec D, Kamath A, et al. The dendritic cell populations of mouse lymph nodes. *J Immunol* 2001;**167**:741-8.
- 18 Belyakov IM, Hel Z, Kelsall B, et al. Mucosal AIDS vaccine reduces disease and viral load in gut reservoir and blood after mucosal infection of macaques. *Nat Med* 2001;**7**:1320-6.
- 19 Kelsall BL, Strober W. Dendritic cells of the gastrointestinal tract. *Springer Semin Immunopathol* 1997;**18**:409-20.
- 20 Malmstrom V, Shipton D, Singh B, et al. CD134L expression on dendritic cells in the mesenteric lymph nodes drives colitis in T cell-restored SCID mice. *J Immunol* 2001;**166**:6972-81.
- 21 Stagg AJ, Hart AL, Knight SC, et al. The dendritic cell: its role in intestinal inflammation and relationship with gut bacteria. *Gut* 2003;**52**:1522-9.
- 22 Mayer L. Oral tolerance: new approaches, new problems. *Clin Immunol* 2000;**94**:1-8.
- 23 Duchmann R, Kaiser I, Hermann E, et al. Tolerance exists towards resident intestinal flora but is broken in active inflammatory bowel disease (IBD) (see comments). *Clin Exp Immunol* 1995;**102**:448-55.
- 24 MacDonald TT. Breakdown of tolerance to the intestinal bacterial flora in inflammatory bowel disease (IBD). *Clin Exp Immunol* 1995;**102**:445-7.
- 25 Baumgarth N, Roederer M. A practical approach to multicolor flow cytometry for immunophenotyping. *J Immunol Methods* 2000;**243**:77-97.
- 26 Steinman RM. DC-SIGN: a guide to some mysteries of dendritic cells. *Cell* 2000;**100**:491-4.
- 27 Jameson B, Baribaud F, Pohlmann S, et al. Expression of DC-SIGN by dendritic cells of intestinal and genital mucosae in humans and rhesus macaques. *J Virol* 2002;**76**:1866-75.
- 28 Dzionek A, Fuchs A, Schmidt P, et al. BDCA-2, BDCA-3, and BDCA-4: three markers for distinct subsets of dendritic cells in human peripheral blood. *J Immunol* 2000;**165**:6037-46.
- 29 Dzionek A, Sohma Y, Nagafune J, et al. BDCA-2, a novel plasmacytoid dendritic cell-specific type II C-type lectin, mediates antigen capture and is a potent inhibitor of interferon alpha/beta induction. *J Exp Med* 2001;**194**:1823-34.
- 30 Truelove SC, Witts LJ. Cortisone in ulcerative colitis; preliminary report on a therapeutic trial. *BMJ* 1954;**4884**:375-8.
- 31 Truelove SC, Witts LJ. Cortisone in ulcerative colitis; final report on a therapeutic trial. *BMJ* 1955;**4947**:1041-8.
- 32 Harvey RF, Bradshaw JM. A simple index of Crohn's-disease activity. *Lancet* 1980;**1**:514.
- 33 Maier K, von Gaisberg U, Kraus B. Ulcerative colitis. Activity index for the clinical and histological classification of inflammatory activity. *Schweiz Med Wochenschr* 1988;**118**:763-6.
- 34 Seo M, Okada M, Yao T, et al. Evaluation of disease activity in patients with moderately active ulcerative colitis: comparisons between a new activity index and Truelove and Witts' classification. *Am J Gastroenterol* 1995;**90**:1759-63.
- 35 Li B, Bassiri H, Rossman MD, et al. Involvement of the Fas/Fas ligand pathway in activation-induced cell death of mycobacteria-reactive human gamma delta T cells: a mechanism for the loss of gamma delta T cells in patients with pulmonary tuberculosis. *J Immunol* 1998;**161**:1558-67.
- 36 Baumgart DC, Olivier WA, Reya T, et al. Mechanisms of intestinal epithelial cell injury and colitis in interleukin 2 (IL2)-deficient mice. *Cell Immunol* 1998;**187**:52-66.
- 37 Dzionek A, Fuchs A, Schmidt P, et al. BDCA-2, BDCA-3, and BDCA-4: three markers for distinct subsets of dendritic cells in human peripheral blood. *J Immunol* 2000;**165**:6037-46.
- 38 Krug A, Towarowski A, Britsch S, et al. Toll-like receptor expression reveals CpG DNA as a unique microbial stimulus for plasmacytoid dendritic cells which synergizes with CD40 ligand to induce high amounts of IL-12. *Eur J Immunol* 2001;**31**:3026-37.
- 39 Krug A, Rothenfusser S, Hornung V, et al. Identification of CpG oligonucleotide sequences with high induction of IFN-alpha/beta in plasmacytoid dendritic cells. *Eur J Immunol* 2001;**31**:2154-63.
- 40 Krieg AM. CpG motifs in bacterial DNA and their immune effects. *Annu Rev Immunol* 2002;**20**:709-60.
- 41 Farstad IN, Halstensen TS, Lien B, et al. Distribution of beta 7 integrins in human intestinal mucosa and organized gut-associated lymphoid tissue. *Immunology* 1996;**89**:227-37.
- 42 Wagner N, Lohler J, Kunkel EJ, et al. Critical role for beta7 integrins in formation of the gut-associated lymphoid tissue. *Nature* 1996;**382**:366-70.
- 43 te Velde AA, van Kooyk Y, Braat H, et al. Increased expression of DC-SIGN+IL-12+IL-18+ and CD83+IL-12-IL-18- dendritic cell populations in the colonic mucosa of patients with Crohn's disease. *Eur J Immunol* 2003;**33**:143-51.
- 44 Krug A, Rothenfusser S, Hornung V, et al. Identification of CpG oligonucleotide sequences with high induction of IFN-alpha/beta in plasmacytoid dendritic cells. *Eur J Immunol* 2001;**31**:2154-63.
- 45 Krug A, Towarowski A, Britsch S, et al. Toll-like receptor expression reveals CpG DNA as a unique microbial stimulus for plasmacytoid dendritic cells which synergizes with CD40 ligand to induce high amounts of IL-12. *Eur J Immunol* 2001;**31**:3026-37.
- 46 Banchereau J, Steinman RM. Dendritic cells and the control of immunity. *Nature* 1998;**392**:245-52.
- 47 Steinman RM, Nussenzweig MC. Avoiding horror autotoxicus: the importance of dendritic cells in peripheral T cell tolerance. *Proc Natl Acad Sci U S A* 2002;**99**:351-8.
- 48 Hawiger D, Inaba K, Dorsett Y, et al. Dendritic cells induce peripheral T cell unresponsiveness under steady state conditions in vivo. *J Exp Med* 2001;**194**:769-79.
- 49 Steinman RM, Nussenzweig MC. Avoiding horror autotoxicus: the importance of dendritic cells in peripheral T cell tolerance. *Proc Natl Acad Sci U S A* 2002;**99**:351-8.
- 50 Cederblad B, Blomberg S, Vallin H, et al. Patients with systemic lupus erythematosus have reduced numbers of circulating natural interferon-alpha-producing cells. *J Autoimmun* 1998;**11**:465-70.
- 51 Blanco P, Palucka AK, Gill M, et al. Induction of dendritic cell differentiation by IFN-alpha in systemic lupus erythematosus. *Science* 2001;**294**:1540-3.
- 52 Chehimi J, Campbell DE, Azzoni L, et al. Persistent decreases in blood plasmacytoid dendritic cell number and function despite effective highly active antiretroviral therapy and increased blood myeloid dendritic cells in HIV-infected individuals. *J Immunol* 2002;**168**:4796-801.
- 53 Liu YJ, Kanzler H, Soumelis V, et al. Dendritic cell lineage, plasticity and cross-regulation. *Nat Immunol* 2001;**2**:585-9.
- 54 Liu YJ. Dendritic cell subsets and lineages, and their functions in innate and adaptive immunity. *Cell* 2001;**106**:259-62.
- 55 Penna G, Vulcano M, Sozzani S, et al. Differential migration behavior and chemokine production by myeloid and plasmacytoid dendritic cells. *Hum Immunol* 2002;**63**:1164-71.
- 56 Steinman R, Hoffman L, Pope M. Maturation and migration of cutaneous dendritic cells. *J Invest Dermatol* 1995;**105**(1 Suppl):2S-7S.
- 57 Jonuleit H, Knop J, Enk AH. Cytokines and their effects on maturation, differentiation and migration of dendritic cells. *Arch Dermatol Res* 1996;**289**:1-8.
- 58 Baumgart DC, Dignass AU. Intestinal barrier function. *Curr Opin Clin Nutr Metab Care* 2002;**5**:685-94.
- 59 Baumgart DC, Dignass AU. Mucosal dendritic cells: reaching out for the unknown (antigen). *Z Gastroenterol* 2002;**40**:375-7.
- 60 Blumberg RS, Strober W. Prospects for research in inflammatory bowel disease. *JAMA* 2001;**285**:643-7.
- 61 Neurath MF, Finotto S, Glimcher LH. The role of Th1/Th2 polarization in mucosal immunity. *Nat Med* 2002;**8**:567-73.
- 62 Lagaraine C, Lebranchu Y. Effects of immunosuppressive drugs on dendritic cells and tolerance induction. *Transplantation* 2003;**75**(9 Suppl):37S-42S.
- 63 Ikeda Y, Akbar F, Matsui H, et al. Characterization of antigen-presenting dendritic cells in the peripheral blood and colonic mucosa of patients with ulcerative colitis. *Eur J Gastroenterol Hepatol* 2001;**13**:841-50.
- 64 Vuckovic S, Florin TH, Khalil D, et al. CD40 and CD86 upregulation with divergent CMRF44 expression on blood dendritic cells in inflammatory bowel diseases. *Am J Gastroenterol* 2001;**96**:2946-56.