
Gut June 1965



In cases of dyspepsia with or without a radiologically proven ulcer Roter will normally provide prompt symptomatic relief and lead to rapid healing of the lesion.

The reputation of Roter is world wide. It is a complete treatment in itself and requires no additional antispasmodics or anticholinergics.



FORMULA: Each Roter tablet contains: Bismuth Subnitrate

Roter (350 mg.), Magnesium Carbonate (400 mg.), Sodium

Bicarbonate (200 mg.) and Cortex Rhamni Frangulae (25 mg.).

PACKINGS OF ROTER TABLETS: Tins of 40 and 120, also dispensing sizes, 360 and 720 (P.T. exempt).

ROTER IN PEPTIC ULCER

1



FOR COMPLETE PARENTERAL NUTRITION

. . . without risk of Thrombophlebitis



literature and suggested regimes on request

PAINES & BYRNE LTD., GREENFORD, MIDDLESEX pre-eminent in Parenteral Nutrition

Lomotil does just one thing it stops diarrhoea

... patients began to feel more comfortable ... after approximately one hour.'

Hock, C.W. (1961) J.med.Ass.Ga., **50**,485

Most diarrhoeas are non-specific in origin. Lomotil (brand of diphenoxylate hydrochloride with atropine sulphate) provides the rapid, symptomatic control needed in such cases.

In diarrhoea of specific origin, Lomotil conserves electrolytes and prevents dehydration while antibacterial treatment takes effect.

Lomotil curbs the intestinal hypermotility characteristic of diarrhoea. It is effective in virtually all types of diarrhoea — acute, chronic or recurrent.

Lomotil is a registered trade mark

THE RAPID-ACTING ANTI-DIARRHOEAL Lomotil
Searle

The success of BIOGASTRONE in gastric ulcer

further proof in 1965



66 It is the only drug therapy for gastric ulcer which has been demonstrated conclusively to accelerate healing, which can be achieved with patients remaining up and about, often at work, and with minimum change in diet ??

from editorial synopsis: GUT (1965) 6:19



In 1959, the first clinical trial of Biogastrone* was begun in patients with gastric ulcer. The results indicated that Biogastrone significantly reduced the size of ulcer niche.

"It is suggested that Biogastrone promotes the healing of gastric ulcers."

Lancet (1962) ii: 793

Now, in a further trial, the same author proves beyond any doubt that Biogastrone is successful in assisting gastric ulcers to heal. The average degree of healing was similar in both trials —72 and 78% in the Biogastrone groups and 35 and 39% for the control.


"It is concluded that carbenoxolone sodium [Biogastrone] facilitates the healing of gastric ulcers."

Gut (1965) 6: 19

More detailed information from Berk Pharmaceuticals Limited, Catteshall Lane Godalming, Surrey. * trade mark BR 3775/J817

Have Philosophers you a new autors necessity that is, a read a part of second for a part of s





2 TABLETS TWICE A DAY— PROVEN MAINTENANCE THERAPY

In Ulcerative Colitis

available as a 0.5 g. tablet, either plain or enteric coated

FIRST DEMONSTRATION . . . RESULTS SIGNIFICANT

"This is the first demonstration in a formal trial that any treatment reduces the relapse-rate in ulcerative colitis, 24 out of 34 patients taking 2 g. of sulphasalazine daily remained free of symptoms for a year. This result is significantly different from that obtained in a comparable group of patients who received a placebo."

PREFERABLE TO SYSTEMIC CORTICOSTEROIDS

"We have shown that sulphasalazine, 0.5 g. four times daily, is often effective in maintaining remission . . . and it therefore appears preferable to systemic corticosteroid drugs for this purpose."

REMISSION FOR ONE YEAR

"24 patients remained in symptomatic remission for a year while taking 2 g. of sulphasalazine daily, whereas only 8 remained symptom-free in the placebo group."

NORMAL MUCOSA

"2 of the 24 patients who remained symptom-free for a year while on sulphasalazine had a haemorrhagic mucosa at the end of the trial. The remaining 22 patients had a non-haemorrhagic mucosa which in many cases appeared normal."

MINIMAL SIDE EFFECTS

"In the present trial, using a dose of 2 g. daily, only 3 patients out of 34 had to discontinue treatment because of side-effects."

WHITE CELL COUNT

"In the patients treated with sulphasalazine there was no difference in the haemoglobin level before and after treatment; but the mean white-cell count was lower after six months or a year than at the start of treatment, though in no patient was it less than 4500 per c.mm."

The Lancet Jan 23rd 1965 \cdot Vol I \cdot Pages 185-188

SALAZOPYRIN



Literature and further information are supplied on request SINCLAIR HOUSE \cdot THE AVENUE \cdot WEST EALING \cdot LONDON W13 Distributed by Martindale Samoore Ltd \cdot 60-61 Welbeck St \cdot London W1



Strength in hand



with London Hospital Sutures




Strong, sure, supple — packaged for speed and safety.

Available in a full range of catgut and non-absorbable materials with or without eyeless needles



IX Gut June 1965



LONGER-LASTING DRIP THERAPY

Gastrils—first ever antacid pastilles. Gastrils do not break up in the mouth like tablets; they dissolve slowly and smoothly, so give sustained drip therapy for the treatment of peptic ulcer.

Gastrils contain a co-dried gel of aluminium hydroxide and magnesium carbonate, a buffer antacid comparable in activity to freshly prepared liquid aluminium hydroxide gel. Gastrils rapidly buffer gastric pH to an optimum level, and keep it there, without risk of acid

To relieve the tedium of constant antacid therapy, every Gastrils prescription is dispensed in equal quantities of two delicious flavours-mint and fruit. Few antacids are as effective as Gastrils—none is as good to take!

GASTRILS

ANTACID PASTILLES

INDICATIONS Hyperacidity, peptic ulcer, gastritis, heartburn, oesophagitis and hiatus hernia.

FORMULA


Each pastille contains 500 mg. co-dried gel of aluminium hydroxide and magnesium carbonate. BASIC N.H.S. 24 individually wrapped Gastrils in carton 3/-

24 individually wrapped Gastrils from dispensing pack: 2/61d. Gastrils are prescribable on form E.C.10

'Gastrils' is a registered trade mark of T. J. Smith & Nephew Ltd.



WELWYN GARDEN CITY · HERTFORDSHIRE · ENGLAND



Pancreatitis may be difficult to differentiate

from a coronary thrombosis, perforated peptic ulcer, a dissecting aneurysm of the aorta or acute cholecystitis, but when the diagnosis is established, TRASYLOL the proteinase inactivator is indicated.

TRASYLOL specifically inhibits the proteolytic enzymes of the pancreas, and administered in adequate dosage arrests the autodigestive processes.

A recent analysis of all cases of pancreatitis, those treated with TRASYLOL showed a reduction in mortality from 22% to 9%.

TRASYLOL is therefore indicated in the treatment of acute pancreatitis, pancreatic fistulae, and pancreatitis accompanying disorders of the gall bladder and bile ducts.



don't use Merbentyl to inhibit gastric secretion

because

Merbentyl is not an antisecretory drug. Do use it to control gastrointestinal spasm because here Merbentyl is a remarkably effective antispasmodic. Moreover, Merbentyl is so selective that its use is rarely followed by side effects commonly seen with other drugs used for this purpose. It can even be administered to patients with chronic simple glaucoma or prostatic hypertrophy.

Merbentyl (dicyclomine hydrochloride B.P.C.) was first described in 1947, and has been widely used clinically to control gastrointestinal spasm and hypermotility in many countries for over 12 years. It acts both on the neuroreceptor sites innervated by post-ganglionic parasympathetic fibres, and directly on smooth muscle tissue.

Indications

- 1 Primary functional disorders of the gastro-intestinal tract: spastic colon, pylorospasm, post-cholecystectomy syndrome.
- 2 Adjunctive therapy in organic disorders, e.g. ulcerative colitis, diverticulitis.
- 3 Infant colic (here the drug is regarded as specific therapy).

Low NHS. cost An average week's treatment—2/8d.

Merbentyl tablets each tablet contains 10mg. dicyclomine hydrochloride B.P.C. Recommended adult dosage, 2 t.d.s.

Also available with phenobarbitone and as Merbentyl Syrup.

Full clinical information available on request.




Merrell-National (Laboratories) Limited, 20 Savile Row, London, W.1.

*Prodexin.

continuous drip control of ulcer

by tablet



PRODEXIN tablets give all the benefits of continuous intragastric drip therapy¹ without any of its disadvantages.

The slow suck down tablets are specially formulated to give prolonged antacid buffer effect, unaffected by stomach emptying rate.²

Immediate relief of pain is achieved by rapidly raising pH to 2 neutralising excess acid and controlling the acid secretion by slowly building up the pH to a desirable level of pH3.5 and pH4.5

DOSAGE:

For continuous therapeutic effect one tablet at a time should be sucked slowly. For peptic ulcer one or more tablets an hour.

References:

- 1. Practitioner (1954), 173, 46.
- 2. Post Grad. med. J. (1960) 36, 722.

*Aluminium Glycinate 0.9G. Light Magnesium Carbonate B.P. 0.1G.

PRODEXIN is a product of British research at

Beecham Research Laboratories



VELACTIN

A readily assimilated milk substitute of vegetable origin for Milk Allergies



M441

Velactin contains no gluten or lactose The biological value of the soya protein has been improved by the addition of DL-Methionine to make it equivalent to that of first-class animal protein.

Literature gladly sent on request A. Wander Limited, 42 Upper Grosvenor Street, London W.1. **Indications:** Allergy or intolerance to human or cow's milk.¹ As basic feeding pending identification of specific food allergens.¹ Galactosaemia.^{2,3} Gluten enteropathy. Velactin may also be tried in cases of undetermined metabolic or non-specific intolerance.

References: (1) Nutritio et Dieta 1961; 3, 89.

(2) Lancet 1959, 2 849; 1960 1, 336.

(3) Ugeskrift f. Laeger 1960, 122, 1229.

Presentation: Tin containing 9 oz. net.

ANTIBIOTICS AND THE INTESTINAL FLORA

Treatment with broad spectrum antibiotics suppresses the normal flora, which may be replaced by abnormal resistant organisms causing diarrhoea.

This change can be prevented or reversed by the implantation of *Lactobacillus acidophilus* which has been rendered antibiotic-resistant, and is thus able to multiply freely under these abnormal conditions. The products of growth of this organism are inimical to that of many other bacteria^{1,2} and thus contribute to its rapid preponderance over other species.

Oral treatment with antibiotic-resistant *L. acidophilus* in the form of ENPAC has been shown to eliminate *Candida* from the faeces of infants with antibiotic-induced diarrhoea, with relief of symptoms succeeded by rapid gain in weight^{3,4} and greatly to reduce the numbers of staphylococci in the faeces of patients treated with tetracyclines.⁵

¹ (1958) Ann. Inst. Pasteur **95**, 194. ² (1959) J. Bact. **78**, 477. ³ (1957) Klin. Wschr. **35**, 198. ⁴ (1959) Medizinische **7**, 296. ⁵ (1957) Lancet (i), 899.



LACTOBACILLUS ACIDOPHILUS PREPARATION

Samples and literature will be sent to the medical profession on request to:-

APLIN and BARRETT LIMITED . BIOLOGICALS DIVISION . TROWBRIDGE . WILTS

SECOND SYMPOSIUM OF THE FACULTY OF MEDICINE UNIVERSITY OF ALBERTA

"Gastric Secretion — Mechanisms and Control"

An international symposium to be held on September 13, 14 and 15, 1965, at the University of Alberta, Edmonton, Canada.

The following authorities will participate:

Andersson, Dr. Sven Blair, Dr. E. L. Card, Dr. W. I. Clarke, Dr. J. S. Code, Dr. C. F. Davies, Dr. R. E. Dragstedt, Dr. L. R. Elliott, Dr. Dan W. Ellison, Dr. E. H. Glass, Dr. G. B. J. Gregory, Dr. R. E. Grossman, Dr. M. I. Hollander, Dr. F. Irvine, Prof. W. T. Stockholm Newcastle upon Tyne Edinburgh Albuquerque Rochester Philadelphia Gainesville, Fla. Pittsburgh Milwaukee New York Liverpool Los Angeles New York London Ito, Dr. S.
Jeffries, Dr. Graham H.
Jordan, Dr. Paul H.
Leonard, Dr. A. S.
Menguy, Dr. René
Murray, Prof. J. G.
Olbe, Dr. Lars
Rasanen, Dr. Toimi
Rudick, Dr. J.
Schofield, Dr. B.
Sircus, Dr. W.
Teir, Dr. Harold
Thompson, Dr. J. C.
Webster, Dr. D. R.

Boston
New York
Houston
Minneapolis
Lexington
London
Gothenburg
Helsinki
Seattle
Newcastle upon Tyne
Edinburgh
Helsinki
Torrance, California

Montreal

Registration Fee: \$50.00

Further details may be obtained from Chairman of Symposium, Room C148, University Hospital, Edmonton, Alberta, Canada.

BRITISH



A comprehensive general medical journal. Essential for every medical man or student who wishes to keep informed of research in medical science, of trends in clinical medicine and of professional activities.

Weekly. Annual Subscription, United Kingdom, Irish Republic, £9 9s.; Abroad £5 5s. (\$15.00).

and its associated specialist journals

BRITISH JOURNAL OF OPHTHALMOLOGY

Designed for the clinician and research worker and includes original articles, case notes, information on new appliances, book reviews.

Monthly. Annual Subscription, Inland £6 6s.; Overseas £7(\$20.00). (Combined subscription for the two Ophthalmic journals, Inland £11 11s.; Overseas £13 (\$37.00)).

OPHTHALMIC LITERATURE

Abstracts of articles on the wide field of ophthalmology published throughout the world. Each volume contains some 5,000 abstracts—covering one year.

Six issues a year plus index. Annual Subscription, Inland £6 6s.; Overseas £7 (\$20.00). (For combined subscriptions

see above.)

JOURNAL OF CLINICAL PATHOLOGY

Original and specially commissioned articles on each branch of pathology, with prominence given to its clinical application. Technical methods, notes on new or improved apparatus, book reviews.

**Annual Subscription (six issues), Inland £4 4s.; Overseas £4 15s. (\$14.00).

BRITISH HEART JOURNAL

Original work on the heart and circulation—anatomical, physiological and pathological.

Annual Subscription (six issues), Inland £4 4s.; Overseas £4 15s. (\$14.00).

ARCHIVES OF DISEASE IN CHILDHOOD

Devoted to child health and disease, with emphasis on clinical paediatrics. Original articles by paediatricians and other specialists.

Annual Subscription (six issues), Inland £4 4s.; Overseas £4 15s. (\$14.00).

BRITISH JOURNAL OF PHARMACOLOGY AND CHEMOTHERAPY

Reports on original work in all branches of the subject, including experimental chemotherapy. Biochemical and pathological studies.

Annual Subscription (six issues), Inland £10 10s.; Overseas £11 (\$32.00).

JOURNAL OF NEUROLOGY, NEUROSURGERY AND PSYCHIATRY
Original articles on current research, throwing light on practice in neurology, neurosurgery and psychiatry.

Annual Subscription (six issues), Inland £4 4s.; Overseas £4 15s. (\$14.00).

ANNALS OF THE RHEUMATIC DISEASES
Original articles by authorities in the U.K., U.S.A. and elsewhere on every manifestation of the rheumatic diseases.
Summaries in English, French and Spanish. Annual Subscription (six issues), Inland £4 4s.; Overseas £4 15s. (\$14.00).

THORAX

Original work on the anatomy, physiology and pathology of the chest and heart. Descriptions of techniques and apparatus used in modern cardiac surgery. Annual Subscription (six issues), Inland £4 4s.; Overseas £4 15s. (\$14.00).

Journal of the British Society of Gastroenterology. A journal for physicians, surgeons, radiologists, and pathologists, carrying authoritative articles on all aspects of gastroenterology.

Annual Subscription (six issues), Inland £4 4s.; Overseas £4 15s. (\$14.00).

JOURNAL OF MEDICAL GENETICS

A new journal publishing articles on clinical, statistical, pathological, cytological and biochemical aspects of medical genetics.

Quarterly. Annual Subscription, Inland £3 3s.; Overseas £3 10s. (\$10.00).

BRITISH JOURNAL OF VENEREAL DISEASES

Designed to disseminate, exchange and increase knowledge among consultants, practitioners, research workers, teachers and others concerned with the problem of venereal diseases.

Quarterly. Annual Subscription, Inland £3 3s.: Overseas £3 10s. (\$10.00).

BRITISH JOURNAL OF INDUSTRIAL MEDICINE

Authoritative papers on all aspects of occupational medicine, e.g., the pneumoconioses, occupational cancers, industrial toxicology and psychology. Quarterly. Annual Subscription, Inland £3 3s.; Overseas £3 10s. (\$10.00). BRITISH JOURNAL OF PREVENTIVE AND SOCIAL MEDICINE

Original work in the field, the emphasis on epidemiological methods and medical research.

Quarterly. Annual Subscription, Inland £3 3s.; Overseas £3 10s. (\$10.00).

MEDICAL AND BIOLOGICAL ILLUSTRATION

Illustration and it application to the requirements of the medical and biological sciences.

Quarterly. Annual Subscription, Inland £3 3s.; Overseas £3 10s. (\$10.00).

ABSTRACTS OF WORLD MEDICINE

Critical abstracts of important articles appearing in medical periodicals in every part of the world. Accent on what is new.

Monthly. Annual Subscription, Inland and Overseas £6 6s. (\$18.00).

Write to The Subscription Manager, BRITISH MEDICAL JOURNAL, B.M.A. House, Tavistock Square, London W.C.1

or order through any leading subscription agent or bookseller (All dollar rates quoted refer to the U.S. dollar)

ALUHYDE

for prompt, safe symptomatic medication and hyperchlorhydria and peptic alcor

'ALUHYDE' tablets present a new combination with these advantages...

'ALUHYDE' effects safe, reliable, nonsystemic neutralization by means of two of the most preferred modern antacid adsorbents...



sedative-effect by its inclusion of a barbiturate recognized for minimal cumulative potential...



'ALUHYDE' provides antispasmodic action through its content of an agent long known for its favourable influence against hypermotility...



Formula-

Each tablet: Dried Aluminium

Hydroxide Gel. B.P.C. - 3.75 gr.

Magnesium trisilicate B.P. - 3.75 gr.

Quinalbarbitone sodium B.P. - 0.50 gr.

Tinct. Belladonnæ B.P. - 3 m.

Packings-Regular packs of 50; bottles of 500 (dispensing)

Samples available on personally signed request of qualified physicians only

MULTIPAX CHEMICALS LTD

121 NORWOOD ROAD, LONDON, S.E.24

Phone: TULse Hill 3276

NDON, S.E.24

IA53