PTU-112

A COMPARISON OF INCIDENCE OF COLORECTAL CANCER AND POLYPS IN BOWEL CANCER SCREENING PROGRAMME PATIENTS WITH SYMPTOMATIC PATIENTS REFERRED UNDER 2 WEEK WAIT IN A DEMOGRAPHICALLY MATCHED POPULATION

doi:10.1136/gut.2011.239301.240

P Harrow,* S Nayagam, N Stafford, K Hopkins, R Loke *Gastroenterology, Kent and Sussex Hospital, Tunbridge Wells, UK*

Introduction Traditionally the majority of colorectal cancer has been diagnosed following investigations triggered by symptoms suggestive of malignancy. The National Bowel Cancer Screening Programme has created an alternative route to diagnosis of asymptomatic individuals.

Methods This prospective study compared the incidence of colorectal cancer and adenomatous polyps at colonoscopy and virtual colonoscopy in a population demographically matched for age (60–70) and geographical area (limited to 33 local GP practices). Subjects were either asymptomatic under the Bowel Cancer Screening Programme (BCSP) or symptomatic patients referred under the 2 week wait target (2ww). All endoscopy took place at the Kent & Sussex Hospital between August 2009 and August 2010. Data from colonoscopy reports and histology were analysed.

Results 155 BCSP (mean age 66.0, 61% male) and 127 2ww patients (mean age 64.6, 40% male) fulfilled the entry criteria. Sub-group analysis of both gender groups also found a significant difference in the incidence of adenomatous polyps and no significant difference in the incidence of malignancy.

Table 1 PTU-112 Findings at endoscopy

	BCSP (155)	2 ww (127)	
Cancer	11%	7.1%	p>0.05
-Duke A-B	59%	33%	p>0.05
Polyps	47%	30%	p<0.005
-% Adenomas	79%	47%	p<0.005

Conclusion This study demonstrates comparable incidence of colorectal cancers in the two groups. The incidence of polyps was significantly higher in the BCSP group and importantly the polyps found in BCSP were of higher risk of progression to cancer according to the classical adenoma-carcinoma sequence.¹ This study demonstrates the efficacy of the BCSP at detecting significant colorectal pathology. The detection of colorectal cancer and high risk polyps are thought to reduce mortality from colorectal cancer.²

The similar detection rates between the symptomatic and asymptomatic groups highlights the poor positive predictive value of symptoms and the assessment currently used for emergency referrals for the diagnosis of colorectal cancer.

Competing interests None.

Keywords bowel cancer screening, colonic polyps, colorectal cancer.

REFERENCES

1. Fearon ER, Vogelstein B. A genetic model for colorectal tumorigenesis. *Cell*. 1990 Jun 1;61(5):759–767.
2. Mandel, Church, Bond *et al.* *N Engl J Med*. 1993 May 328(19):1365–7.