PTU-020

**DOES BOWEL CANCER SCREENING DETECT
COLORECTAL CANCERS AT AN EARLIER STAGE?**

doi:10.1136/gut.2011.239301.148

P Patel,^{1*} T Pirani,¹ R Leicester,¹ M Bouwer,¹ A Poullis¹ ¹Endoscopy Unit, St. George's Hospital, London, UK

Introduction The NHS bowel cancer screening program (NHS BCSP) currently screens persons between the ages of 60 and 69 for bowel cancer with the aim to detect early cancer and reduce colorectal cancer (CRC) mortality. The inclusion age is currently being extended to 74. The NHS BCSP began in 2006 in England and Nationwide coverage was achieved in 2010. Individuals are invited biennially to provide a stool sample for faecal occult blood test (FOBT) analysis. Bowel cancer usually presents once symptoms occur and as a result at an advanced stage (Duke's stage at diagnosis A 11% vs C/D 54%¹). Pilot studies of the NHS BCSP showed a reduction in the stage of cancer at diagnosis (Duke's stage at diagnosis (42% stage A, C/D 28.2%²). This abstract examines whether the introduction of the NHS BCSP has a real-life effect on stage of CRC at diagnosis in patients from a large screening centre.

Table 1 PTU-020

	BCSP patients	Non-BCSP patients
Total number of cancers	134	456 (3 synchronous)
Duke's Stage A	58 (43%)	58 (13%)
Duke's Stage B	24 (18%)	172 (38%)
Duke's Stage C	36 (27%)	171 (38%)
Duke's Stage D	16 (12%)	55 (12%)

Methods Data from BCSP patients from January 2007 to December 2009 were obtained from the database at the South West London BCS centre based at St. Georges'. Data on non-BCSP patients from January 2003 to December 2007 was obtained from local cancer network databases.

Results The table below shows the numbers of cancers and their stage at diagnosis for both patient groups.

A higher proportion of Duke's A CRC are diagnosed in patients through the BCSP (43% vs 13%; $p < 0.05$ χ^2 test). 26 cancers in the BCSP were in fact polyp cancers which have been included in the Duke's A group.

Conclusion Our results are in keeping with previous pilot data² and a previously presented abstract³ and show that the BCSP detects CRC at a relatively earlier stage compared to non-BCSP patients. This study needs to be repeated at intervals to see whether this difference changes with further BCSP rounds.

Competing interests None.

Keywords Bowel cancer screening, colorectal cancer.

REFERENCES

1. Improving Outcome in Colorectal Cancers: Manual update. *NICE*. June 2004
2. Evaluation of the UK colorectal Screening Pilot, Final Report. February 2003
3. Lee TJW, Zwart C, Rees C, *et al*. Reduced stage at diagnosis of colorectal cancer in the NHS Bowel Cancer Screening Programme. *Gut* 2009;58(Suppl II):A36.