PWE-065 SHOULD FOCAL COLONIC ACTIVITY FOUND ON ¹⁸FDG PET CT BE FOLLOWED-UP WITH ENDOSCOPIC EVALUATION?

doi:10.1136/gutjnl-2012-302514d.65

¹C L Y Wong,* ¹W J Jafar, ²B Rajashanker, ¹R C Prudham, ²M Prescott, ³P A Hulse, ¹J K Limdi. ¹Department of Gastroenterology, Pennine Acute Hospitals NHS Trust, Manchester, UK; ²Department of Radiology, Central Manchester Hospitals NHS Trust, Greater Manchester, Manchester, UK; ³Department of Radiology, The Christie NHS Foundation Trust, Manchester, UK

Introduction PET CT is a non-invasive imaging modality that is widely used in the imaging work up of malignant disease. It has been postulated that due to ¹⁸FDG accumulation in adenomatous polyps, PET CT using ¹⁸FDG may detect early premalignant colonic lesions. The aim of our study was to assess the performance of ¹⁸FDG PET CT in the detection of significant colonic lesions namely adenomas or villous adenomas >1 cm in size and to assess whether endoscopy in all patients with focal FDG colonic uptake on PET CT is justified by the detection rate.

Methods A retrospective review of patients who had ¹⁸FDG PET CT at the Pennine Acute Hospitals NHS Trust, Manchester as part of their staging for various cancers between February 2007 and October 2011 was conducted by a review of the reports of the examinations.

Results 1553 patients between the ages of 17 and 93 years (median age 55 years) had ¹⁸FDG PET CT performed during this period. Increased focal FDG uptake in the colon was detected in 179 of 1553 patients studied (11.5%). Endoscopic evaluation was performed in 71 (39.7%) of these patients, flexible sigmoidoscopy in 25 (35.2%) and colonoscopy in 46 (64.8%). Non-malignant pathology was found in 45 of 179 patients (25.1%); 17 (9.5%) were tubulovillous adenomas; 6 (3.4%) had villous adenomas; 9 (5.0%) were tubular adenomas; 4 (2.2%) hyperplastic polyps; 11 (6.1%) were normal colonic tissue and 1 (0.6%) was active inflammatory bowel disease. From the patients with increased ¹⁸FDG uptake in the colon, 19 (10.6%) were subsequently found to have adenocarcinoma on histology. This represents 26.8% of the 71 patients who received endoscopic evaluation.

Conclusion The presence of focal colonic ¹⁸FDG uptake on PET CT scan justifies endoscopic evaluation in all patients where treatment would be clinically appropriate.

Competing interests None declared.

PWE-066 MAGNETIC RESONANCE ENTEROGRAPHY FOR THE ASSESSMENT OF CROHN'S DISEASE: CHANGING IMAGING PARADIGMS?

doi:10.1136/gutjnl-2012-302514d.66

¹K A Mcwhirter,* ¹J Limdi, ²R Filobos. ¹Department of Gastroenterology, Pennine Acute Hospitals, Manchester, UK; ²Department of Radiology, Pennine Acute Hospitals, Manchester, UK

Introduction Recent advances in the immunopathogenesis and therapy of inflammatory bowel disease (IBD) coupled with bolder definitions of disease control have led to increasing reliance on imaging. Increased awareness of the potential downstream effects of ionising radiation has placed more emphasis on radiation-free imaging. We aimed to assess the role of magnetic resonance enterography (MRE) in assessing Crohn's disease.

Methods We conducted a retrospective review of 141 consecutive MRE studies performed between June 2009 and November 2010. Clinical data were obtained from electronic patient record review. Inflammatory markers, radiological investigations and ileocolonoscopy when performed within 90 days of MR enterography were recorded. MRE reports were recorded using accepted activity criteria- small bowel dilatation, stenosis, wall thickening, enhance-

ment, mucosal irregularity, mesenteric inflammation, hyper-vascularity, lymph node enlargement, abscesses, fistulation and extraintestinal features.

Results Of 67 patients with IBD, 60 had Crohn's disease and 59 examinations were complete. Thirty- nine of 67 patients were female, mean age 34 (range 16–68) and median disease duration of 5 years (range 0–39). Abnormalities were noted in 47 MRE scans; 34 had active non-stricturing, 12 active stricturing and one fibrostenotic disease. Within the active groups, there were four fistulae and three abscesses in four patients. Ileo-colonoscopy was performed in 14 of these patients with 12/14 showing active colitis and raised CRP in 11/16 within 90 days of MRE. Treatment was increased in 47% of the active non-stricturing group, 3/16 to azathioprine, 8/16 to infliximab, 4/16 to surgery with no change to treatment in the remaining 53%, of whom 4/8 had normal ileo-colonoscopy and 13/16 normal CRP. In 83% of active stricturing patient treatment was increased, one to azathioprine, six to biologics, three to surgery. Four of 12 patients in this group had an elevated CRP and 4/7 had active colitis at ileo-colonoscopy. Of 12 normal MRE, treatment was not increased in 92%. Of these, CRP was normal in 10/11 and ileo-colonoscopy normal in 4/7. Treatment was increased in one to biologics, with an elevated CRP and moderately severe colitis at colonoscopy. The fibrostenotic subject had normal CRP and mild colitis at colonoscopy, and proceeded to surgery. All the abscess/fistula subjects had raised inflammatory markers. Two were referred for surgery, one started biologics and one treated with antibiotics.

Conclusion The small bowel remains difficult to assess endoscopically. MRE adds to the assessment of patients with Crohn's disease, in addition to endoscopy and biological markers identifying patients with active disease where treatment escalation may result in meaningful benefit.

Competing interests None declared.

PWE-067 MRI TO DISTINGUISH HIDRADENITIS SUPPURATIVA FROM PERIANAL CROHN'S DISEASE

doi:10.1136/gutjnl-2012-302514d.67

¹N Griffin, ²K M Taylor,* ³A B Williams, ²S H Anderson, ²P M Irving, ²J D Sanderson, ⁴N Desai, ¹V Goh. ¹Department of Radiology, Guy's & St Thomas' NHS Foundation Trust, London, UK; ²Department of Gastroenterology, Guy's & St Thomas' NHS Foundation Trust, London, UK; ³Department of Colorectal Surgery, Guy's & St Thomas' NHS Foundation Trust, London, UK; ⁴Department of Dermatology, Guy's & St Thomas' NHS Foundation Trust, London, UK

Introduction Hidradenitis suppurativa is a chronic inflammatory disease of apocrine gland-bearing skin. It has a predilection for the ano-genital skin and is characterised clinically by inflammatory abscesses and sinus tract formation. Clinically, the condition can mimic cutaneous Crohn's disease, and furthermore it may co-exist with Crohn's disease.¹ MRI is often used to assess perianal Crohn's disease and has been comprehensively described in the literature, however to date, there has only been one case report to describe the MRI features of perianal hidradenitis suppurativa.²

Methods The electronic patient records, and radiology and inflammatory bowel disease databases were searched for patients with hidradenitis suppurativa who had undergone short-tau inversion recovery (STIR) MR imaging to define disease severity, assess for fistulating disease and exclude Crohn's disease, between 2005 and 2012.

Results A total of 188 patients with hidradenitis suppurativa were identified (129 female, six with concomitant IBD), of whom 18 (15 male, three with concomitant IBD) had undergone perianal MRI. On baseline MRI, multiple tracts were typically seen in the natal cleft (15/18), the perianal (11/18), the perineal (10/18) and gluteal (8/18) regions. Abscesses were most commonly gluteal (3/18). The