REVIEW OF CT COLONOGRAPHY: REAL-LIFE EXPERIENCE OF ONE THOUSAND CASES IN A TERTIARY REFERRAL CENTRE

A E Smyth, C F Healy, P MacMathuna, H M Fenlon *Departments of Gastroenterology and Radiology, Mater Misericordiae University Hospital, Dublin, Ireland*

10.1136/gutjnl-2013-305143.36

Introduction CT Colonography is increasingly being utilised to investigate patients with lower gastrointestinal symptoms and in screening for colorectal carcinoma in the at risk population.

Aims/Background This review aimed to quantify results from a real life experience of 1000 CT Colonography examinations in a tertiary referral hospital campus.

Method Data on patient demographics, indications and findings were analysed retrospectively from a dedicated database.

Results Over a five year period (January 2008–December 2012) 1000 CT Colonography examinations were performed. The median age was 70 years (IQR 60–79). Patients were symptomatic in 86% of cases, screening accounted for 7% and surveillance in patients with a history of polyps or colorectal cancer for 7%. In 45% of patients it was documented that the study was performed following an incomplete optical colonoscopy.

CT Colonography had normal or benign colonic findings in 75% of patients. 6% had incomplete examinations performed. Neoplasia was observed in 14%: colorectal carcinoma 6%, polyps >1cm in 3% and 5% were found to have <3 polyps of 6–9 mm. 5% had findings of wall thickening or strictures that were suspected to be benign but required further endoscopy and biopsy.

There was an extracolonic finding that required further evaluation in 19% with 6% found to have a potentially clinically significant finding.

There were no significant procedure related complications.

Conclusion This review describes the real life experience of CT Colonography in a largely elderly, symptomatic patient group. In this patient group, CT Colonography safely confirmed the need for no further bowel tests in 75%, identified colorectal cancer in 6% and clinically significant extracolonic findings in 6% of cases without any significant complications.