OPEN ACCESS

ORIGINAL ARTICLE

α -Conotoxin Vc1.1 inhibits human dorsal root ganglion neuroexcitability and mouse colonic nociception via GABA_B receptors

Joel Castro,¹ Andrea M Harrington,¹ Sonia Garcia-Caraballo,¹ Jessica Maddern,¹ Luke Grundy,¹ Jingming Zhang,² Guy Page,² Paul E Miller,² David J Craik,³ David J Adams,⁴ Stuart M Brierley¹

► Additional material is published online only. To view please visit the journal online (<http://dx.doi.org/10.1136/gutjnl-2015-310971>).

For numbered affiliations see end of article.

Correspondence to

Professor Stuart M Brierley, Visceral Pain Group, Centre for Nutrition and Gastrointestinal Diseases, The University of Adelaide, Level 7, South Australian Health and Medical Research Institute (SAHMRI), North Terrace, Adelaide, SA 5000, Australia; stuart.brierley@adelaide.edu.au

JC, AMH and SG-C contributed equally.

Received 27 October 2015
Revised 22 December 2015
Accepted 14 January 2016
Published Online First
17 February 2016

ABSTRACT

Objective α -Conotoxin Vc1.1 is a small disulfide-bonded peptide from the venom of the marine cone snail *Conus victoriae*. Vc1.1 has antinociceptive actions in animal models of neuropathic pain, but its applicability to inhibiting human dorsal root ganglion (DRG) neuroexcitability and reducing chronic visceral pain (CVP) is unknown.

Design We determined the inhibitory actions of Vc1.1 on human DRG neurons and on mouse colonic sensory afferents in healthy and chronic visceral hypersensitivity (CVH) states. In mice, visceral nociception was assessed by neuronal activation within the spinal cord in response to noxious colorectal distension (CRD). Quantitative-reverse-transcription-PCR, single-cell-reverse-transcription-PCR and immunohistochemistry determined γ -aminobutyric acid receptor B (GABA_BR) and voltage-gated calcium channel (Ca_v2.2, Ca_v2.3) expression in human and mouse DRG neurons.

Results Vc1.1 reduced the excitability of human DRG neurons, whereas a synthetic Vc1.1 analogue that is inactive at GABA_BR did not. Human DRG neurons expressed GABA_BR and its downstream effector channels Ca_v2.2 and Ca_v2.3. Mouse colonic DRG neurons exhibited high GABA_BR, Ca_v2.2 and Ca_v2.3 expression, with upregulation of the Ca_v2.2 exon-37a variant during CVH. Vc1.1 inhibited mouse colonic afferents ex vivo and nociceptive signalling of noxious CRD into the spinal cord in vivo, with greatest efficacy observed during CVH. A selective GABA_BR antagonist prevented Vc1.1-induced inhibition, whereas blocking both Ca_v2.2 and Ca_v2.3 caused inhibition comparable with Vc1.1 alone.

Conclusions Vc1.1-mediated activation of GABA_BR is a novel mechanism for reducing the excitability of human DRG neurons. Vc1.1-induced activation of GABA_BR on the peripheral endings of colonic afferents reduces nociceptive signalling. The enhanced antinociceptive actions of Vc1.1 during CVH suggest it is a novel candidate for the treatment for CVP.

Significance of this study**What is already known on this subject?**

- Patients with IBS suffer from chronic visceral pain (CVP); however, there are limited analgesic therapeutic options currently available for treatment.
- A rich source of novel agents to treat chronic pain is the α -conotoxin family of peptides from the venom of marine cone snails.
- α -Conotoxin Vc1.1 has antinociceptive and antihyperalgesic actions in neuropathic pain models; however, its ability to inhibit human sensory dorsal root ganglion (DRG) neurons remains unknown.
- Vc1.1's applicability in reducing CVP is also unknown.

What are the new findings?

- Vc1.1 reduces human sensory DRG neuroexcitability, via a γ -aminobutyric acid receptor B (GABA_BR)-mediated mechanism.
- We show that human DRG neurons express GABA_BR and the voltage-gated calcium channels Ca_v2.2, and Ca_v2.3, which are the direct and downstream targets of Vc1.1, respectively.
- Vc1.1 inhibits mouse colonic nociceptors and also low-threshold distension-sensitive colonic afferents with greatest effect during chronic visceral hypersensitivity (CVH).
- Peripheral in vivo Vc1.1 administration inhibits the signalling of noxious information from the colon to the spinal cord. This antinociceptive effect is also greater in mice with CVH.
- During CVH, mouse colonic DRG neurons show upregulation of the Ca_v2.2 exon-37a variant, which may explain the increased inhibitory effect of Vc1.1 in CVH states.

INTRODUCTION

IBS is a prevalent, chronic GI disorder that negatively impacts the quality of life in up to 14% of the population.^{1,2} It is characterised by abdominal pain and discomfort associated with altered bowel habits.^{3–5} Although the pathophysiology of IBS is not completely understood, it is becoming clear

that changes to peripheral cellular and sensory mechanisms play key roles in the associated pain.^{6,7} In particular, chronic visceral hypersensitivity (CVH) of colonic afferents is implicated in the development and maintenance of chronic visceral

CrossMark

To cite: Castro J, Harrington AM, Garcia-Caraballo S, et al. *Gut* 2017;**66**:1083–1094.

Significance of this study

How might it impact on clinical practice in the foreseeable future?

- ▶ Vc1.1 has been tested in human clinical trials for the treatment of neuropathic pain, where it has been demonstrated to have a clean safety and side-effect profile.
- ▶ Our current findings show Vc1.1 inhibits human DRG neurons, via activation of the GABA_BR, which is a key finding for clinical translatability.
- ▶ This inhibitory effect in human neurons combined with the enhanced antinociceptive action of Vc1.1 in colonic pathways in a mouse model of CVH suggests it is a novel candidate for the treatment for CVP associated with IBS.
- ▶ We show that by reducing nociceptive signalling from the periphery, Vc1.1 has potential therapeutic value in treating CVP.
- ▶ These findings put GABA_BR agonists in the spotlight as potential peripheral agents for the treatment of CVP.

pain (CVP) in IBS.^{4–5} Characteristic features of CVH include nociceptor hypersensitivity⁸ and increased signalling of noxious colorectal distension (CRD) within the spinal cord.^{9–11} Recent evidence suggests sensory afferents display upregulation of numerous ion channels and receptors in animal models of CVH,^{7–10–12} making them targets for analgesic treatment.

A recently introduced treatment for patients with IBS and constipation involves a small disulfide-rich peptide that is restricted to the GI tract, where it inhibits peripheral nociceptive pathways and produces clinically relevant pain relief.⁹ Given the limited treatments available for patients with other subtypes of IBS, additional analgesic therapeutic options are needed. A rich source of novel small disulfide-rich agents comes from the α -conotoxin family of peptides from the venom of marine cone snails.¹³ These peptides target a wide variety of membrane receptors and ion channels.¹⁴ In particular α -conotoxin Vc1.1, a 16-amino acid synthetic version of a peptide derived from the marine cone snail *Conus victoriae*, has antinociceptive actions in vitro and antihyperalgesic actions in numerous in vivo neuropathic pain models.^{15–17} Interestingly, in a chronic constriction injury model of neuropathic pain, Vc1.1 relieves tactile allodynia.¹⁷ These inhibitory effects were similar to those obtained with gabapentin, a ligand recently proposed as a potential IBS therapeutic,¹⁸ but were achieved at far lower doses.¹⁷ Notably, Vc1.1 (also called ACV1) has been used in phase I and phase IIA clinical trials for the treatment of neuropathic pain.^{19–21} These studies showed Vc1.1 was safe and well tolerated with a clean safety and side-effect profile. Despite this promise, therapeutic trials were discontinued as Vc1.1 was shown to be less potent at the human $\alpha 9\alpha 10$ nicotinic acetylcholine receptor (nAChR), which was thought to mediate the inhibitory action of Vc1.1. However, more recent recombinant cell line studies have clearly demonstrated that the human γ -aminobutyric acid receptor B (GABA_BR) is the primary and high affinity target for Vc1.1.^{17–22–25–27} These studies also demonstrated GABA_BR activation by Vc1.1 causes downstream inhibition of the voltage-gated calcium channels Ca_v2.2 and Ca_v2.3, which underlies Vc1.1's inhibitory actions.^{14–28} These recent findings are intriguing; as both oral and intravenous administration of baclofen, the archetypal GABA_BR agonist has been shown to reduce the

pseudo-affective responses to CRD in animal models.^{29–30} Although it is unclear if this baclofen-induced inhibition is centrally or peripherally mediated, we wondered if Vc1.1 represents a novel peripheral gut analgesic for the treatment of CVP. Therefore, we determined if Vc1.1 inhibits human sensory dorsal root ganglion (DRG) neurons, the primary transducers at the start of the pain-processing pathway. Second, we determined if Vc1.1 inhibits sensory pathways within the splanchnic and pelvic innervation of the colon and whether these actions are enhanced in an animal model of CVH. Third, we determined if the inhibitory actions of Vc1.1 are mediated via activation of GABA_BR on the peripheral endings of colonic afferents.

MATERIALS AND METHODS

For comprehensive descriptions of the methodologies used, see the online supplementary information.

Human DRG

Thoracolumbar (TL) DRG (T9–L1) were acquired from five (three female, two male) human adult organ donors (22.2 $\pm$ 2.08 years of age) during the removal of the vital organs for transplantation. The harvested DRG were immediately processed for downstream patch clamp or RNA studies. Intact DRG were kept for quantitative-reverse-transcription-PCR (qRT-PCR) mRNA expression studies from each spinal level (T9, T10, T11, T12, L1) while additional DRG were dissociated to allow individual DRG neurons to be studied with single-cell-reverse-transcription-PCR (RT-PCR) studies, or to allow patch clamp recordings to be performed.

Human DRG patch clamp recordings

Whole-cell patch clamp recordings of cultured human DRG neurons were performed in current clamp mode in response to depolarising current pulses (20 or 50 pA current steps, 500 ms duration). This allowed the rheobase (amount of current needed to initiate action potential generation) to be assessed in the presence and absence of Vc1.1 (1000 nM) and a synthetic analogue of Vc1.1 ([P6O]Vc1.1; 1000 nM), which is inactive at GABA_BR. An increased rheobase indicates more current is required to fire an action potential and therefore the neuron displays reduced excitability.

Mouse model of CVH

Intracolonic trinitrobenzene-sulfonic acid (TNBS) was administered as described previously.^{8–10} TNBS-treated mice were allowed to recover for 28 days, at which stage inflammation had resolved and chronic colonic afferent mechanical hypersensitivity was evident.^{8–10–12}

Ex vivo electrophysiology

Recordings of splanchnic and pelvic afferents were made from healthy control and CVH mice as described previously.^{8–10} Briefly, colonic nociceptors were recorded from the splanchnic pathway. They respond to noxious distension (40 mm Hg), stretch (≥ 7 g) or von Frey hair filaments (2 g)^{8–31} and become mechanically hypersensitive in models of CVP.^{8–10–12} Muscular-mucosal afferents were recorded from the pelvic pathway and respond to both low-intensity circular stretch (<5 g) and fine mucosal stroking (10 mg).^{8–31–33} Once baseline responses had been established, mechanosensitivity was retested after application of Vc1.1 (1, 10, 100, 1000 nM) for 10 min at each dose. To determine the mechanism of action of Vc1.1 the selective GABA_BR antagonist (CGP55845; 5 μ M), Ca_v2.2 blocker (ω -conotoxin CVID; 1 μ M) or Ca_v2.3 blocker

(SNX-482:200 nM) were applied alone, or in combination, at maximally effective concentrations for 10 min prior to incubation with Vc1.1 (1000 nM).

CRD and pERK immunohistochemistry

Healthy control or CVH mice received an intracolonic enema of either saline or Vc1.1 (1000 nM). Ten minutes later, under anaesthesia, a 4 cm CRD balloon catheter was inserted transanally into healthy or CVH mice.^{9–11} After regaining consciousness CRD was performed (80 mmHg for 10 s, deflated for 5 s, repeated five times). Following sacrifice via anaesthetic overdose, mice underwent fixation by transcardial perfusion and the TL (T10–L1) and lumbosacral (LS:L6–S1) spinal cord removed and cryoprotected. Frozen sections were cut and incubated with monoclonal rabbit anti-phosphorylated MAP kinase ERK 1/2 (pERK) and AlexaFluor-488 was used for visualisation.^{9–11}

Isolation of mouse colonic DRG neurons

TL and LS DRG were removed from healthy control and CVH mice 4 days after retrograde tracing from the colon with AlexaFluor-555-conjugated cholera-toxin subunit-B (CTB-AF555). DRG were dissociated and single or pooled colonic DRG neurons isolated for downstream mRNA expression analysis.¹⁰

Quantitative-reverse-transcription-PCR

RNA was extracted from either whole human DRG and single human DRG neurons or mouse whole DRG, pooled colonic DRG neurons and single colonic DRG neurons from healthy control and CVH mice using specific isolation kits. QRT-PCR was performed using either human-specific or mouse-specific primers for GABA_BR1, GABA_BR2, Ca_v2.2 and Ca_v2.3.^{10 12 32 34} The comparative cycle threshold method was used to quantify the abundance of target transcripts to reference genes.^{10 12 32 34}

Immunohistochemistry

In both perfused-fixed frozen mouse DRG sections and dissociated mouse DRG neurons specific antibodies for GABA_BR1, GABA_BR2, Ca_v2.2 or Ca_v2.3 were used to determine the expression of these targets in retrogradely traced colonic DRG neurons. Antibody preabsorption and omission of primary antibodies were used as controls (see online supplementary figure S1). AlexaFluor-488 or AlexaFluor-594 conjugated secondary antibodies were used for visualisation.

RESULTS

Vc1.1 reduces the excitability of human DRG neurons

To determine if Vc1.1 reduces the excitability of human DRG neurons we used whole-cell patch clamp recordings to assess neuronal excitability. Vc1.1 (1000 nM) inhibited a specific population (40%) of human DRG neurons, which was indicated by a significant increase in the amount of injected current required to fire an action potential (figure 1Ai). In this population of neurons, Vc1.1 increased the rheobase by 20% compared with control responses (figure 1Aii, B and see online supplementary figure S2). The average of cell capacitance for all the recorded human DRG neurons was 131.48 ± 18.03 pF, with no significant difference in cell capacitance observed between neurons which were affected by Vc1.1 and those that were not.

To determine if this inhibition was mediated via GABA_BR or $\alpha 9\alpha 10$ -nAChR we used a modified version of Vc1.1 ([P6O] Vc1.1), which is inactive at the GABA_BR, but active at the $\alpha 9\alpha 10$ -nAChR.³⁵ [P6O]Vc1.1 had no effect on human DRG

neuronal excitability (figure 1C), suggesting Vc1.1 exerts its inhibitory effects on human DRG neurons via a GABA_BR mechanism. Recent recombinant cell line studies have demonstrated that the human GABA_BR is the high affinity (nanomolar) target for Vc1.1 and that GABA_BR activation by Vc1.1 causes downstream inhibition of the voltage-gated calcium channels Ca_v2.2 and Ca_v2.3.¹⁴ In order to determine if the same mechanism applied in human DRG neurons, we determined the expression of GABA_BR and Ca_v channels in whole TL DRG from five spinal levels from four human adult donors. We showed that subunits R1 and R2 for GABA_BR were expressed as well as Ca_v2.2 and Ca_v2.3 (figure 1D). Expression levels for each of the targets were remarkably consistent between the different DRG levels across the four human donors (figure 1D). Single-cell-RT-PCR of individual human TL DRG neurons demonstrated that 46% coexpressed GABA_BR and Ca_v2.2, the minimum components required for Vc1.1-induced inhibition (figure 1E). This was consistent with our patch clamp observations where 40% of the human DRG neurons tested were inhibited by Vc1.1. Overall, these functional and expression studies indicate Vc1.1 inhibits human DRG neurons via a GABA_BR-mediated mechanism.

Vc1.1 inhibits mouse colonic afferents with greater efficacy in CVH

Given Vc1.1's inhibitory actions on human DRG neurons in the current study and rat somatosensory neurons in previous studies,^{16 35 36} we hypothesised Vc1.1 may also inhibit colonic afferents. To test this hypothesis we performed in vitro single-unit colonic afferent recordings.^{9 10 32 33} Specifically, we recorded from mouse splanchnic nerves, which supply the mid-to-distal colon and signal predominantly nociceptive information,^{8 33} and the pelvic nerves supplying the colorectum, which signal a mixture of physiological and nociceptive information.^{8 33} Vc1.1 significantly and dose-dependently decreased healthy colonic nociceptor activity, with a maximum reduction in response to mechanical stimulation of 32% at the highest concentration tested (figure 2Ai). We then asked if these Vc1.1-induced antinociceptive effects were maintained, or indeed augmented, in CVH. This question was assessed in a mouse model where colonic nociceptor mechanical hypersensitivity^{7–10} and colonic mechanical hyperalgesia and allodynia are evident long after resolution of TNBS-induced colitis.^{7 37 38} We found that colonic nociceptors in the CVH model displayed pronounced hypersensitivity and that Vc1.1 significantly reduced nociceptor mechanosensitivity, showing significant reductions at 100 nM and 1000 nM, with a maximal reduction of 44% at 1000 nM (figure 2Aii). Overall, Vc1.1's inhibitory effect was greatly enhanced in CVH nociceptors compared with healthy nociceptors (figure 2B, C). We also tested whether the inhibitory effects of Vc1.1 extended to low-threshold distension sensitive pelvic afferents and found that Vc1.1 dose-dependently inhibited pelvic muscular–mucosal afferent responses to circular stretch in healthy mice (figure 2Di, Fi). The inhibitory effect of Vc1.1 on pelvic afferents was also enhanced in afferents from CVH mice (figure 2Dii, E, F).

Vc1.1 reduces in vivo processing of noxious CRD in the mouse TL and LS spinal cord

Vc1.1 inhibits mouse colonic nociceptors in the splanchnic pathway and low-threshold distension sensitive afferents in the pelvic pathway. We therefore hypothesised these actions should correspondingly reduce signalling of noxious CRD relayed into the TL and LS spinal cord in vivo. In response to noxious CRD,

Figure 1 α -Conotoxin Vc1.1 inhibits human dorsal root ganglion (DRG) neurons. (A) (i) Group data showing that Vc1.1 (1000 nM) significantly increases the rheobase of a subpopulation (40%) of human DRG neurons, indicating Vc1.1 inhibits neuronal excitability and more current is required to initiate an action potential (** $p < 0.001$, $n = 10$, paired t-test). (ii) In this population of neurons, Vc1.1 increased the rheobase by 20% compared with baseline response, meaning the neurons are less excitable (** $p < 0.001$). (B) Representative examples of human DRG neuronal responses in the absence (control solutions) and in the presence of Vc1.1. Note in this example more current is required to fire an action potential from a human DRG neuron in the presence Vc1.1 (1970 pA) relative to control (1540 pA). (C) [P6O]Vc1.1 (1000 nM), a synthetic analogue of Vc1.1 that does not act at γ -aminobutyric acid receptor B (GABA_BR), did not affect human DRG neuronal excitability when expressed as either i) rheobase or ii) % of rheobase ($p > 0.05$, $n = 10$, paired t-test) indicating Vc1.1 induces its inhibitory effect via the GABA_BR. (D) Group data of quantitative reverse-transcription-PCR analysis from thoracolumbar (T9, T10, T11, T12, L1) DRG from four human adult donors indicating expression of the GABA_BR subunits R1, R2 and the voltage-gated calcium channels Ca_v2.2 and Ca_v2.3 in human DRG. (E) (i) Examples of gel electrophoresis following single-cell-PCR analysis from individual human DRG neurons. (ii) Combined analysis of expression and coexpression of GABA_BR and Ca_v channels in 39 human DRG neurons. Of human DRG neurons, 46.2% (18/39) coexpress GABA_BR and Ca_v2.2, the minimum components required for Vc1.1-induced inhibition. Combined these studies indicate Vc1.1 inhibits human DRG neurons via a GABA_BR-mediated mechanism. GAPDH, glyceraldehyde-3-phosphate dehydrogenase.

Figure 2 α -Conotoxin Vc1.1 inhibits colonic afferents in ex vivo recordings from healthy and chronic visceral hypersensitivity (CVH) mice. (A) (i) Vc1.1 significantly inhibits splanchnic colonic nociceptors from healthy mice. Compared with baseline, Vc1.1 at 1000 nM significantly reduced colonic nociceptor mechanosensitivity ($^{**}p < 0.01$, $n = 10$ afferents, one-way ANOVA, Bonferroni post hoc). (ii) In a model of CVH, nociceptors are potently and concentration-dependently inhibited by Vc1.1, with significant reductions in mechanical responses at 100 nM and 1000 nM ($^{**}p < 0.01$, $n = 10$ afferents, one-way ANOVA, Bonferroni post hoc tests). (B) Change in mechanosensitivity induced by Vc1.1 in healthy and CVH nociceptors compared with their respective baseline responses. Vc1.1 caused significantly more inhibition at 100 nM ($^{**}p < 0.01$) and 1000 nM ($^{*}p < 0.05$) in CVH nociceptors than healthy nociceptors (healthy: $n = 10$; CVH: $n = 10$, two-way ANOVA, Bonferroni post hoc). (C) Single-unit recordings from the splanchnic innervation showing inhibition of (i) a healthy nociceptor, and (ii) a CVH nociceptor following application of Vc1.1 (1000 nM). (D) (i) Vc1.1 inhibited pelvic low-threshold muscular-mucosal afferents from healthy mice. Compared with baseline, significant reductions in the mechanosensory response evoked by 5 g circular stretch were observed at 100 nM and 1000 nM Vc1.1 ($^{*}p < 0.05$, $n = 6$ afferents). (ii) Low-threshold muscular-mucosal afferents from CVH mice were also concentration-dependently inhibited by Vc1.1, with significant reductions at 100 nM ($^{*}p < 0.05$) and 1000 nM ($^{**}p < 0.01$, $n = 6$ afferents). (E) Compared with their respective baseline responses, Vc1.1 causes significantly more inhibition of muscular-mucosal afferents ($^{*}p < 0.05$ at 1000 nM) from CVH mice ($n = 6$) relative to healthy ($n = 6$) mice. (F) Single-unit recordings from the pelvic innervation showing inhibition of (i) a healthy low-threshold muscular-mucosal afferent and (ii) a CVH muscular-mucosal afferent following application of Vc1.1 (1000 nM).

pERK-immunoreactivity (pERK-IR) identifies activated neurons in the dorsal horn (DH) of the spinal cord.^{9–11} In healthy mice given noxious CRD, prior intracolonic administration of 1000 nM Vc1.1 resulted in significantly fewer pERK-IR DH neurons in both the TL and LS spinal cord (figure 3A–C).

In response to noxious CRD, CVH mice displayed greater numbers of pERK-IR DH neurons than healthy mice, which corresponds with the mechanical hypersensitivity of colonic nociceptors observed in our afferent recording studies. CVH mice pretreated with intracolonic Vc1.1 displayed significantly reduced numbers of pERK-IR DH neurons in both the TL and LS spinal cord following noxious CRD (figure 3D,E), with the extent of inhibition greater within the TL pathway (figure 3D). Overall, these results suggest Vc1.1 reduces the signalling of

noxious stimuli from the colon and reduces the CVH observed in vivo.

Vc1.1-induced inhibition of mouse colonic afferents is mediated via the GABA_BR

To elucidate the site and mechanism of action of Vc1.1 in colonic pathways, we first used [P6O]Vc1.1, which is inactive at GABA_BR. As per our recordings in human DRG neurons, [P6O] Vc1.1 did not inhibit mouse colonic nociceptors (figure 4A), suggesting that the inhibitory effects of Vc1.1 on colonic afferents are mediated via GABA_BR.

We then asked if the archetypal GABA_BR agonist, baclofen, inhibited colonic nociceptors. Baclofen caused a dose-dependent inhibition of colonic nociceptors from both healthy (figure 4Bi)

Figure 3 Intracolonic administration of Vc1.1 reduces nociceptive signalling in the dorsal horn (DH) of the spinal cord in response to noxious colorectal distension (CRD). (A) Noxious CRD (80 mm Hg) in healthy mice results in activation of DH neurons in the thoracolumbar (T10–L1; splanchnic innervation) and lumbosacral (L6–S1; pelvic innervation) spinal cord, as indicated by pERK-immunoreactivity (pERK-IR). Mice pretreated with intracolonic Vc1.1 (1000 nM) display significantly fewer DH neurons in the thoracolumbar spinal cord, specifically T11–T12 (** $p < 0.01$) and T13–L1 (** $p < 0.01$), and the lumbosacral spinal cord (** $p < 0.001$, one-way ANOVA, $n = 6$: healthy+saline, $n = 6$: healthy+1000 nM Vc1.1). (B) Schematic representation of laminae I–V (LI–LV) in the DH of the spinal cord. (C) (i) Healthy thoracolumbar DH. Left panel: following noxious CRD, pERK-IR (yellow arrows) neurons were predominantly located in the superficial DH (laminae I) and laminae V. Right panel: in healthy mice pretreated with Vc1.1 (1000 nM) fewer pERK-IR neurons are evident following noxious CRD. (ii) Healthy lumbosacral DH. Left panel: following noxious CRD, pERK-IR (yellow arrows) neurons were located in laminae I, II, IV and V. Right panel: healthy mice pretreated with Vc1.1 (1000 nM) displayed fewer pERK-IR neurons following noxious CRD, particularly within laminae I. (D) In chronic visceral hypersensitivity (CVH) mice, more neurons are activated by noxious CRD at baseline in the thoracolumbar DH. Pretreatment with intracolonic Vc1.1 (1000 nM) significantly reduces the number of pERK-IR DH neurons within the T10–T11 (* $p < 0.05$), T11–T12 (** $p < 0.01$), T13–L1 (** $p < 0.01$) and lumbosacral DH (** $p < 0.01$; CVH +intracolonic saline: $n = 6$, CVH+intracolonic Vc1.1: $n = 6$). (E) (i) Left panel: in CVH mice, following noxious CRD, pERK-IR neurons in the thoracolumbar DH were predominantly located in laminae I–II and throughout laminae III–V. Right panel: CVH mice pretreated with Vc1.1 (1000 nM) display fewer pERK-IR neurons following noxious CRD, particularly in the superficial laminae. (ii) CVH mice pretreated with intracolonic Vc1.1 (1000 nM) display fewer pERK-IR neurons in the lumbosacral DH.

and CVH (figure 4Bii) mice. Interestingly, and similarly to Vc1.1, baclofen also inhibited CVH colonic nociceptors to a greater degree (figure 4C). To confirm that inhibition of colonic nociceptors by Vc1.1 was mediated by the GABA_BR, we first administered the selective GABA_BR antagonist CGP55845. In the presence of CGP55845, Vc1.1 no longer inhibited colonic nociceptors from either healthy (figure 4D) or CVH (figure 4E) mice. Finally, we confirmed the expression of GABA_BR subunits GABA_BR1 and GABA_BR2 in colonic DRG neurons by using immunohistochemistry. More than 80% of colonic DRG neurons expressed both GABA_BR1 and GABA_BR2 subunits (figure 5A and see online supplementary figure S3). Taken together, these data suggest that the antinociceptive action of Vc1.1 on colonic afferents is mediated via GABA_BR expressed on colonic afferents.

Coexpression of GABA_BR and Ca_v2.2 and Ca_v2.3 in mouse colonic DRG neurons

Recent studies in mammalian cell lines show that Vc1.1-induced inhibition via the GABA_BR also requires the presence of either Ca_v2.2^{35–39} or Ca_v2.3.⁴⁰ To examine whether these channels contribute to the Vc1.1-induced inhibition of colonic afferents, we determined their expression profile in both the TL (splanchnic) and LS (pelvic) pathways innervating the colon. Using qRT-PCR, we found that GABA_BR1, GABA_BR2, Ca_v2.2 and Ca_v2.3 were abundantly expressed in both mouse TL and LS DRG (see online supplementary figure S4). As colonic DRG neurons represent only approximately 5% of the neurons in these ganglia, we performed retrograde tracing to identify colonic innervating neurons.^{10–12–34} Single-cell-RT-PCR analysis and

Figure 4 The inhibitory effect of Vc1.1 on mouse colonic afferents is mediated via the γ -aminobutyric acid receptor B ($GABA_B$ R). (A) The modified peptide (P60)Vc1.1, which is inactive at the $GABA_B$ R, does not inhibit colonic nociceptors from chronic visceral hypersensitivity (CVH) mice (*Not Significant* (NS), $n=6$ afferents, paired t-test). (B) (i) Application of the $GABA_B$ R agonist baclofen caused dose-dependent inhibition of healthy colonic nociceptors, with significant reductions in mechanosensitivity observed at 20 μ M ($*p<0.05$) and 200 μ M baclofen ($***p<0.001$), respectively. (ii) Similarly, in colonic nociceptors from CVH mice, baclofen caused significant inhibition at 2 μ M ($*p<0.05$), 20 μ M ($***p<0.001$) and 200 μ M ($***p<0.001$), respectively. (C) Change in mechanosensitivity induced by baclofen in healthy and CVH nociceptors, compared with their respective baseline responses. Baclofen caused significantly more inhibition at 20 μ M ($*p<0.05$) and 200 μ M ($**p<0.01$) in CVH nociceptors compared with healthy nociceptors (healthy: $n=7$; CVH: $n=6$, two-way ANOVA, Bonferroni post hoc). (D) (i) A single dose of Vc1.1 (1000 nM) caused significant inhibition of colonic nociceptors from healthy mice ($***p<0.001$, $n=10$, paired t-test). (ii) Prior incubation with the selective $GABA_B$ R antagonist CGP-55845 (5 μ M) prevented the Vc1.1-induced inhibition of healthy colonic nociceptors (NS, $n=5$, one-way ANOVA). (E) (i) CVH colonic nociceptors were also inhibited by a single high dose (1000 nM) of Vc1.1 ($***p<0.001$, $n=7$, paired t-test). (ii) Prior incubation of CGP-55845 (5 μ M) also prevented the Vc1.1-induced inhibition of CVH nociceptors (NS; $n=9$, one-way ANOVA). ANOVA, analysis of variance.

immunohistochemistry determines the expression and importantly the coexpression of these targets specifically within colonic DRG neurons (figure 5B–D, see online supplementary figures S5 and S6). Immunohistochemistry demonstrated that the vast majority of colonic DRG neurons express $Ca_v2.2$ (see online supplementary figure S5) and $Ca_v2.3$ (see online supplementary figure S6). Single-cell-RT-PCR confirmed the majority of colonic DRG neurons expressed $GABA_B$ R, $Ca_v2.2$ and $Ca_v2.3$ (figure 5Ci,ii), with 85% of colonic DRG neurons from healthy and CVH mice coexpressing high levels of $GABA_B$ R and $Ca_v2.2$ (figure 5Di), with 80% coexpressing all three targets, $GABA_B$ R, $Ca_v2.2$ and $Ca_v2.3$ (figure 5Dii).

Recent studies indicate there are two isoforms of $Ca_v2.2$: exon-37a and exon-37b.^{41 42} Using isoform-specific primers and qRT-PCR from pooled colonic DRG neurons, we found a significant increase in the $Ca_v2.2$ -exon-37a variant in CVH mice in both TL and LS pathways (figure 5E). This upregulation may explain the increased efficacy of Vc1.1 in CVH states.

$Ca_v2.2$ and $Ca_v2.3$ contribute to Vc1.1-mediated inhibition of mouse colonic nociceptors

Recombinant cell line studies indicate Vc1.1-mediated activation of $GABA_B$ R results in the downstream inhibition of both $Ca_v2.2$ and $Ca_v2.3$ via second messenger pathways.^{35 40} To determine how Vc1.1 inhibits colonic nociceptors, we hypothesised blocking $Ca_v2.2$ and $Ca_v2.3$, either alone or in combination with maximally effective concentrations of toxin blockers, should also inhibit mouse colonic nociceptors. Using either a selective $Ca_v2.2$ (ω -conotoxin CVID) or $Ca_v2.3$ (SNX-482) blocker inhibited

healthy nociceptors (figure 6Ai,Bi, see online supplementary figures S7A and S8A) and caused greater inhibition of CVH nociceptors (figure 6Aii,Bii, see online supplementary figures S7B and S8B). In separate experiments a combination of CVID and SNX-482 caused pronounced inhibition of healthy nociceptors (figure 6Ci, see online supplementary figure S9A) and even greater inhibition of CVH nociceptors (figure 6Cii, see online supplementary figure S9B). Application of Vc1.1 in the presence of both CVID and SNX-482 had little additional inhibitory effects in both states (figure 6Ci,ii, see online supplementary figures S9A and S9B). Overall, these findings suggest Vc1.1-induced activation of $GABA_B$ R results in the downstream blockade of $Ca_v2.2$ and $Ca_v2.3$, which inhibits colonic nociceptor excitability (figures 6D and 7).

DISCUSSION

This study provides evidence that the α -conotoxin Vc1.1 inhibits human DRG neurons via activation of the $GABA_B$ R. It also demonstrates that the peripheral administration of Vc1.1 in mice strongly inhibits the processing of nociceptive information within colonic sensory pathways. We show that both human DRG neurons and mouse colonic DRG neurons express the molecular targets of Vc1.1, the $GABA_B$ R and its downstream effector channels $Ca_v2.2$ and $Ca_v2.3$. Correspondingly, we show that Vc1.1 inhibits colonic afferents in both the splanchnic and pelvic pathways and that blocking $Ca_v2.2$ and $Ca_v2.3$ causes inhibition comparable with that of Vc1.1 alone. These findings highlight the potential therapeutic value of Vc1.1 in the treatment of CVP.

Figure 5 Colonic dorsal root ganglion (DRG) neurons express γ -aminobutyric acid receptor B (GABA_B) subunits and the voltage-gated calcium channels Ca_v2.2 and Ca_v2.3. (A) Immunohistochemistry for (i) GABA_B1 and (iii) GABA_B2 in frozen sections of thoracolumbar DRG from mice that had previously undergone colonic retrograde tracing with CTB-555. A high percentage of traced colonic DRG neurons from both healthy and chronic visceral hypersensitivity (CVH) mice express (ii) GABA_B1 and (iv) GABA_B2, respectively (healthy: n=6; CVH: n=5). (B) In separate experiments healthy and CVH mice underwent retrograde tracing from the colon with CTB-555. After 4 days thoracolumbar DRG neurons were dissociated and individual colonic DRG neurons were isolated for single-cell-PCR analysis. Gel electrophoresis indicates individual colonic DRG neurons from healthy and CVH mice and their respective expression of GABA_B1, GABA_B2, Ca_v2.2 and Ca_v2.3. Bath controls, perfusate collected during the isolation of single cells, show no expression of any of the targets or reference genes. (C) A high proportion of colonic DRG neurons from (i) healthy and (ii) CVH mice express mRNA for GABA_B1, GABA_B2, Ca_v2.2 and Ca_v2.3. (D) (i) Coexpression of Ca_v2.2 and GABA_B mRNA is found in the majority (>85%) of thoracolumbar colonic DRG neurons from healthy and CVH mice. (ii) mRNA for the GABA_B, Ca_v2.2 and Ca_v2.3 are coexpressed in the majority (80%) of colonic thoracolumbar DRG neurons from healthy and CVH mice. (E) Quantitative-reverse-transcription-PCR from isolated and pooled (200) colonic DRG neurons shows (i) a significant upregulation of the Ca_v2.2 exon-37a splice variant in both thoracolumbar and lumbosacral colonic DRG neurons from CVH mice (*p<0.05; healthy: n=4; CVH: n=4). (ii) There is no overall change in total Ca_v2.2 (exon-37a+37b) levels.

Figure 6 Vc1.1-induced inhibition can be replicated by blocking both $\text{Ca}_v2.2$ (CVID) and $\text{Ca}_v2.3$ (SNX-482). (A) (i) In ex vivo preparations colonic nociceptors from healthy mice are inhibited following incubation of the $\text{Ca}_v2.2$ antagonist CVID (1 μM), although this effect is not significant, whereas (ii) chronic visceral hypersensitivity (CVH) colonic nociceptors are significantly inhibited by CVID (* $p < 0.05$, $n = 10$). In both healthy and CVH colonic nociceptors, the subsequent application of Vc1.1 (1000 nM) in the presence of CVID caused further inhibition (healthy: * $p < 0.05$, $n = 10$; CVH: ** $p < 0.01$, $n = 10$). (B) The $\text{Ca}_v2.3$ blocker SNX-482 (200 nM) inhibited both (i) healthy (* $p < 0.05$, $n = 7$) and (ii) CVH (* $p < 0.05$, $n = 5$) splanchnic colonic nociceptor mechanosensitivity. In both healthy and CVH states the subsequent application of Vc1.1 (1000 nM) in the presence of SNX-482 (200 nM) caused further inhibition of healthy colonic nociceptors (** $p < 0.01$, $n = 7$). (C) The combined application of the $\text{Ca}_v2.2$ and $\text{Ca}_v2.3$ blockers, CVID (1 μM) and SNX-482 (200 nM), respectively, significantly inhibits both (i) healthy (* $p < 0.05$, $n = 8$) and (ii) CVH (** $p < 0.001$, $n = 6$) colonic nociceptors with subsequent application of Vc1.1 (1000 nM) causing little additional inhibition. (D) Inhibition of (i) healthy and (ii) CVH colonic nociceptors by single doses of Vc1.1 (1000 nM), CVID (1 μM) and SNX-482 (200 nM) or in the presence of a combination of CVID (1 μM) and SNX-482 (200 nM) expressed as percentage inhibition from healthy nociceptor baseline. CVID causes significantly lesser inhibition than Vc1.1 in (i) healthy (* $p < 0.05$) and (ii) CVH (** $p < 0.01$) nociceptors. However, blocking both $\text{Ca}_v2.2$ and $\text{Ca}_v2.3$ in combination with CVID (1 μM) and SNX-482 (200 nM) causes similar inhibition to Vc1.1 alone.

Vc1.1 inhibits sensory DRG neurons, which are the primary transducers of nociceptive information at the start of the pain pathway

A crucial finding of this study was Vc1.1's ability to inhibit a subpopulation of human DRG neurons. This indicates Vc1.1 has an antinociceptive effect in these neurons, which is a key discovery for clinical translatability. These findings were complemented with animal studies where we observed significant Vc1.1-induced inhibition of both colonic nociceptors and low-threshold stretch sensitive afferents that can encode into the noxious range. Crucially, we found that these antinociceptive

actions were augmented in a mouse model of CVH. These ex vivo findings translate in vivo as, in response to noxious CRD, mice administered intracolonic Vc1.1 have reduced numbers of activated DH neurons within the TL and LS spinal cord. This finding indicates, in the presence of Vc1.1, a reduced capacity to detect and signal nociceptive events from the colon into the central nervous system. In particular, we observed fewer activated neurons within the superficial lamina of the DH. This is the major termination zone for nociceptive afferents and consists of nociception-specific neurons. Importantly, our findings suggest that Vc1.1 reverses the chronic visceral mechanical

Figure 7 Proposed antinociceptive mechanism by which α -conotoxin Vc1.1 activates γ -aminobutyric acid receptor B (GABA_B). Vc1.1 activates GABA_B expressed on human dorsal root ganglion (DRG) neurons and on colonic afferent endings resulting in $\text{G}\beta\gamma$ and c-Src kinase-mediated inhibition of $\text{Ca}_v2.2$ and $\text{Ca}_v2.3$ and subsequent inhibition of neuronal excitability. Modified from Adams et al.¹⁴

hypersensitivity evident in our ex vivo and in vivo studies, rather than completely blocking nociceptive responses. This is important as ideal analgesic agents reverse pathological pain, rather than removing protective pain signalling completely.

Vc1.1 activates GABA_BR on human DRG neurons and on mouse colonic afferents to inhibit nociceptive signalling

In this study, we have demonstrated for the first time that Vc1.1 inhibits human DRG neuronal excitability via GABA_BR activation. This was evident as [P6O]Vc1.1, which is inactive at GABA_BR, does not alter neuronal excitability. Similarly, [P6O] Vc1.1 did not affect mouse colonic afferents, whereas the selective GABA_BR antagonist CGP55845 prevented Vc1.1-induced inhibition of colonic nociceptors, both in healthy and CVH states. To confirm our proposal that Vc1.1 acts via GABA_BR, we used baclofen, the archetypal GABA_BR agonist, and showed that it also inhibits colonic nociceptors. Notably, this inhibitory effect was greater during CVH. The significance of this finding is fourfold. First, these findings closely match those with Vc1.1 and conclusively demonstrate that activation of GABA_BR on the peripheral endings of colonic DRG neurons within the colon wall results in nociceptor inhibition. Second, although it is known that baclofen inhibits vagal afferents in the upper gut,⁴³ and low-threshold distension sensitive pelvic colonic afferents,⁴⁴ it has not been previously shown to inhibit colonic nociceptors, or afferents in a model of CVH. Third, in rats, both oral and intravenous administration of baclofen reduces visceral pain-related pseudo-affective responses to CRD.^{29–30} Fourth, baclofen also reduces colonic inflammation-induced neuronal activation within the spinal cord and the brainstem.⁴⁵ Overall, these observations are consistent with our current in vitro, ex vivo and in vivo findings on the antinociceptive actions of Vc1.1. In response to noxious colonic stimuli, we show inhibition of both colonic nociceptors and low-threshold afferents and a reduction in neuronal activation to noxious CRD in the TL and LS DH of healthy and CVH mice.

Although GABA_BR have been localised within the rat and human GI tract,^{46–47} crucially we demonstrate for the first time, definitive expression of both GABA_BR subunits in human DRG neurons and in colonic DRG neurons from healthy and CVH mice. Taken together these studies demonstrate that activation of GABA_BR on human DRG neurons reduces nociceptive signalling, while activation of GABA_BR on the peripheral endings of colonic afferents reduces nociception and visceral pain in both healthy and hyperalgesic states. These are important findings and complement studies in other fields of neuroscience, whereby in pyramidal neurons in the cortex, somatic and dendritic GABA_B receptors regulate neuronal excitability via different mechanisms.⁴⁸ Specifically, these studies show that activation of somatic GABA_B receptors leads to a reduction in neuronal output, primarily by increasing the rheobase, whereas activation of dendritic GABA_B receptors blocks burst firing, decreasing action potential output.⁴⁸ Our studies recording from the soma of DRG neurons and primary afferent endings in the colon support these mechanisms, where we have observed Vc1.1 increasing the action potential rheobase and decreasing action potential output, respectively.

Vc1.1 as a novel antinociceptive peptide for the treatment of CVP

Although we have shown that the overall antinociceptive effect induced by baclofen and Vc1.1 are similar, it is clear from other studies that Vc1.1 and baclofen act via different mechanisms, in terms of their binding to GABA_BR and also their downstream

targets. For example, Vc1.1 does not compete with baclofen for binding at the 'Venus fly trap' on the GABA_BR, but is proposed to target the interface between the GABA_BR subunit ectodomains (figure 7).⁴⁹ Furthermore, whereas baclofen is able to inhibit several different neuronal calcium channels, including Ca_v2.1, Ca_v2.2 and Ca_v2.3, and activate G-protein-coupled inwardly rectifying potassium channels (GIRK) channels, Vc1.1 is more specific by only acting via Ca_v2.2 or Ca_v2.3.^{26–35–39–40} Because baclofen crosses the blood–brain barrier, some of its previously reported antinociceptive effects may be mediated centrally.⁵⁰ This presents a problem in terms of its off-target effects, which include centrally mediated neurological side-effects, including dizziness.⁵¹ In contrast, we show that peripheral administration of Vc1.1 ex vivo and in vivo reduces nociceptive signalling, suggesting a peripheral mechanism of action. Furthermore, as Vc1.1 is a peptide, if delivered peripherally it is unlikely to cross the blood–brain barrier and therefore may be less likely to cause central side-effects. Notably, Vc1.1 has been tested in human clinical trials for treatment of neuropathic pain.^{19–21} However, its development was discontinued due to lack of potency at its (at the time) proposed molecular target, the human $\alpha 9\alpha 10$ -nAChR.⁵² The emergence of an action mediated via the human GABA_BR suggests its development for chronic pain treatment could resume. Given our current finding of an enhanced Vc1.1 antinociceptive action during CVH, we suggest it is a novel candidate for the treatment for CVP, particularly as cyclised versions of Vc1.1 have impressive stability and are resistant to proteolysis.¹⁷

Human DRG neurons and mouse colonic DRG neurons express Ca_v2.2 and Ca_v2.3, the key downstream targets of GABA_BR activation by Vc1.1

The GABA_BR is a G-protein-coupled receptor and therefore must couple to downstream channels to exert its inhibitory actions. Previous studies demonstrate Vc1.1-mediated activation of GABA_BR is coupled to both Ca_v2.2 and Ca_v2.3 via G $\beta\gamma$ and c-Src kinase second messenger systems.^{35–40} Ca_v2.2 channels mediate the neuronal N-type calcium current,⁴⁹ whereas Ca_v2.3 channels typically conduct a small proportion of the whole-cell calcium current, known as the R-type calcium current.⁵³ Inhibition of Ca_v2.2, by either gene knockout or selective channel antagonists, causes analgesia in neuropathic pain models.⁴⁹ Notably, Ca_v channels have been demonstrated in several studies to contribute to the rheobase of neurons.^{54–55} While Ca_v channels have been identified as potential therapeutic targets for treating neuropathic pain,⁵⁶ little is known about the roles of Ca_v2.2 and Ca_v2.3 in visceral pain. Here we show that the vast majority of colonic DRG neurons express mRNA and protein for GABA_BR, Ca_v2.2 and Ca_v2.3. Notably, in colonic DRG neurons from CVH mice, we identified a significant upregulation of the Ca_v2.2 variant exon-37a. Although the Ca_v2.2 exon-37a variant is expressed at relatively low levels, this is an important finding as exon-37a has been reported to be highly expressed in nociceptors, where it acts as part of a molecular switch controlling N-type current density and G-protein-mediated voltage-independent inhibition.^{41–42–57} Accordingly, upregulation of this variant may contribute to the processes responsible for mechanical hypersensitivity and neuronal hyperexcitability in CVH. Furthermore, this upregulation may help to explain the increased inhibitory effects of both Vc1.1 and the Ca_v2.2 antagonist, ω -conotoxin CVID, in nociceptors from CVH mice. The high expression levels for Ca_v2.3 in colonic DRG neurons and a considerable inhibitory action of SNX-482 on colonic nociceptors suggests a key role for Ca_v2.3

in colonic pain pathways. Although $\text{Ca}_v2.3$ channels are present in some somatic nociceptors⁵⁸ and can contribute to somatic pain behaviour via spinal and supraspinal mechanisms,^{59–60} our findings are consistent with previous studies suggesting lower $\text{Ca}_v2.3$ expression in DRG neurons that innervate the epidermis than those innervating deep tissues.⁶¹ As we could not specifically identify colonic innervating DRG neurons from human donors, this may explain why we observed less expression of $\text{Ca}_v2.3$ in whole human DRG and less human DRG neurons expressing $\text{Ca}_v2.3$.

In conclusion, our findings demonstrate an antinociceptive action for Vc1.1 in human DRG neurons and in colonic sensory afferents. This antinociceptive action is stronger in a model of CVH than in healthy mice and is mediated via the G-protein-coupled receptor, GABA_BR , which is abundantly expressed in human DRG neurons and mouse colonic DRG neurons. Because altered visceral sensory function is a hallmark of IBS, Vc1.1 represents a potential novel therapy to reduce nociceptive stimuli from the colon and rectum to the central nervous system. Our findings highlight the potential therapeutic value of Vc1.1 and its optimised analogues⁶² and also identifies the GABA_BR as a potential target for the treatment of CVP associated with GI disorders.

Author affiliations

¹Visceral Pain Group, Centre for Nutrition and Gastrointestinal Diseases, Discipline of Medicine, Faculty of Health Sciences, The University of Adelaide, South Australian Health and Medical Research Institute (SAHMRI), Adelaide, South Australia, Australia

²Anabios, San Diego, California, USA

³Division of Chemistry and Structural Biology, Institute for Molecular Bioscience, University of Queensland, Brisbane, Queensland, Australia

⁴Illawarra Health & Medical Research Institute (IHMRI), University of Wollongong, Wollongong, NSW, Australia

Acknowledgements Dr Andrea Ghetti (Anabios) is acknowledged for discussions on the human patch clamp recordings.

Contributors SMB, DJA and DJC conceived the study. SMB, DJA, AMH, LG, GP, JZ, PEM and JC designed, conducted and analysed experiments. SG-C and JM also conducted and analysed experiments. DJC synthesised Vc1.1 and associated analogues and assisted with critical revision of the manuscript for important intellectual content. SMB wrote the paper and all authors helped with revising the manuscript.

Funding This work was funded by the National Health and Medical Research Council (NHMRC) of Australia Project Grant #1049928 awarded to DJA, SMB and DJC. AMH received funding via the Australian Research Council (ARC) Discovery Early Career Research Award. DJC is an NHMRC Senior Principal Research Fellow. DJA is an ARC Australian Professorial Fellow. SMB is an NHMRC RD Wright Biomedical Research Fellow.

Competing interests None declared.

Patient consent Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

Data sharing statement We have included all data relevant to the submission in the manuscript and the supplementary information.

Open Access This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

REFERENCES

- Hungin AP, Chang L, Locke GR, *et al.* Irritable bowel syndrome in the United States: prevalence, symptom patterns and impact. *Aliment Pharmacol Ther* 2005;21:1365–75.
- Drossman DA, Camilleri M, Mayer EA, *et al.* AGA technical review on irritable bowel syndrome. *Gastroenterology* 2002;123:2108–31.
- Azpiroz F, Bouin M, Camilleri M, *et al.* Mechanisms of hypersensitivity in IBS and functional disorders. *Neurogastroenterol Motil* 2007;19:62–88.
- Lembo T, Munakata J, Mertz H, *et al.* Evidence for the hypersensitivity of lumbar splanchnic afferents in irritable bowel syndrome. *Gastroenterology* 1994;107:1686–96.
- Ritchie J. Pain from distention of the pelvic colon by inflating a balloon in the irritable bowel syndrome. *Gut* 1973;6:105–12.
- Barbara G, Stanghellini V, De Giorgio R, *et al.* Activated mast cells in proximity to colonic nerves correlate with abdominal pain in irritable bowel syndrome. *Gastroenterology* 2004;126:693–702.
- Brierley SM, Linden DR. Neuroplasticity and dysfunction after gastrointestinal inflammation. *Nat Rev Gastroenterol Hepatol* 2014;11:611–27.
- Hughes PA, Brierley SM, Martin CM, *et al.* Post-inflammatory colonic afferent sensitisation: different subtypes, different pathways and different time courses. *Gut* 2009;58:1333–41.
- Castro J, Harrington AM, Hughes PA, *et al.* Linaclotide inhibits colonic nociceptors and relieves abdominal pain via guanylate cyclase-C and extracellular cyclic guanosine 3',5'-monophosphate. *Gastroenterology* 2013;145:1334–46.e11-11.
- de Araujo AD, Mobli M, Castro J, *et al.* Selenoether oxytocin analogues have analgesic properties in a mouse model of chronic abdominal pain. *Nat Commun* 2014;5:3165.
- Harrington AM, Brierley SM, Isaacs N, *et al.* Sprouting of colonic afferent central terminals and increased spinal mitogen-activated protein kinase expression in a mouse model of chronic visceral hypersensitivity. *J Comp Neurol* 2012;520:2241–55.
- Hughes PA, Castro J, Harrington AM, *et al.* Increased κ -opioid receptor expression and function during chronic visceral hypersensitivity. *Gut* 2014;63:1199–200.
- Schroeder CI, Craik DJ. Therapeutic potential of conopeptides. *Future Med Chem* 2012;4:1243–55.
- Adams DJ, Callaghan B, Berecki G. Analgesic conotoxins: block and G protein-coupled receptor modulation of the N-type ($\text{Ca}_v2.2$) calcium channels. *Br J Pharmacol* 2012;166:486–500.
- Klimis H, Adams DJ, Callaghan B, *et al.* A novel mechanism of inhibition of high-voltage activated calcium channels by α -conotoxins contributes to relief of nerve injury-induced neuropathic pain. *Pain* 2011;152:259–66.
- Satkunanathan N, Livett B, Gayler K, *et al.* Alpha-conotoxin Vc1.1 alleviates neuropathic pain and accelerates functional recovery of injured neurones. *Brain Res* 2005;1059:149–58.
- Clark RJ, Jensen J, Nevin ST, *et al.* The engineering of an orally active conotoxin for the treatment of neuropathic pain. *Angew Chem Int Ed Engl* 2010;49:6545–8.
- Gale JD, Houghton LA. Alpha-2-delta ($\alpha(2)\delta$) ligands, gabapentin and pregabalin: what is the evidence for potential use of these ligands in irritable bowel syndrome. *Front Pharmacol* 2011;2:28.
- ASX. Metabolic pharmaceuticals: metabolics neuropathic pain drug, ACV1- clinical trials update. 2006. <http://www.asx.com.au/asxpdf/20061129/pdf/3zv2c96tyh1nx.pdf>.
- ASX. Metabolic pharmaceuticals: metabolics neuropathic pain drug ACV1-additional preclinical studies reveal greater potential. 2006. <http://www.asx.com.au/asxpdf/20061123/pdf/3zqm91n1jhpff.pdf>.
- ASX. Metabolic pharmaceuticals: metabolic discontinues clinical trial programme for neuropathic pain drug, ACV1. 2007. <http://www.asx.com.au/asxpdf/20070814/pdf/313yjpg7j4lg.pdf>.
- Halai R, Clark RJ, Nevin ST, *et al.* Scanning mutagenesis of alpha-conotoxin Vc1.1 reveals residues crucial for activity at the $\alpha 9\alpha 10$ nicotinic acetylcholine receptor. *J Biol Chem* 2009;284:20275–84.
- Napier IA, Klimis H, Rycroft BK, *et al.* Intrathecal α -conotoxins Vc1.1, AulB and MII acting on distinct nicotinic receptor subtypes reverse signs of neuropathic pain. *Neuropharmacology* 2012;62:2202–7.
- van Lierop BJ, Robinson SD, Kompella SN, *et al.* Dicarba α -conotoxin Vc1.1 analogues with differential selectivity for nicotinic acetylcholine and GABAB receptors. *ACS Chem Biol* 2013;8:1815–21.
- Yu R, Kompella SN, Adams DJ, *et al.* Determination of the α -conotoxin Vc1.1 binding site on the $\alpha 9\alpha 10$ nicotinic acetylcholine receptor. *J Med Chem* 2013;56:3557–67.
- Callaghan B, Adams DJ. Analgesic α -conotoxins Vc1.1 and RglA inhibit N-type calcium channels in sensory neurons of $\alpha 9$ nicotinic receptor knockout mice. *Channels (Austin)* 2010;4:51–4.
- Sharpe IA, Gehrmann J, Loughnan ML, *et al.* Two new classes of conopeptides inhibit the $\alpha 1$ -adrenoceptor and noradrenaline transporter. *Nat Neurosci* 2001;4:902–7.
- Berecki G, McArthur JR, Cuny H, *et al.* Differential Cav2.1 and Cav2.3 channel inhibition by baclofen and α -conotoxin Vc1.1 via GABAB receptor activation. *J Gen Physiol* 2014;143:465–79.
- Brusberg M, Ravnefjord A, Martinsson R, *et al.* The GABA(B) receptor agonist, baclofen, and the positive allosteric modulator, CGP7930, inhibit visceral pain-related responses to colorectal distension in rats. *Neuropharmacol* 2009;56:362–7.
- Lindström E, Brusberg M, Ravnefjord A, *et al.* Oral baclofen reduces visceral pain-related pseudo-affective responses to colorectal distension in rats: relation between plasma exposure and efficacy. *Scand J Gastroenterol* 2011;46:652–62.

- 31 Brierley SM, Page AJ, Hughes PA, *et al.* Selective role for TRPV4 ion channels in visceral sensory pathways. *Gastroenterology* 2008;134:2059–69.
- 32 Brierley SM, Hughes PA, Page AJ, *et al.* The ion channel TRPA1 is required for normal mechanosensation and is modulated by algescic stimuli. *Gastroenterology* 2009;137:2084–2095.e3.
- 33 Brierley SM, Jones RC III, Gebhart GF, *et al.* Splanchnic and pelvic mechanosensory afferents signal different qualities of colonic stimuli in mice. *Gastroenterology* 2004;127:166–78.
- 34 Hughes PA, Harrington AM, Castro J, *et al.* Sensory neuro-immune interactions differ between irritable bowel syndrome subtypes. *Gut* 2013;62:1456–65.
- 35 Callaghan B, Haythornthwaite A, Berecki G, *et al.* Analgesic alpha-conotoxins Vc1.1 and Rg1A inhibit N-type calcium channels in rat sensory neurons via GABA_BR activation. *J Neurosci* 2008;28:10943–51.
- 36 Lang PM, Burgstahler R, Haberberger RV, *et al.* A conus peptide blocks nicotinic receptors of unmyelinated axons in human nerves. *Neuroreport* 2005;16:479–83.
- 37 Adam B, Liebrechts T, Gschossmann JM, *et al.* Severity of mucosal inflammation as a predictor for alterations of visceral sensory function in a rat model. *Pain* 2006;123:179–86.
- 38 Gschossmann JM, Liebrechts T, Adam B, *et al.* Long-term effects of transient chemically induced colitis on the visceromotor response to mechanical colorectal distension. *Dig Dis Sci* 2004;49:96–101.
- 39 Cuny H, de Faoite A, Huynh TG, *et al.* γ -Aminobutyric acid type B (GABAB) receptor expression is needed for inhibition of N-type (Cav2.2) calcium channels by analgesic α -conotoxins. *J Biol Chem* 2012;287:23948–57.
- 40 Berecki G, McArthur JR, Cuny H, *et al.* Differential Cav2.1 and Cav2.3 channel inhibition by baclofen and α -conotoxin Vc1.1 via GABABR activation. *J Gen Physiol* 2014;143:465–79.
- 41 Bell TJ, Thaler C, Castiglioni AJ, *et al.* Cell-specific alternative splicing increases calcium channel current density in the pain pathway. *Neuron* 2004;41:127–38.
- 42 Raingo J, Castiglioni AJ, Lipscombe D. Alternative splicing controls G protein-dependent inhibition of N-type calcium channels in nociceptors. *Nat Neurosci* 2007;10:285–92.
- 43 Page AJ, Blackshaw LA. GABA_BR inhibit mechanosensitivity of primary afferent endings. *J Neurosci* 1999;19:8597–602.
- 44 Sengupta JN, Medda BK, Shaker R. Effect of GABA(B) receptor agonist on distension-sensitive pelvic nerve afferent fibers innervating rat colon. *Am J Physiol Gastrointest Liver Physiol* 2002;283:1343–51.
- 45 Lu Y, Westlund KN. Effects of baclofen on colon inflammation-induced Fos, CGRP and SP expression in spinal cord and brainstem. *Brain Res* 2001;889:118–30.
- 46 Nakajima K, Tooyama I, Kuriyama K, *et al.* Immunohistochemical demonstration of GABABR in the rat gastrointestinal tract. *Neurochem Res* 1996;21:211–15.
- 47 Uezono Y, Kaibara M, Hayashi H, *et al.* Characterization of GABABR in the human colon. *J Pharmacol Sci* 2004;94:211–13.
- 48 Breton JD, Stuart GJ. Somatic and dendritic GABA(B) receptors regulate neuronal excitability via different mechanisms. *J Neurophysiol* 2012;108:2810–18.
- 49 Adams DJ, Berecki G. Mechanisms of conotoxin inhibition of N-type (Ca(v)2.2) calcium channels. *Biochim Biophys Acta* 2013;1828:1619–28.
- 50 Davis MP. Drug management of visceral pain: concepts from basic research. Pain research and treatment. Hindawi Publishing Corporation, 2012.
- 51 Hyland NP, Cryan JF. A gut feeling about GABA: focus on GABA(B) receptors. *Front Pharmacol* 2010;1:124.
- 52 McIntosh JM, Absalom N, Chebib M, *et al.* Alpha9 nicotinic acetylcholine receptors and the treatment of pain. *Biochem Pharmacol* 2009;78:693–702.
- 53 Schneider T, Dibué M, Hescheler J. How “Pharmaco-resistant” is Cav2.3, the major component of voltage-gated R-type Ca²⁺ channels. *Pharmaceuticals (Basel)* 2013;6:759–76.
- 54 Hogan Q, Lirk P, Poroli M, *et al.* Restoration of calcium influx corrects membrane hyperexcitability in injured rat dorsal root ganglion neurons. *Anesth Analg* 2008;107:1045–51.
- 55 Lirk P, Poroli M, Rigaud M, *et al.* Modulators of calcium influx regulate membrane excitability in rat dorsal root ganglion neurons. *Anesth Analg* 2008;107:673–85.
- 56 Zamponi GW, Striessnig J, Koschak A, *et al.* The physiology, pathology, and pharmacology of voltage-gated calcium channels and their future therapeutic potential. *Pharmacol Rev* 2015;67:821–70.
- 57 Altier C, Dale CS, Kisilevsky AE, *et al.* Differential role of N-type calcium channel splice isoforms in pain. *J Neurosci* 2007;27:6363–73.
- 58 Fang Z, Park CK, Li HY, *et al.* Molecular basis of Ca(v)2.3 calcium channels in rat nociceptive neurons. *J Biol Chem* 2007;282:4757–64.
- 59 Saegusa H, Kurihara T, Zong S, *et al.* Altered pain responses in mice lacking alpha 1E subunit of the voltage-dependent Ca²⁺ channel. *Proc Natl Acad Sci USA* 2000;97:6132–7.
- 60 Terashima T, Xu Q, Yamaguchi S, *et al.* Intrathecal P/Q- and R-type calcium channel blockade of spinal substance P release and c-Fos expression. *Neuropharmacology* 2013;75:1–8.
- 61 Yang FC, Tan T, Huang T, *et al.* Genetic control of the segregation of pain-related sensory neurons innervating the cutaneous versus deep tissues. *Cell Rep* 2013;5:1353–64.
- 62 Yu R, Seymour VA, Berecki G, *et al.* Less is more: design of a highly stable disulfide-deleted mutant of analgesic cyclic α -conotoxin Vc1.1. *Sci Rep* 2015;5:13264.