



OPEN ACCESS

ORIGINAL ARTICLE

Outcome measures in coeliac disease trials: the Tampere recommendations

Jonas F Ludvigsson,^{1,2} Carolina Ciacci,³ Peter HR Green,⁴ Katri Kaukinen,^{5,6} Ilma R Korponay-Szabo,^{7,8} Kalle Kurppa,^{9,10} Joseph A Murray,¹¹ Knut Erik Aslaksen Lundin,^{12,13} Markku J Maki,^{14,15} Alina Popp,^{16,17} Norelle R Reilly,^{18,19} Alfonso Rodriguez-Herrera,²⁰ David S Sanders,²¹ Detlef Schuppan,^{22,23} Sarah Sleet,²⁴ Juha Taavela,²⁵ Kristin Voorhees,²⁶ Marjorie M Walker,²⁷ Daniel A Leffler²⁸

► Additional material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/gutjnl-2017-314853>).

For numbered affiliations see end of article.

Correspondence to

Dr Jonas F Ludvigsson, Department of Medical Epidemiology and Biostatistics, Karolinska Institutet, Stockholm 171 77, Sweden; jonasludvigsson@yahoo.com

Received 14 July 2017

Revised 19 December 2017

Accepted 8 January 2018

Published Online First

13 February 2018

ABSTRACT

Objective A gluten-free diet is the only treatment option of coeliac disease, but recently an increasing number of trials have begun to explore alternative treatment strategies. We aimed to review the literature on coeliac disease therapeutic trials and issue recommendations for outcome measures.

Design Based on a literature review of 10 062 references, we (17 researchers and 2 patient representatives from 10 countries) reviewed the use and suitability of both clinical and non-clinical outcome measures. We then made expert-based recommendations for use of these outcomes in coeliac disease trials and identified areas where research is needed.

Results We comment on the use of histology, serology, clinical outcome assessment (including patient-reported outcomes), quality of life and immunological tools including gluten immunogenic peptides for trials in coeliac disease.

Conclusion Careful evaluation and reporting of outcome measures will increase transparency and comparability of coeliac disease therapeutic trials, and will benefit patients, healthcare and the pharmaceutical industry.

INTRODUCTION

Coeliac disease (CD) is an immune-mediated disease triggered by gluten exposure.¹ Although characterised by small intestinal inflammation, consequences are widespread and linked to diverse manifestations that include osteoporosis,² lymphoma,^{3–4} pneumonia⁵ and increased mortality.⁶ Symptoms vary, with some patients having diarrhoea and malabsorption (often termed ‘classical CD’), others suffering from constipation, fatigue and depression (non-classical CD) and some are asymptomatic (subclinical CD).⁷ The global prevalence of CD is about 1%–2%,^{8,9} but seems to be increasing.^{10,11}

Lifelong adherence to a gluten-free diet (GFD) is the only available treatment for CD.¹ For several reasons, patients find the GFD to be exceedingly burdensome,¹² that is, it is socially restrictive¹³ and more expensive than ordinary food.^{14–16} Patients differ in their ability to adapt psychologically to CD. Some people have little difficulty in adopting the GFD, whereas for others, living with CD is a

Significance of this study

What is already known about this subject?

- A gluten-free diet is the only treatment option of coeliac disease, but recently an increasing number of trials have begun to explore alternative treatment strategies.
- A large number of trials of non-dietary treatments for coeliac disease are ongoing or under way.
- There is no consensus on outcome measures in coeliac disease trials.

What are the new findings?

- After an extensive literature review, 17 researchers and 2 patient representatives from 10 countries reviewed the use and suitability of histology, serology, clinical outcome assessment (including patient-reported outcomes), quality of life and immunological tools that comprised gluten immunogenic peptides for trials in coeliac disease.
- In this paper, we make expert-based recommendations for use of these outcomes in coeliac disease therapeutic trials.

How might it impact on clinical practice in the foreseeable future?

- Following the outlined recommendations of this paper, will increase transparency and comparability of coeliac disease therapeutic trials with benefit to patients, healthcare and the pharmaceutical industry.

daily struggle.¹³ In addition to the burden of treatment, patients with CD frequently have ongoing symptoms and mucosal healing is slow and often incomplete. For these reasons, there is a need for alternative treatments of CD, as suggested by the intensive research efforts undertaken in different laboratories.¹⁷ Potential targets for treatment include glutenases, modified or pretreated gluten, gluten sequestrants, neutralising antibodies, inhibitors of intestinal permeability, lymphocyte blockers, including anti-interleukin-15, tissue transglutaminase (TG2) inhibitors, immune tolerance induction, exposure to hookworms and



To cite: Ludvigsson JF, Ciacci C, Green PHR, *et al.* *Gut* 2018;**67**:1410–1424.

DQ2-blocking peptide analogues.^{17 18} Several of these drugs are now being tested in phase I or phase II trials, and a recent study found that novel therapies attract the interest of patients with CD more than any other disease-related topic.¹⁹

Of importance is that treatment effects are measured against robust standards. A recent systematic review²⁰ identified six histological CD activity indices,^{21–26} five patient-reported outcomes (PROs)^{27–31} and four indices for endoscopic CD activity^{32–35} that have been used for coeliac trials.

In the present paper, we have explored clinical, serological, histological and immunological outcome measures for performing clinical trials in CD, and importantly, have included the patient perspectives concerning recommendations for their use.

METHODS

Task force

Coauthors were invited by JFL and DAL with the aim to obtain a group with knowledge, experiences and interests that reflect the heterogeneity of outcome measures used in trials of CD. Most of the participating researchers were adult gastroenterologists (PHRG, CC, DSS, KKa, DS, JAM, JT, KEAL, DAL) but our group also included six paediatricians (JFL, NRR, Kku, MJM, AP, IRK-S), one pathologist (MMW), one basic scientist (AR-H) and two representatives of patient organisations (SS and KV). Members of this diverse collaboration originated from 10 countries. Most of the coauthors participated in the CD meeting organised by MJM in Tampere, Finland on 24–25 November 2016 (which provides the motivation for the subtitle of this paper).

Literature review

Coauthors were divided into seven teams of three to four individuals who jointly reviewed five domains: serology, histology, immunology, PROs and other outcome measures. The Karolinska Institutet library carried out literature searches for relevant papers up until 1 December 2016 (see online supplementary appendix). This search yielded 10 062 references. After reviewing titles and abstracts for these 10 062 references (online supplementary appendix), there remained 941 publications that were considered potentially relevant for this review and subsequently read in detail.

In this paper, we issued a number of recommendations. Where appropriate these were graded according to the method suggested by the Oxford Centre for Evidence-based Medicine,³⁶ where grade A evidence represents the highest level of evidence and grade D the lower available evidence (generally based on the opinion of experts but with no preceding randomised trials, cohort or case-control studies). The appendix contains a detailed description of grade A–D. All recommendations were subject to a post hoc voting on a five-level scale (strongly disagree, disagree, not agree or disagree, agree and strongly agree).

Manuscript draft

JFL wrote the first draft of the paper. The text was then extensively revised by the coauthors. JFL and DAL supervised these revisions, but all authors contributed and agreed on the conclusions and the final wording of the paper.

This is a series of expert-based recommendations and did not meet the requirements of a formal systematic review (eg, the Preferred Reporting Items for Systematic Reviews and Meta-Analyses statement³⁷). We aimed to highlight the state-of-the-art of designing intervention trials in CD.

RESULTS

Histology

Clinicopathological correlation is key to the diagnosis of CD in adults and children. In adults, confirmation of the diagnosis by duodenal biopsy is the gold standard.^{7 38 39} In Europe, a ‘biopsy-sparing’ protocol (with defined limitations for use) has been adopted for symptomatic children defined by an anti-TG2 titre ≥ 10 times the upper limit of normal, positive endomysial antibody (EMA) on a second blood draw and positivity for human leucocyte antigen haplotypes HLA-DQ2 and/or HLA-DQ8.⁴⁰ To evaluate effective treatment for CD, quantitative histological assessment (morphometry) outperforms qualitative histology (eg, the Marsh score) in a trial setting.²⁰ In clinical trials, optimised biopsy protocols should be followed for assessment of mucosal damage or healing.

Well-known classifications in histological assessment are described by Marsh and modified by Oberhuber^{7 38} and Corazza and Villanacci.⁴¹ Although grouped classifications are practical in clinical work, recent studies have shown imperfect reproducibility and reliability.^{26 42}

Recently published recommendations for biopsy diagnosis of CD in adult patients for the number of biopsies and sites are available,^{38 39} with optimal laboratory processing alongside structured reports to include validated morphometric analysis.^{26 38}

It is recommended to take at least five duodenal biopsies, one or two from the duodenal bulb (D1) and four from the second part of the duodenum (D2).^{38 39} These biopsies should be taken across circular folds to avoid a crushing artefact.⁴³ Endoscopists should take one biopsy specimen per pass of the forceps in that a single-biopsy technique improves the yield of well-oriented duodenal biopsy specimens.⁴⁴ Biopsies from D1 and D2 should be reviewed separately by a pathologist.⁴⁵

When processed in the laboratory, biopsies should be oriented correctly and sectioned at three levels. In trial settings it is appropriate to always count intraepithelial lymphocytes (IELs) and state the number present/100 enterocytes (normal counts are $\leq 25/100$ enterocytes).^{7 26 39 45 46} IELs can be counted in H&E stained sections; however, immunohistochemistry with CD3 is preferred by some pathologists.^{26 47} Frozen tissue specimens have been used to evaluate T-cell receptor gamma delta positive ($\gamma\delta+$) T cells,⁴⁸ but new antibodies for use on paraffin-embedded specimens are now available.^{49 50} Identification of a high density of $\gamma\delta+$ T cells is relatively specific for CD and can be useful when histological diagnosis remains equivocal.^{48 49}

Immunohistochemistry to show deposited immunoglobulin A targeting TG2 in the small bowel mucosa is accurate in the detection of CD if patients are on a gluten-containing diet.⁵¹ These deposits have had 100% sensitivity⁵² and a mean specificity of 94%⁵³ for CD. The deposits have been used in several gluten challenge studies to measure gluten reactivity.⁵² This technique requires the use of frozen tissue. The use of frozen tissue in clinical trials has shown variable results, which mirrored serum TG2 antibodies with increased deposits.⁵⁴

Morphometry, in which continuous variables such as the villous height-to-crypt depth ratio and IEL density are measured separately, overcomes certain problems encountered when using grouped classifications.^{26 55} Of note, a threshold change of 0.4 represents a measurable and likely clinically relevant difference between villous height: crypt depth ratio measurements. A villous height: crypt depth ratio of < 2 is indicative of atrophy, active disease. Patients with treated CD have values above 3. Similarly, $\geq 30\%$ change in T-cell IEL densities is considered clinically significant.²⁶

Importantly, morphometry, which has produced excellent reproducibility and reliability,²⁶ has a significant role in clinical trials in which reliable and accurate measurements are a requirement.⁵⁶ Whichever classification is used, two blinded observers should read the histology to ensure reliability in clinical trials.²⁶ It should be noted that CD can be patchy and there is some intra-subject and even intrabiospy variability in villous architecture and lymphocyte numbers contributing to sampling error and difficulties in interpretation. Given this, and that only a *small* proportion of the proximal small intestinal mucosa is evaluated by conventional biopsy review, new tools are needed for assessment of mucosal health. The optimal timing of biopsy to evaluate healing should be a clinicopathological decision dependent on treatment offered and taking into account possible sampling errors by following protocols for biopsy sites.

In CD, there may be concurrent upper GI pathologies (eg, *Helicobacter pylori* infection,^{57 58} lymphocytic gastritis⁵⁹ and eosinophilic oesophagitis/oesophageal eosinophilia⁶⁰) that should be assessed at initial endoscopy if clinical history is suggestive, and if present, reassessed post-treatment because these may contribute to ongoing symptoms not related to small intestinal damage.

Patients included in studies for CD therapy must have had an initial robust diagnosis. Occasionally, patients will have been diagnosed without histological confirmation.⁶¹ These patients should not be included in gluten challenge studies but instead included in trials of active CD treatment. Patients with a study entry biopsy confirming villous atrophy (VA) and a record of positive serology and permissive HLA status should be eligible for treatment studies in CD.

As per the current European Society for Paediatric Gastroenterology Hepatology and Nutrition guidelines,⁴⁰ we are reluctant to suggest timelines for control biopsies for children, although a recent paper found that mucosal healing may not be as complete as previously assumed.⁶² For now, follow-up biopsy in children should be dictated by clinical needs.

Table 1 summarises changes in mucosal histopathology, serology and symptoms in clinical trials with gluten challenge.

Recommendations: Histology is an essential outcome measure in any trials of CD treatment (grade B). Histology should be performed both before and at the end of the trial when healing or histological relapse is the primary outcome measure (grade B). In a gluten challenge study, successful treatment may be characterised as no change, or, in a treatment study, as histological improvement by a significant increase in the villous height:crypt ratio (>0.4 being considered relevant) and/or a $\geq 30\%$ change in IEL densities (grade D). Additionally, histology may be useful as a criterion for study inclusion in which participants in gluten challenge studies should have a high villous height:crypt depth of >2–3 and participants in treatment trials should have a decreased villous height:crypt depth ratio of <2–3 (grade D). Histological evaluation should follow a priori histology protocols using quantitative measures.

Vote: agree: 7; strongly agree: 12.

Serology

Serology is a cornerstone in the diagnostic workup for CD.^{38 39 63 64} IgA auto-antibodies to TG2 and IgG antibodies to deamidated gliadin peptides (DGPs) are central diagnostic tests for active CD. In IgA-sufficient patients, IgA-anti-TG2 is the most predictive and reproducible single test, although IgA EMA performs similarly well in some expert laboratories and is often used as a confirmatory test. IgG anti-DGP displays similar

sensitivity as IgA anti-TG2 but has lower specificity. Selection of optimal serological tests is mandatory because not all commercial assays perform equally well.^{65 66} Importantly, calculation of results and thus numerical values for the same samples⁵⁹ also may differ and only tests with a multipoint calibration curve give values proportional to serum antibody concentration.⁴⁰ Differences in assays can make interpretation in and comparisons between clinical trials difficult. This difficulty is in part due to different epitopes that are detected by different tests, different calibration or to antibodies with lower avidity and specificity.^{67–69} Thus, an important shortcoming when using serology to evaluate the outcome of gluten challenge is the wide range of response. When the UK National External Quality Assessment consortium tested the same positive samples in 14 commercial anti-TG2 assays, large differences in antibody levels were found, consistent with substantial variability for antibodies used in the diagnosis of CD.⁶⁵

Overall, a correlation exists between IgA-anti-TG2 antibody titres and the severity of mucosal damage by histology, as well as the histological outcome on a GFD.^{70–73} Yet, a recent meta-analysis found that serum TG2 and EMA often underestimate the degree of VA.⁷⁴ Antibody titres below diagnostic cut-offs, thus, do not predict a normal or near-normal (Marsh I) histology. The one caveat is that biopsies usually sample a short segment of the (descending) duodenum, whereas active CD can affect large portions of the small intestine that occasionally extend down to the ileum.⁷⁵ Therefore, a patient may be in clinical and serological remission with residual inflammation in the proximal duodenum but which is quantitatively much less extensive than before.⁷⁴ Thus, in one of the largest studies to date,⁷⁶ IgA anti-TG2 failed to detect 44% of persistent VA (Marsh III) in patients with CD on a GFD for >1 year.

Normal serology is generally required for entry into a gluten challenge study to ensure that participants do not have severe VA prior to gluten challenge. Conversely, serology above diagnostic cut-offs has not been used as an inclusion criterion for treatment studies in non-responsive CD because many people with a normal serology will still have VA and ongoing symptoms.⁷⁴ Participants with elevated serologic tests may respond better to some therapies. While this remains to be confirmed, stratification by CD antibody levels at study entry should be considered.

In clinical trials, serology may be used in assessing change during gluten challenge or to monitor longer treatment studies. The antibody response to gluten challenge depends on four factors: duration of the previous GFD, daily amount of ingested gluten,⁵⁴ duration of gluten intake and individual factors.

Patients with CD may tolerate different levels of gluten exposure. When low (1–3 g/day) or moderate (3–5 g/day) amounts of gluten were administered to 25 Finnish patients with CD in remission for 12 weeks, only 67% of the patients with CD showed signs of mucosal inflammation and 43% developed positive IgA-anti-TG2 antibodies.⁵⁴ However, in a US study of 20 adult patients with CD in remission challenged with 3 or 7.5 g gluten/day for 2 weeks, Marsh III histology developed in 68% of the patients with CD, whereas anti-TG2 and anti-DGP antibodies increased in only 25% for anti-TG2 and 30% for anti-DGP. Remarkably, positivity for both antibodies increased to 55% and 45% 2 weeks after the end of gluten challenge.⁷⁷ A recent study from Norway showed even lower proportion of patients responding serologically after 2 weeks.⁷⁸ Therefore, the histology at week 2 and serology at week 4 combined showed a gluten response in nearly 90%, with no difference between both doses.⁷⁷ Notably, some rare patients who had been on a GFD for years may develop a tolerance to gluten ingestion that may last

Table 1 Coeliac disease: results of clinical trials with gluten challenge, changes in mucosal histopathology, serology and symptoms

	Patient, n	Amount of gluten	Challenge duration	Mean reduction in VhCRD*	Mean increase in CD3+IELs	Marsh-Oberhuber classification	Serology	Symptoms	Other
Lähdeaho <i>et al</i> 2014 ⁸⁹	16	1.5 g	6 wk	0.6†	NA	NA	NA	NA	–
	21	2.0 g	6 wk	0.8†	Statistically significant increase†	NA	1/21 seroconverted to positive EMA	Trend towards more symptomatic in GSRS, CDQ and SF-36 questionnaires	$\alpha\beta$ and $\gamma\delta$ T cells per mm of epithelium elevated. No change in TG2 IgA deposits.
	13	3.0 g	6 wk	1.1†	NA	NA	NA	NA	–
	13	6.0 g	6 wk	1.7‡	NA	NA	NA	NA	–
Leffler <i>et al</i> 2013 ⁷⁸	10+10§	3 and 7.5g§	Day 3	0.52¶	3.1 per 100 ECs¶	NA	Mean increase in IgA-TG2 titre was 1.03¶	Significant change in both CSI† and GSRSt	No significant change in lactulose to mannitol ratio.
			Day 14	1.09‡	19.22 per 100 ECs¶	10/20 showed histological response	Mean increase in IgA-TG2 titre was 2.86†	Significant change in both CSI† and GSRSt	
Tack <i>et al</i> 013 ¹⁴⁴	7	7 g	2 wk	NA	NA	3/7 deteriorated¶	1/7 seroconverted to positive IgA-TG2	No significant change in CDQ	5/5 had increased deposits of TG2 IgA at end.
Brottweit <i>et al</i> 2013 ⁸²	11	Four slices of bread	Day 4	NA	Data not presented¶	6/11 deteriorated	NA	4/11 had increased abdominal symptoms	IFN- γ , IL-8 and TNF- α genes increased.†
Lähdeaho <i>et al</i> 2011 ⁵⁴	10	3–5 g	12 wk	1.3†	39 per mm of epithelium‡	NA	5/10 had increase in IgA TG2 titre†	8 developed mild to moderate symptoms*	5/10 had increased deposits of TG2 IgA.
	11	1–2 g	12 wk	1.0†	20¶	NA	4/11 had increase in IgA TG2 titre†	7 developed mild to moderate symptoms*	4/11 had increased deposits of TG2 IgA.
Catassi <i>et al</i> 2007 ¹⁴⁵	13	10 mg	3 months	–1%¶	0%	None	No change	NA	–
	13	50 mg		–20%††	+12%¶	2/13 deteriorated	IgA-TG2¶ IgG-AGA decreased†		
Holm <i>et al</i> 2006 ¹⁴⁶	10	14 g (range 7–19 g)	3–24 months	Statistically significant increase†	Statistically significant increase†	NA	EMA and IgA-TG2 positive in 5/10 at 1 month	4/10 developed symptoms at histological relapse	–
Catassi <i>et al</i> 1993 ¹⁴⁷	10	100 mg	4 wk	0.2†	8 per 100 ECs†	NA	4/10 seroconverted to positive in IgA AGA	None	–
	10	500 mg		0.5†	15 per 100 ECs†		6/8 seroconverted to positive IgA AGA	3/10 showed clinical symptoms	

*Over 0.4 is considered a statistically significant change.²⁶†Significant change compared with baseline ($P \leq 0.05$).‡Significant change compared with baseline ($P < 0.001$).

§The study consisted of a low and high gluten group with no significant differences between the groups in histology or serology. In the study, the data were presented after combining the two groups.

¶Non-significant change ($P > 0.05$).

**Data not presented.

††Significant change compared with placebo ($P < 0.05$).

CDQ, Coeliac Disease Questionnaire; CSI, Coeliac Symptom Index; EC, epithelial cells; EMA, endomysial antibody; GSRS, Gastrointestinal Symptom Rating Scale; IFN, interferon; IL, interleukin; NA, not available; SF-36, 36-Item Short-Form; TG2, tissue transglutaminase; TNF, tumour necrosis factor; VgCRD, villous height crypt depth ratio; wk, week.

for several years.⁷⁹ Overall, high serologic titres, or significant increases in serologic titres, are predictive of VA, but substantial mucosal changes may occur without a significant change in serology.

It is important to note that the serological tests were developed for the diagnosis of CD. Currently, the Food and Drug Administration (FDA) have only cleared use of coeliac serologies as an aid for diagnosis of patients with suspected CD.⁸⁰ This restriction limits how serologies can be used in regulatory trials, although they are routinely used for monitoring in clinical practice. To date, no manufacturer has submitted a claim for use of serological tests for disease monitoring and the FDA is only able 'to review submitted claims'.⁸⁰ Nonetheless, well validated IgA anti-TG2 and IgG anti-DGP tests will be important assets to clinical studies by helping to monitor CD activity during short and especially longer duration gluten challenge trials. Their further validation in ongoing trials may lead to FDA approval as secondary or combined (with histology or symptoms) primary end points in phase II and III clinical studies (table 1).

Recommendations: Although serology is not approved for use as a primary pivotal clinical trial outcome by the FDA, IgA TG2 and IgG DGP should, at a minimum, be measured at study entry and at completion in trials of CD (grade B). For entry into a gluten challenge study, participants should have near-normal titres, whereas for treatment studies, titres may be either elevated or normal, with stratification by serologic titre at study entry possibly as an a priori analysis (grade B). Choice of assay should be made with care and attention be paid to dynamic range, especially around or below the cut-off for normal ranges. Preferably, an assay with a calibration curve should be applied. Although cut-off ranges for diagnosis may not be optimal for monitoring response or predicting VA, any significant increase during a trial suggests increasing CD activity and may be used as a key outcome in some studies.

Vote: agree: 2; strongly agree: 17.

Immunological tools to measure treatment outcomes

Known innate, and particularly adaptive, immune mechanisms in CD, are prime candidates to measure treatment outcomes with and without gluten challenge. These may be non-invasive blood tests, duodenal biopsies with histological assessment by immunocytochemistry, gene expression signatures or in vitro culture.

With in vivo gluten challenge, there is rapid immune activation in the duodenum.^{81–83} One study found that interferon (IFN)- γ was increased both at baseline and with gluten challenge and for this reason does not appear to be a useful measure of disease activity.⁸² A whole-blood IFN- γ release assay is a much more promising measure for identifying immune responsiveness to gluten.^{84 85}

Treated patients with CD and healthy individuals show highly variable differences in serum cytokines and chemokine levels.⁸⁶ Gluten challenge, however, induces a wave of cytokine release.⁷⁸

In the lamina propria of the duodenal mucosa gluten peptides are taken up by dendritic (antigen-presenting) cells with surface HLA-DQ2 or DQ8 MHC molecules to stimulate gluten peptide reactive CD4+ T cells.^{87–90} On day 6 of postgluten challenge an increase of active disease gluten responsive T cells was seen in peripheral blood.^{91 92} These T cells can be demonstrated in Enzyme-Linked ImmunoSpot (ELISPOT) assays of cytokines when rechallenged with gluten ex vivo. Overall, 80%–90% of treated patients with CD in remission will show a positive response on challenge. T cells can also be demonstrated by binding to HLA-DQ-gliadin peptide tetramers, a

construct consisting of multimers of HLA-DQ2 or HLA-DQ8 molecules bound to a gluten peptide and a reporter molecule giving a signal in flow cytometry.^{93 94} Here, also 80%–90% of challenged patients with CD in remission will have a positive test.⁹⁵ However, both the ELISPOT and the HLA-DQ2-gluten tetramer tests are confounded by large interindividual differences and small number of studied patients. Recently, the HLA-DQ2-gluten tetramer technology has demonstrated disease-specific T cells in the peripheral blood even without a gluten challenge.⁹⁶ This, together with the demonstration of restricted T-cell receptors,⁹⁷ may lead to new outcome measurements.

Serum IgA anti-TG2 antibody levels as biomarkers of disease activity seem to be a useful tool (see the 'Serology' section). Peripheral blood B cells may also prove to be a potential source of future biomarkers.⁹⁸ The local mucosal production of antibodies targeting extracellular TG2 in vivo has shown potential in diagnosis,⁵¹ but was not informative beyond standard histology in a clinical drug trial with gluten challenge.⁹⁹

Recommendations: Although several immunological markers are under development as potential outcome measures, they have not been validated for therapeutic trials. At this point, they should only be used as exploratory outcomes in phase II and III clinical trials (grade D).

Vote: agree: 2; strongly agree: 17.

Gluten immunogenic peptides as a compliance measure

Symptom monitoring, serology and histology are at best indirect measures of GFD adherence with imperfect overall accuracy.¹⁰⁰ Similarly, diet questionnaires are poor predictors of gluten exposure.¹⁰⁰

Gluten immunogenic peptides (GIPs), including the 33-mer peptide from α 2-gliadin, are resistant to GI digestion.^{101 102} Because of this resistance, GIP can be detected in faeces or urine and thus provides direct evidence and likely quantitation of intake.¹⁰³ A clinical trial examined correlations between faecal GIPs and traditional methods to monitoring the GFD.¹⁰⁴ The majority (85.7%) of children with CD under 3 years of age had faeces negative for GIPs. Among those aged ≥ 13 years, faecal positivity for GIPs rose to 39.2%. More males were positive for GIPs in faeces compared with females (60% vs 31.5%, $P=0.034$). Serum IgA anti-TG2 was found negative in 40 of the 56 patients with GIP-positive stools. Today's data suggest that GIP testing may be superior to questionnaires or anti-TG2 antibodies.¹⁰⁴ Furthermore, a strong correlation has been demonstrated between the absence of GIPs in urine and healing of the intestinal epithelium.¹⁰⁵ The first therapeutic clinical trials using the technology are ongoing (NCT02637141, NCT02633020). Whereas coeliac trial investigators and sponsors had to previously guess whether patients were consuming gluten or adhering to the GFD, this technology decreases the guesswork with data to accurately interpret results and outliers. The assay could possibly be developed for quantitative detection of the rate of glutamine residue deamidation in trials aiming at interfering with transglutaminase activity.¹⁰⁶

Recommendations: GIP testing is a promising tool for evaluating and selecting patients for clinical trials in CD aimed at reducing toxicity related to gluten exposure (grade D). Hence, it should be considered in future trials, especially for trials in non-responsive CD for therapies that are designed to prevent symptoms because of inadvertent gluten exposure (grade D).

Vote: agree: 4; strongly agree: 15.

Clinical trial end points

The ideal clinical trial end point should be clearly linked to an outcome important to patients, reliable, responsive to treatment, clinically or physiologically proximal to the outcome of interest and efficient and scalable for use in diverse clinical trial settings. Although in some areas biomarkers may be used as primary outcomes in clinical trials, these rarely have sufficient data for regulatory acceptance.¹⁰⁷ In contrast, clinical outcome assessments can be more easily linked to patient well-being: clinical outcome assessments are grouped into PROs, clinician-reported outcomes, observer-related outcomes and performance outcomes. Clinician-reported outcomes, such as physician global assessment, are of limited value as they do not directly assess patient status and generally do not correlate well with PROs. Observer-related outcomes can be vital in specific populations, such as young children, where direct response is not possible. Symptom-focused PROs are the main clinical outcome assessment in use in CD and in gastroenterology overall and are the focus of this section, although other clinical outcomes will also be discussed.

From a clinical/practical perspective, PROs can be helpful to monitor patient status and target quality improvement initiatives.¹⁰⁸ A growing number of digital devices allow patients to track and transmit symptom data to their physician; however, for these to be useful for practising clinicians, PROs must be easy to administer and interpret, as well as to allow feasible clinical interventions.¹⁰⁸ In research and clinical trials, the key features of PROs that should be considered are high responsiveness to change and low participant burden.

For patients, the ideal PRO must focus on the symptoms or disease attributes most meaningful to them while again minimising time and complexity of use. In CD, this is particularly important given the significant impact on emotional, mental and social well-being due to the constant vigilance required by the GFD. Finally, from a regulatory standpoint, the ability to use a PRO in a pivotal clinical trial to support a labelling claim depends on whether its characteristics (eg, concept being measured, content validity, conceptual framework, intended population, format, scoring) are satisfactory and clearly

documented in a regulatory dossier, which is now available for only a few PROs.¹⁰⁹

Presently available PROs frequently used in CD or developed for CD include the Gastrointestinal Symptom Rating Scale (GSRS),^{110 111} the Celiac Symptom Index (CSI),³¹ the Celiac Disease PRO (CeD-PRO),²⁷ the Celiac Disease Symptom Diary (CDSYD)¹¹² and the Celiac Disease Assessment Questionnaire (CDAQ).¹¹³ Of these, the GSRS has been used most frequently, ranging from natural history to the effects of the GFD to clinical trials of novel therapeutics.^{110 114–116} The GSRS was developed originally for peptic disease and irritable bowel syndrome,¹¹⁷ but because the symptoms of many GI disorders overlap, it has proven useful for a variety of GI disorders, including CD.¹¹⁸ However, it is not optimised for CD and would not be appropriate for use in pivotal trials. Conversely, the CSI was developed specifically for CD and has been used in many cross-sectional and interventional studies.^{119 120} However, its development was completed before the 2009 FDA guidance¹⁰⁹ and thus the CSI lacks much of the documentation necessary for regulatory clearance. Conversely, both the CDSYD and the CeD-PRO were developed specifically for regulatory approval of CD therapeutics and are preferred instruments for this purpose.

The CDAQ was recently developed and assesses a variety of domains: symptoms, dietary burden, worry, stigma, and social isolation.¹²¹ As such, it is a hybrid of the health-related quality of life (HRQoL) tools discussed below and a symptom PRO. It is unclear if this instrument were developed and documented in line with regulatory guidance: in the CDAQ both constipation and diarrhoea are evaluated in a single combined question that may make changes in these important symptoms difficult to assess. However, the incorporation of these or similar HRQoL domains related to disease burden is critical in ensuring that outcomes are relevant and meaningful to patients.

Across these instruments, there is significant overlap, which is expected given the limited number of GI symptoms in general (table 2): common to all the PROs are diarrhoea, abdominal pain, bloating, and nausea. It should be acknowledged that while PROs in general may be developed and tested in specific diseases, they will not discriminate between diseases and therefore scores

Table 2 Symptoms assessed across PROs in CD

Symptom	CDSYD ¹¹²	CD-PRO ²⁷	CSI ³¹	CDAQ ¹²¹	GSRS ¹¹⁷	CD-GSRS ¹⁴⁸
Abdominal pain	X	X	X	X	X	X
Diarrhoea	X	X	X	X†	X	X
Bloating	X	X	X	X	X	X
Nausea	X	X	X	X	X	X
Constipation	X			X†	X	
Fatigue*	X	X	X	X		
Headache*	X	X	X			
Skin rash	X					
Flatulence	X	X			X	X
Burping					X	X
Problems thinking*	X			X		
Incomplete emptying			X		X	
Loss of appetite			X			
Hunger pains			X		X	X
Stomach rumbling		X	X		X	X
Reflux					X	

*Removed from final PRO as per the Food and Drug Administration recommendations.

†Assessed in a single domain.

CD, coeliac disease; CDAQ, Celiac Disease Assessment Questionnaire; CDSYD, Celiac Disease Symptom Diary; CSI, Celiac Symptom Index; GSRS, Gastrointestinal Symptom Rating Scale; PRO, patient-reported outcome.

for patients with different GI disorders will overlap.¹²² There is also a poor correlation between symptoms, histology, and serology^{80 123} that is due in part to a different time to response after gluten exposure or coexisting symptoms due to irritable bowel syndrome or undetected food allergy.¹²⁴ Thus, it is also clear from recent clinical trials that many symptomatic patients have no histological or serologic evidence of active CD and many patients with significant enteropathy have few or no symptoms.^{78 80 112 116 125} Meanwhile complete recovery of small intestinal lesions is very rare in adult CD patients despite symptom disappearance.¹²⁶

In addition, PRO use in CD can be challenging because of symptom heterogeneity (eg, asymptomatic or paucisymptomatic patients) and variable extraintestinal manifestations, for which no PROs are available. In contrast to disorders such as chronic constipation or headache in which one symptom defines the condition, symptoms in CD can vary substantially between individuals and hence careful attention to PRO use is mandatory. Moreover, responsiveness to change of non-symptom-based dimensions in PROs can vary significantly and must be carefully assessed in relation to the intervention under investigation. For example, measures of quality of life (QoL) may be less amenable to change if the intervention does not reduce dietary burden, social isolation and stigma, which are closely linked to managing the GFD. Comparing overall PRO mean scores at baseline and postintervention may dilute the treatment effect if not all domains change or if changes cancel out each other (eg, diarrhoea improves but results in constipation by disrupting bowel movements). Another option is to compare the means only in prespecified domains (eg, include diarrhoea but not constipation). However, this approach may result in a highly selected population that is not representative. A more sophisticated approach is to limit primary assessment of the effect of intervention to the symptoms most bothersome to a specific patient and then to include all individuals with bothersome symptoms in the final PRO assessment. Even with this approach, for treatment trials, only patients with symptoms measured by the primary PRO outcome can be enrolled. Given the heterogeneity of CD, it is likely that sequential trials will need to be performed with different outcomes in order to understand the utility of a particular therapy. For example, a therapy found to be useful in improving GI symptoms in CD could be assessed in a later trial assessing itch in patients with dermatitis herpetiformis.

Particularly noteworthy is that while great progress has been made in PROs in CD, there is limited experience using the more recent PROs in languages other than English and none has been validated for use in paediatric populations. Although it is expected that the PROs developed for CD in adults will be applicable for paediatric populations, this still requires validation. For example, young children and teens may define improved QoL differently because of unique challenges in school, social settings and peer relationships. Furthermore, improvement in extraintestinal symptoms, including behavioural changes, may be more relevant among paediatric patients. Development of formats suitable for caregivers of children unable to independently complete the questionnaires must also be advanced. Development of responder definitions and minimal clinically important change are additional areas requiring development to realise the potential of CD PROs in clinical research.

Given that the relative SD of histology is substantially smaller than that of symptoms, for a study where several hundred patients are needed to adequately detect differences in symptoms, fewer patients may be needed for the histology end point. Under such circumstances, researchers may choose to perform

histology on a random sample of study participants provided that they have sufficient study power for the histology end point.

Recommendations: Clinical end points must be included in trials of CD and generally PROs should be a primary outcome in studies of treatment of active CD, generally late phase II and III (grade D). Thus far, there is insufficient evidence to recommend one specific scale, although the CSDS and the CeD-PRO appear most likely to meet regulatory requirements (grade D). Given the heterogeneity of symptoms in patients with CD, it is adequate to limit analyses to certain domains, either in the study overall or to allow for participant-specific symptom assessments. However, such decisions should be made at study entry and rigorously documented.

Vote: agree: 3; strongly agree: 16.

Health-related quality of life assessment

Clinical trials must also consider the ongoing psychological burden of CD to better understand the outcomes. Many studies in different settings suggest that CD has a considerable impact on HRQoL.^{12 127} In untreated CD, GI symptoms and extraintestinal issues, such as isolated anaemia, fatigue and malaise, may be responsible for reduced HRQoL.¹²⁸ In general, the treatment of CD results in significant improvement in the HRQoL of symptomatic patients.¹²⁹ Even in patients with silent or asymptomatic screening-detected CD, improvement in both symptoms and HRQoL has been shown in numerous studies.^{116 130-132}

Impairment in HRQoL may also contribute to and be impacted by psychological disorders (eg, anxiety and depression) seen in patients with CD.^{12 133} Whereas anxiety generally improves after diagnosis and treatment CD, depression may exist before and persist after diagnosis.¹³⁴ Moreover, anxiety and depression alone or through their impact on HRQoL can impair dietary compliance.^{12 135} The interaction between mood disorders, GFD adherence and HRQoL is incompletely understood and should be addressed in future trials. Additionally, the burden of a GFD may lead to ongoing HRQoL impairment despite symptom relief and improved physical well-being after dietary intervention.^{13 19}

Because of small numbers of studies and variations in study designs, populations and HRQoL measures, we lack a complete understanding of degrees and drivers of HRQoL in individuals with untreated and treated CD. There are also few studies about the effect of CD treatment in the outcome of depression and anxiety. Therefore, further studies are required if we wish to know more about this specific aspect of CD.

Although HRQoL is generally not accepted by regulatory agencies as a primary outcome in pivotal trials, this is a key outcome for both patients and clinicians and may determine whether a therapy succeeds or fails. Arguably, a main goal of therapeutics in CD, in addition to histological and symptom improvement, is to improve HRQoL. Indeed, it is probable that some patients in histological and symptomatic remission could have a significant HRQoL benefit from pharmacological therapy owing to a reduction in burden of the GFD and anxiety regarding potential exposures. There are several HRQoL scales that have been specifically developed for or used in CD (table 3).

Recommendations: HRQoL in CD is complex and multidimensional and may be a more relevant concern to patients than specific symptoms. In any trial aiming to improve CD control (as opposed to gluten challenge studies that aim to prevent worsening), measurement of HRQoL should be considered a critical end point that may help to determine the overall value of a therapy or intervention to both patients and payors (grade D).

Vote: agree: 4; strongly agree: 15.

Table 3 Quality of life instruments relevant to coeliac disease

Adult instruments			
Instrument	Author	Participants	Instrument description
Generic			
Zung Self Rating Depression Scale ¹⁴⁹	Ciacchi 2003 ¹⁵⁰	581 CD	20-Item self-report, 4-point Likert scale. Assesses four characteristics of depression.
Gastrointestinal Quality of Life Index ¹⁵¹	Casellas 2008 ¹⁵²	340 CD- 177 untreated vs 163 on GFD	36-Item self-report, 4-point Likert scale. Assesses five domains: symptoms, physical function, emotional function, social function and medical function.
COPE (brief) ¹⁵³	Smith 2011 ¹⁵⁴	156 CD	28-Item self-report, 4-point Likert scale. Measures 14 coping responses.
	Sainsbury 2013 ¹⁵⁵	189 CD	
EuroQol-5D ¹⁵⁶	Casellas 2008 ¹⁵²	340 CD	5-Item self-report, 5-point Likert scale plus a general health rating (scored on a 20 cm visual analogue scale). Measures five dimensions: mobility, self-care, daily activities, pain/discomfort and anxiety/depression.
	Grey 2010 ¹⁵⁷	783 CD	
	Norström 2011 ¹⁵⁸	1031 CD	
	Casellas 2015 ¹⁵⁹	366 CD on GFD>1 year	
Beck Depression Inventory ¹⁶¹	Ramírez-Cervantes 2015 ¹⁶⁰	80 CD on GFD≥6 months	21-Item self-report, 4-point Likert scale. Evaluates 21 symptoms of depression, including emotions, behavioural changes and somatic symptoms.
	Borghini 2016 ¹⁶²	210 CD (70 untreated vs 70 on GFD 6–12 months vs 70 on GFD>12 months)	
	Nachman 2009 ¹²⁸	132 Newly diagnosed CD	
Psychological General Well-Being Index ¹⁶⁴	Nachman 2010 ¹⁶³	53 Newly diagnosed CD	22-Item self-report, 6-point Likert scale. Includes six dimensions: anxiety, depressed mood, positive well-being, self-control, general health and vitality.
	Ford 2012 ¹⁶⁵	288 CD	
	Smith 2011 ¹⁵⁴	156 CD	
	Mustalahti 2002 ¹³¹	19 screen-detected CD vs 21 symptom-detected CD	
	Mahadev 2016 ¹²⁷	211 (71 screen-detected CD vs 140 symptom-detected CD)	
	Borghini 2016 ¹⁶²	210 CD (70 untreated vs 70 on GFD 6–12 months vs 70 on GFD>12 months)	
	Paavola 2011 ¹³⁰	466 CD (96 screen detected CD vs 370 symptom-detected CD vs 110 non-CD controls)	
	Peräaho 2004 ¹⁶⁶	39 CD	
	Vilijamaa 2005 ¹⁶⁷	98 CD (54 screen-detected CD vs 44 symptom-detected CD)	
	Ukkola 2011 ¹³²	698 CD	
Hospital Anxiety and Depression Scale ¹⁶⁹	Smith 2011 ¹⁵⁴	156	14-Item self-report, 4-point Likert scale. Seven items pertain to anxiety and seven to depression.
	Paarlahti 2013 ¹⁶⁸	596 CD	
	Häuser 2007 ¹⁷⁰	516 CD	
	Häuser 2006 ¹⁷¹	446 CD	
	Barratt 2011 ¹⁷²	225 CD	
The Short-Form 36-Item QoL measure ¹⁷⁴	Barratt 2011 ¹⁷³	225 CD	36- Item self-report, 3-point, 5-point and 6-point Likert scales as well as binary (yes/no) response items. Measures eight domains: physical functioning, role limitations due to physical health, role limitations due to emotional problems, energy/fatigue, emotional well-being, social functioning, pain and general health.
	Ramírez-Cervantes 2015 ¹⁶⁰	80 CD on GFD≥6 months	
	Bakker 2013 ¹⁷⁵	57 CD+T1D	
	Usai 2002 ¹⁷⁶	58 CD	
	Usai 2007 ¹⁷⁷	129 CD	
	Barratt 2011 ¹⁷²	225 CD	
	Barratt 2011 ¹⁷³	225 CD	
	Hallert 1998 ¹¹⁵	89 CD	
	Vilijamaa 2005 ¹⁶⁷	98 CD	
	Johnston 2004 ¹⁷⁸	14 CD	
	Häuser 2006 ¹⁷⁰	516 CD	
	Häuser 2006 ¹⁷¹	446 CD	
	Nachman 2009 ¹²⁸	132 Newly diagnosed CD	
	Nachman 2010 ¹⁶³	53 Newly diagnosed CD	
Hopman 2009 ¹⁷⁹	53 CD		
Tontini 2010 ¹⁸⁰	43 CD		
Aksan 2015	205 CD		
CD-specific scales			

Continued

Table 3 Continued

Adult instruments			
Instrument	Author	Participants	Instrument description
Celiac Disease Questionnaire ³⁰	Häuser 2006 ¹⁷⁰	516 CD	28-Item self-report, 7-point Likert scale. Measures four domains: emotional and social problems, disease-related worries and GI symptoms.
	Zampieron 2011 ¹⁸¹	187 CD	
	Ford 2012 ¹⁶⁵	288 CD	
	Marchese 2013 ¹⁸²	171 CD	
	Sainsbury 2013 ¹⁵⁵	189 CD	
	Pouchot 2014 ¹⁸³	211 CD	
Celiac Disease Quality of Life Survey ²⁹	Aksan 2015	205 CD	20-Item self-report, 5-point Likert scale. four subscales: limitations, dysphoria, health concerns and inadequate treatment.
	Mahadev 2016 ¹²⁷	211 CD	
	Zingone 2011 ¹⁸⁴	230 CD	
	Casellas 2013 ¹⁸⁵	298 CD	
	Casellas 2015 ¹⁵⁹	366 CD on GFD>1 year	
	Lee 2016 ¹⁸⁶	2138 CD	
	Rodriguez-Almagro 2016 ¹⁸⁷	1230 CD	
	Russo 2017 ¹⁸⁸	19 CD	
Fatigue scales	Castilhos 2015 ¹⁸⁹	103 CD	
		Dowd 2017 ¹⁹⁰	
The Chronic Fatigue Syndrome Questionnaire ¹⁹¹	Siniscalchi 2005 ¹⁹²	130 CD	8-Item self-report for physical fatigue and five items for mental fatigue. The score ranges from 0 to 26.
The Fatigue Severity Scale ¹⁹³			9-Item self-report (score range 1–7 according to patient' agreement) The scores range from 9 to 63.
Daily Fatigue Impact Scale Questionnaire ¹⁹⁴	Jordà 2010 ¹⁹⁵	51 CD	8-Item self-report questionnaire, five alternative responses per item (from 0=no problem, to 4=extreme problem).
Eating disorder scales			
Binge Eating Staircases ¹⁹⁶	Passananti 2013 ¹⁹⁷	100 CD	16-Item self-report. Measures the behavioural aspects of binge eating, as well as feelings and thoughts associated with binge eating.
Eating Attitudes Test 26 ¹⁹⁸	Passananti 2013 ¹⁹⁷	100 CD	26-Item self-report, multiple choice. Assesses diet-related disorders, bulimia and anxiety-related to food.
Eating Disorder Inventory 2 ¹⁹⁹	Passananti 2013 ¹⁹⁷	100 CD	91-Item self-report (range 0–78); 11 scales, eg, inadequacy, interoceptive awareness, fear of maturity, asceticism, impulsivity and social insecurity.
	Wagner 2015 ²⁰⁰	259 CD adolescents	
	Karwautz 2008 ²⁰¹	283 CD adolescents	
Eating Disorder Examination (EDE) ²⁰²	Wagner 2015 ²⁰⁰	259 CD adolescents	45–60 min semi-structured interview (28-day recall). Four subscales: restraint, eating concern, shape concern, weight concern.
	Karwautz 2008 ²⁰¹	283 CD adolescents	
Eating Disorder Examination Questionnaire (EDE-Q) ²⁰³	Karwautz 2008 ²⁰¹	283 CD adolescents	28-Item, self-report. Uses same subscales as the EDE interview.
Sleep disorders scale			
Pittsburgh Sleep Quality Index ²⁰⁴	Zingone 2010 ²⁰⁵	60 CD	19 self-rated items (15 multiple choice and 4 write in)+5 questions rated by a bed partner or roommate (range 0–57).
Paediatric scales			
TNO-AZL Children's QOL questionnaire (TACQOL) parent ²⁰⁶ and child ²⁰⁷ report	Kolsteren 2001 ²⁰⁸	133 CD children	Generic—two forms for children and parents: The TACQOL comprises seven scales (range0–16): ► Generic: Dutch Children TNO-AZL QOL Questionnaire 25 items with four. ► Domains: emotional, social, familiar, and physical.
	van Koppen 2009 ²⁰⁹	32 CD children	
DUX-25 ²¹⁰	van Koppen 2009 ²⁰⁹	32 CD children	(Ages 5–16) 25-item self-report with four domains: physical, emotional, social and home functioning.
Inventory of Life Quality in Children and Adolescents ²¹¹	Wagner 2008 ²¹²	365 CD adolescents	9-Item self-rating questionnaire Nine areas: school, family, social peer contacts, interests and leisure activities, physical health, psychological health, overall QoL judgement and disease and therapy-associated burden.
Berner Subjective Well-being Inventory ²¹³	Wagner 2008 ²¹²	365 CD adolescents	39-Item self-report questionnaire, with 4-point and 6-point Likert scale questions.
PedsQL core: Paediatric Quality of Life Measurement Inventory ²¹⁴	Sud 2012 ²¹⁵	28 CD+ type 1 diabetes children	Four areas: physical functioning (eight items), emotional functioning (five items), social functioning (five items) and school functioning (five items).
	Biagetti 2015 ²¹⁶	76 CD children	
	Sevinc 2016 ²¹⁷	52 CD children	
WHOQOL-BREF ²¹⁸	de Lorenzo 2012 ²¹⁹	33 CD child/parent dyads vs 62 control child/parent dyads	26-Item self-report (parent), 5-point Likert scale. Six domains: loneliness, general health perception, physical and psychological functioning, social and environmental.
Autoquestionnaire de l'Enfant Imagé -AUQUEI (child form) ²²⁰	de Lorenzo 2012 ²¹⁹	33 CD child/parent dyads vs 62 control child/parent dyads	26-Item self-report (child), explores familial and social relationships, physical activity, health, body functions and temporary separation from the familial environment.
EuroQol-5D ¹⁵⁶	Nordyke 2011 ²²¹	153 screen-detected CD children and 66 known CD vs non-CD controls	5-Item self-report, 5-point Likert scale, plus a general health rating (scored on a 20 cm visual analogue scale) (child friendly pilot and proxy used).
	Nordyke 2013 ²²²	207 CD vs non-CD controls	
DISABKIDS Chronic (short version) ²²³	Bystrom 2012 ²²⁴	160 CD child/parent dyads	12-Item self-report Likert scale for ages 8–18 (additional available version for ages 4–7) mental, social and physical domains.

Continued

Table 3 Continued

Adult instruments			
Instrument	Author	Participants	Instrument description
12-Item Short Form Survey (SF-12) ²²⁵	Altabelli 2013 ²²⁶	140 CD children	12-Item questionnaire, eight domains: physical functioning, role-physical, bodily pain, general health, vitality, social functioning, role-emotional and mental health.
Impact Scale of Childhood Diseases ²²⁷	Di Filippo 2013 ²²⁸	45 CD children	30-Item self-report, four domains: impact of disease and treatment, impact on child development and adjustment, impact on parents and impact on family.
Kidscreen ²²⁹	Myleus 2014 ²³⁰	328 CD children	52-Item self-report, measures overall HRQoL along with 10 HRQoL subdomains capturing physical well-being, psychological well-being, moods and emotions, self-perception, autonomy, parent relation and home life, financial resources, social support and peers, school environment and social acceptance (bullying); 5-point Likert scale.
Children's Depression Inventory ²³¹	Simsek 2015 ¹³⁵	25 CD children	27-Item, self-report, symptom-oriented scale (ages 7–17).
General Purpose Health-Related Quality of Life Questionnaire for Children ²³²	Simsek 2015 ¹³⁵	25 CD children	40-Item, self-report, four domains: psychological well-being, social relationships, physical function and everyday life activities.
CD-specific (paediatric) scales			
TACQOL-COE	Kolsteren 2001 ²⁰⁸	133 CD children	CD-specific instrument (based on TACQOL) TACQOL-COE-DIET: Five questions to address GFD.
CDDUX ²³³	Van Koppen 2009 ²⁰⁹	32 CD children	12-Item questionnaire, three domains: communication, having CD and diet.
	Pico 2014 ²³⁴	118 CD child/parent dyads	
	Torres 2016 ²³⁵	214 CD child/parent dyads and 52 CD children	
	Lins 2015 ²³⁶	33 CD children	
	Vriezinga 2016 ²³⁷	78 CD children	
	Meyer 2016 ²³⁸	34 CD child/parent dyads	
Celiac Disease Paediatric Quality of Life Scale ²³⁹	Jordan 2013 ²³⁹ (validation study)	181 CD children	13-Item (ages 8–12) and 17-item (ages 13–18) scale. Home, school, social and self-esteem (also diet and future in 13–18 tool). Ages 8–18 only.

CDDUX, celiac disease DUX.

DISCUSSION

Strengths and limitations

In this review, a large number of authors reviewed the literature on treatment outcomes in coeliac trials to issue recommendations for future trials. Our research team was multidisciplinary and the treatment outcomes we have evaluated in the paper reflect the expert views of the authors.

We performed an extensive literature review of more than 10 000 papers, and based our deliberations on personal experiences and expertise from our treatment of patients with CD with clinical trials and our research in CD. There are already a number of guidelines for reporting treatment outcome in CD.¹³⁶ Hence, our paper is not meant to give authoritative advice on the study design, which is not yet possible because of the developing nature of the field, but to complement available literature with our expertise with a focus on how to measure response to non-GFD treatment.

CD is a lifelong disease in which the GFD is the only treatment option. However, we suspect that soon other alternative treatments will become commonplace.

Regulatory agencies are responsible for evaluating new therapies based on risks and benefits to patients in how they feel, function or survive.⁸⁰ These aims are of intuitive value to patients and clinicians and sufficiently broad that they should form the foundation of any interventional clinical trial. Despite this, the precedent in many fields, including gastroenterology, has been the use of poorly validated outcomes of limited applicability to clinical practise. Although CD adversely impacts survival^{6 137 138} and function,^{139–143} these outcomes are generally not feasible to assess in clinical trials because of low prevalence and long latency. This leaves the options of measuring how patients feel—mainly using PROs, histology and serology—to assess changes in risk of future adverse events. Ideally, a treatment should result in improvement in more than one outcome measures (PROs, histology and serology), and that is possible using a coprimary end point. However, coprimary

end points should be used with caution since they decrease study power and the number of patients is often limited.

There are currently several well-designed and partially validated PROs developed for CD that should be considered standards for trials in the future. Assessment of extent (degree) of enteropathy (intestinal architecture and IEL assessment) should be considered as a critical outcome in clinical trials of CD. However, it is recognised that technical limitations of duodenal biopsy as a reflection of overall small intestinal mucosal disease limit the potential value of histology as an end point. Development techniques that better evaluate enteropathy across the small intestine are applicable in clinical practise and relevant to patients.

Author affiliations

¹Department of Medical Epidemiology and Biostatistics, Karolinska Institutet, Stockholm, Sweden

²Department of Pediatrics, Örebro University Hospital, Örebro, Sweden

³Coeliac Center at Department of Medicine and Surgery, Scuola Medica Salernitana, University of Salerno, Salerno, Italy

⁴Celiac Disease Center at Columbia University, New York, USA

⁵Celiac Disease Research Center, Faculty of Medicine and Life Sciences, University of Tampere, Tampere, Finland

⁶Department of Internal Medicine, Tampere University Hospital, Tampere, Finland

⁷Coeliac Disease Centre, Heim Pál Children's Hospital, Budapest, Hungary

⁸Department of Paediatrics, Faculty of Medicine, University of Debrecen, Debrecen, Hungary

⁹Celiac Disease Research Center, Faculty of Medicine and Life Sciences, University of Tampere, Tampere, Finland

¹⁰Department of Paediatrics, Tampere University Hospital, Tampere, Finland

¹¹The Mayo Clinic, Rochester, Minnesota, USA

¹²Institute of Clinical Medicine and K.G. Jebsen Coeliac Disease Research Centre, Faculty of Medicine, University of Oslo, Oslo, Norway

¹³Department of Gastroenterology, Oslo University Hospital, Oslo, Norway

¹⁴Science Center, Tampere University Hospital, Tampere, Finland

¹⁵Tampere Centre for Child Health Research, Faculty of Medicine and Life Sciences, University of Tampere, Tampere, Finland

¹⁶Institute for Mother and Child Health Bucharest, University of Medicine and Pharmacy 'Carol Davila', Bucharest, Romania

¹⁷Tampere Centre for Child Health Research, University of Tampere, Tampere University Hospital, Tampere, Finland

¹⁸Division of Pediatric Gastroenterology, Columbia University Medical Center, New York, USA

¹⁹Celiac Disease Center, Department of Medicine, Columbia University Medical Center, New York, USA

²⁰Grupo IHP Pediatría, Sevilla, Spain

²¹Academic Unit of Gastroenterology, Royal Hallamshire Hospital, University of Sheffield, Sheffield, UK

²²Celiac Center, University Medical Center, Johannes-Gutenberg University, Mainz, Germany

²³Division of Gastroenterology, Beth Israel Deaconess Medical Center, Boston, Massachusetts, USA

²⁴Coeliac UK, Buckinghamshire, UK

²⁵Tampere Centre for Child Health Research, University of Tampere, Tampere University Hospital, Tampere, Finland

²⁶Continuum Clinical, Northbrook, Illinois, USA

²⁷Faculty of Health and Medicine, School of Medicine and Public Health, University of Newcastle, Newcastle, New South Wales, Australia

²⁸Celiac Center, Beth Israel Deaconess Medical Center, Harvard Medical School, Boston, Massachusetts, USA

Contributors JFL and DL initiated the study. JFL coordinated the study and wrote the first draft, the text was then extensively revised by the coauthors. All authors contributed to the literature searches, contributed to the writing of the manuscript and approved the final version of the manuscript.

Funding JFL was supported by the Swedish Research Council (522-2A09-195) and the Swedish Society of Medicine while writing the draft of this paper. DS received research support from the German Research Foundation (DFG), the Ministry of Research and Development (BMBF) and the Leibniz Foundation. DSS received an educational grant from Biocard and Simtomax to undertake an investigator-led research study on point-of-care tests and an educational grant from Dr Schär (a gluten-free food manufacturer) to undertake an investigator-led research study on gluten sensitivity. KKa was supported by the Academy of Finland, the Competitive Research Funding of the Tampere University Hospital and The Sigrid Juselius Foundation. IRK-S was supported by the Hungarian Research Fund (Grant NKFI 120392). MJM was supported by the Competitive State Research Financing of the Tampere University Hospital (grant 9U038).

Competing interests PHRG: scientific advisory board of Alvine Pharmaceuticals, ImmunosX and ImmusanT. JAM: serves on the advisory board of Celimmune, was a consultant to BioLineRx, GlaxoSmithKline (GSK), Genentech, UCB Biopharma and Glenmark Pharmaceuticals Ltd and is a consultant to ImmunosanT, Institute for Protein Design (PvP Biologics), Takeda Pharmaceutical Company, Ltd., Innovate Biopharmaceuticals, Inc., Intrexon, 2GPharma Inc., Boeringer-Ingelheim and ImmusanT. KEAL: ImmusanT, Regeneron and Alvine Pharmaceuticals. DSS: holds a patent and receives royalties for the TG2-antibody assay, has received an educational grant from Coeliac UK, Biocard, Simtomax and Dr Schär to undertake an investigator-led research study on CD and/or gluten sensitivity. NRR: clinical advisory board for ImmusanT. DAL: Medical Director for Takeda Pharmaceuticals AR-H: coauthor detecting gluten peptides in human fluids (Patent No. US 20170160288 A1), consultant for Vircell. IRK-S: patent on rapid coeliac antibody detection licensed by the University of Tampere to Labsystems Oy, Finland. MJM: serves on the Advisory Board of Celimmune, USA, ImmusanT, USA and Innovate Pharmaceuticals Inc, USA; is consultant to FinnMedi Oy, Finland and Jilab Oy, Finland via his own company Maki HealthTech Oy, Finland; is an inventor in the patent Methods and Means for Detecting Gluten-Induced Diseases, USA (Patent No. 7,361,480—USA, European Patent No. 1390753). The patent resulted in a commercial product from FinnMedi at the Tampere University Hospital and the University of Tampere, a coeliac disease point-of-care test, Biocard Celiac Test, licensed to Labsystems Diagnostics Oy (former AniBiotech Oy), Finland.

Provenance and peer review Not commissioned; externally peer reviewed.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

© Article author(s) (or their employer(s) unless otherwise stated in the text of the article) 2018. All rights reserved. No commercial use is permitted unless otherwise expressly granted.

REFERENCES

- Lebwohl B, Ludvigsson JF, Green PH. Celiac disease and non-celiac gluten sensitivity. *BMJ* 2015;351:h4347.
- Ludvigsson JF, Michaelsson K, Ekblom A, et al. Coeliac disease and the risk of fractures - a general population-based cohort study. *Aliment Pharmacol Ther* 2007;25:273-85.
- Elfström P, Granath F, Ekström Smedby K, et al. Risk of lymphoproliferative malignancy in relation to small intestinal histopathology among patients with celiac disease. *J Natl Cancer Inst* 2011;103:436-44.
- Ilus T, Kaukinen K, Virta LJ, et al. Incidence of malignancies in diagnosed celiac patients: a population-based estimate. *Am J Gastroenterol* 2014;109:1471-7.
- Zingone F, Abdul Sultan A, Crooks CJ, et al. The risk of community-acquired pneumonia among 9803 patients with coeliac disease compared to the general population: a cohort study. *Aliment Pharmacol Ther* 2016;44:57-67.
- Ludvigsson JF, Montgomery SM, Ekblom A, et al. Small-intestinal histopathology and mortality risk in celiac disease. *JAMA* 2009;302:1171-8.
- Ludvigsson JF, Leffler DA, Bai JC, et al. The Oslo definitions for coeliac disease and related terms. *Gut* 2013;62:43-52.
- Mäki M, Mustalahti K, Kokkonen J, et al. Prevalence of Celiac disease among children in Finland. *N Engl J Med* 2003;348:2517-24.
- Walker MM, Murray JA, Ronkainen J, et al. Detection of celiac disease and lymphocytic enteropathy by parallel serology and histopathology in a population-based study. *Gastroenterology* 2010;139:112-9.
- Lohi S, Mustalahti K, Kaukinen K, et al. Increasing prevalence of coeliac disease over time. *Aliment Pharmacol Ther* 2007;26:1217-25.
- West J, Fleming KM, Tata LJ, et al. Incidence and prevalence of celiac disease and dermatitis herpetiformis in the UK over two decades: population-based study. *Am J Gastroenterol* 2014;109:757-68.
- Zingone F, Swift GL, Card TR, et al. Psychological morbidity of celiac disease: A review of the literature. *United European Gastroenterol J* 2015;3:136-45.
- Shah S, Akbari M, Vanga R, et al. Patient perception of treatment burden is high in celiac disease compared with other common conditions. *Am J Gastroenterol* 2014;109:1304-11.
- Lee AR, Ng DL, Zivin J, et al. Economic burden of a gluten-free diet. *J Hum Nutr Diet* 2007;20:423-30.
- Singh J, Whelan K. Limited availability and higher cost of gluten-free foods. *J Hum Nutr Diet* 2011;24:479-86.
- Burden M, Mooney PD, Blanshard RJ, et al. Cost and availability of gluten-free food in the UK: in store and online. *Postgrad Med J* 2015;91:622-6.
- Sollid LM, Khosla C. Novel therapies for coeliac disease. *J Intern Med* 2011;269:604-13.
- Wungjiranirun M, Kelly CP, Leffler DA. Current status of celiac disease drug development. *Am J Gastroenterol* 2016;111:779-86.
- Aziz I, Evans KE, Papageorgiou V, et al. Are patients with coeliac disease seeking alternative therapies to a gluten-free diet? *J Gastrointest Liver Dis* 2011;20:27-31.
- Hindryckx P, Levesque BG, Holvoet T, et al. Disease activity indices in coeliac disease: systematic review and recommendations for clinical trials. *Gut* 2018;67:61-9.
- Marsh MN. Gluten, major histocompatibility complex, and the small intestine. A molecular and immunobiologic approach to the spectrum of gluten sensitivity ('celiac sprue'). *Gastroenterology* 1992;102:330-54.
- Oberhuber G, Granditsch G, Vogelsang H. The histopathology of coeliac disease: time for a standardized report scheme for pathologists. *Eur J Gastroenterol Hepatol* 1999;11:1185-94.
- Rostami K, Kerckhaert J, von Blomberg BM, et al. SAT and serology in adult coeliacs, seronegative coeliac disease seems a reality. *Neth J Med* 1998;53:15-19.
- Villanacci V, Magazzù G, Pellegrino S, et al. Comparison of the marsh-oberhuber classification with a new grading system in identifying patients with latent celiac disease. *Minerva Gastroenterol Dietol* 2010;56:371-5.
- Ensari A. Gluten-sensitive enteropathy (celiac disease): controversies in diagnosis and classification. *Arch Pathol Lab Med* 2010;134:826-36.
- Taavela J, Koskinen O, Huhtala H, et al. Validation of morphometric analyses of small-intestinal biopsy readouts in celiac disease. *PLoS One* 2013;8:e76163.
- Leffler DA, Kelly CP, Green PH, et al. Larazotide acetate for persistent symptoms of celiac disease despite a gluten-free diet: a randomized controlled trial. *Gastroenterology* 2015;148:1311-9.
- Adelman DC, Leffler D, Lebwohl B, et al. Celiac disease symptom frequency and severity using a disease-specific patient-reported outcome diary: observations from a psychometric validation study in 202 patients. *Am J Gastro* 2012;107:S603.
- Dorn SD, Hernandez L, Minaya MT, et al. The development and validation of a new coeliac disease quality of life survey (CD-QOL). *Aliment Pharmacol Ther* 2010;31:666-75.
- Häuser W, Gold J, Stallmach A, et al. Development and validation of the Celiac Disease Questionnaire (CDQ), a disease-specific health-related quality of life measure for adult patients with celiac disease. *J Clin Gastroenterol* 2007;41:157-66.
- Leffler DA, Dennis M, Edwards George J, et al. A validated disease-specific symptom index for adults with celiac disease. *Clin Gastroenterol Hepatol* 2009;7:1328e1-1334.
- Lee SK, Lo W, Memeo L, et al. Duodenal histology in patients with celiac disease after treatment with a gluten-free diet. *Gastrointest Endosc* 2003;57:187-91.
- Tursi A, Brandimarte G, Giorgetti GM, et al. Endoscopic and histological findings in the duodenum of adults with celiac disease before and after changing to a gluten-free diet: a 2-year prospective study. *Endoscopy* 2006;38:702-7.
- Cammarota G, Cuomo L, Cesaro P, et al. A highly accurate method for monitoring histological recovery in patients with celiac disease on a gluten-free diet using

- an endoscopic approach that avoids the need for biopsy: a double-center study. *Endoscopy* 2007;39:46–51.
- 35 Maiden L, Elliott T, McLaughlin SD, *et al.* A blinded pilot comparison of capsule endoscopy and small bowel histology in unresponsive celiac disease. *Dig Dis Sci* 2009;54:1280–3.
 - 36 Oxford Centre for Evidence-based Medicine. *Levels of evidence*. Oxford: Oxford Centre for Evidence-based Medicine, 2009.
 - 37 Liberati A, Altman DG, Tetzlaff J, *et al.* The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate healthcare interventions: explanation and elaboration. *BMJ* 2009;339:b2700.
 - 38 Rubio-Tapia A, Hill ID, Kelly CP, *et al.* ACG clinical guidelines: diagnosis and management of celiac disease. *Am J Gastroenterol* 2013;108:656–76.
 - 39 Ludvigsson JF, Bai JC, Biagi F, *et al.* Diagnosis and management of adult coeliac disease: guidelines from the British Society of Gastroenterology. *Gut* 2014;63:1210–28.
 - 40 Husby S, Koletzko S, Korponay-Szabó IR, *et al.* European society for pediatric gastroenterology, hepatology, and nutrition guidelines for the diagnosis of coeliac disease. *J Pediatr Gastroenterol Nutr* 2012;54:136–60.
 - 41 Corazza GR, Villanacci V. Coeliac disease. *J Clin Pathol* 2005;58:573–4.
 - 42 Gottlieb K, Dawson J, Hussain F, *et al.* Development of drugs for celiac disease: review of endpoints for Phase 2 and 3 trials. *Gastroenterol Rep* 2015;3:91–102.
 - 43 Babbini BA, Crawford K, Sitaraman SV. Malabsorption work-up: utility of small bowel biopsy. *Clin Gastroenterol Hepatol* 2006;4:1193–8.
 - 44 Latorre M, Lagana SM, Freedberg DE, *et al.* Endoscopic biopsy technique in the diagnosis of celiac disease: one bite or two? *Gastrointest Endosc* 2015;81:1228–33.
 - 45 Ravelli A, Villanacci V. Tricks of the trade: how to avoid histological pitfalls in celiac disease. *Pathol Res Pract* 2012;208:197–202.
 - 46 Rostami K, Marsh MN, Johnson MW, *et al.* ROC-king onwards: intraepithelial lymphocyte counts, distribution & role in coeliac disease mucosal interpretation. *Gut* 2017;66:2080–6.
 - 47 Veress B, Franzén L, Bodin L, *et al.* Duodenal intraepithelial lymphocyte count revisited. *Scand J Gastroenterol* 2004;39:138–44.
 - 48 Järvinen TT, Kaukinen K, Laurila K, *et al.* Intraepithelial lymphocytes in celiac disease. *Am J Gastroenterol* 2003;98:1332–7.
 - 49 Lonardi S, Villanacci V, Lorenzi L, *et al.* Anti-TCR gamma antibody in celiac disease: the value of count on formalin-fixed paraffin-embedded biopsies. *Virchows Arch* 2013;463:409–13.
 - 50 Zanini B, Marullo M, Villanacci V, *et al.* Persistent intraepithelial lymphocytosis in celiac patients adhering to gluten-free diet is not abolished despite a gluten contamination elimination diet. *Nutrients* 2016;8:525.
 - 51 Korponay-Szabó IR, Halttunen T, Szalai Z, *et al.* In vivo targeting of intestinal and extraintestinal transglutaminase 2 by coeliac autoantibodies. *Gut* 2004;53:641–8.
 - 52 Koskinen O, Collin P, Lindfors K, *et al.* Usefulness of small-bowel mucosal transglutaminase-2 specific autoantibody deposits in the diagnosis and follow-up of celiac disease. *J Clin Gastroenterol* 2010;44:483–8.
 - 53 Gatti S, Rossi M, Alfonsi S, *et al.* Beyond the intestinal celiac mucosa: diagnostic role of anti-TG2 deposits, a systematic review. *Front Med* 2014;1:9.
 - 54 Lähdeaho ML, Mäki M, Laurila K, *et al.* Small-bowel mucosal changes and antibody responses after low- and moderate-dose gluten challenge in celiac disease. *BMC Gastroenterol* 2011;11:129.
 - 55 Arquelles-Grande C, Tennyson CA, Lewis SK, *et al.* Variability in small bowel histopathology reporting between different pathology practice settings: impact on the diagnosis of coeliac disease. *J Clin Pathol* 2012;65:242–7.
 - 56 Picarelli A, Borghini R, Donato G, *et al.* Weaknesses of histological analysis in celiac disease diagnosis: new possible scenarios. *Scand J Gastroenterol* 2014;49:1–7.
 - 57 Dixon MF, Genta RM, Yardley JH, *et al.* Classification and grading of gastritis. The updated Sydney system. International workshop on the histopathology of gastritis, Houston 1994. *Am J Surg Pathol* 1996;20:1161–81.
 - 58 Malfertheiner P, Megraud F, O'Morain CA, *et al.* Management of helicobacter pylori infection—the maastricht v/florence consensus report. *Gut* 2017;66:6–30.
 - 59 Leibold B, Green PH, Genta RM. The coeliac stomach: gastritis in patients with coeliac disease. *Aliment Pharmacol Ther* 2015;42:180–7.
 - 60 Verzegnassi F, Bua J, De Angelis P, *et al.* Eosinophilic oesophagitis and coeliac disease: is it just a casual association? *Gut* 2007;56:1029–30.
 - 61 Trovato CM, Montuori M, Anania C, *et al.* Are ESPGHAN “biopsy-sparing” guidelines for celiac disease also suitable for asymptomatic patients? *Am J Gastroenterol* 2015;110:1485–9.
 - 62 Ghazzawi Y, Rubio-Tapia A, Murray JA, *et al.* Mucosal healing in children with treated celiac disease. *J Pediatr Gastroenterol Nutr* 2014;59:229–31.
 - 63 Leffler DA, Schuppan D. Update on serologic testing in celiac disease. *Am J Gastroenterol* 2010;105:2520–4.
 - 64 Bai JC, Fried M, Corazza GR, *et al.* World gastroenterology organisation global guidelines on celiac disease. *J Clin Gastroenterol* 2013;47:121–6.
 - 65 Reeves GE, Squance ML, Duggan AE, *et al.* Diagnostic accuracy of coeliac serological tests: a prospective study. *Eur J Gastroenterol Hepatol* 2006;18:493–501.
 - 66 Naiyer AJ, Hernandez L, Ciaccio EJ, *et al.* Comparison of commercially available serologic kits for the detection of celiac disease. *J Clin Gastroenterol* 2009;43:225–32.
 - 67 Simon-Vecsei Z, Király R, Bagossi P, *et al.* A single conformational transglutaminase 2 epitope contributed by three domains is critical for celiac antibody binding and effects. *Proc Natl Acad Sci U S A* 2012;109:431–6.
 - 68 Iversen R, Mysling S, Hnida K, *et al.* Activity-regulating structural changes and autoantibody epitopes in transglutaminase 2 assessed by hydrogen/deuterium exchange. *Proc Natl Acad Sci U S A* 2014;111:17146–51.
 - 69 Kanchan K, Ergülen E, Király R, *et al.* Identification of a specific one amino acid change in recombinant human transglutaminase 2 that regulates its activity and calcium sensitivity. *Biochem J* 2013;455:261–72.
 - 70 Alessio MG, Tonutti E, Brusca I, *et al.* Correlation between IgA tissue transglutaminase antibody ratio and histological finding in celiac disease. *J Pediatr Gastroenterol Nutr* 2012;55:44–9.
 - 71 Taavela J, Kurppa K, Collin P, *et al.* Degree of damage to the small bowel and serum antibody titers correlate with clinical presentation of patients with celiac disease. *Clin Gastroenterol Hepatol* 2013;11:166–71.
 - 72 Singh P, Kurrray L, Agnihotri A, *et al.* Titers of anti-tissue transglutaminase antibody correlate well with severity of villous abnormalities in celiac disease. *J Clin Gastroenterol* 2015;49:212–7.
 - 73 Pekki H, Kurppa K, Mäki M, *et al.* Predictors and significance of incomplete mucosal recovery in celiac disease after 1 year on a gluten-free diet. *Am J Gastroenterol* 2015;110:1078–85.
 - 74 Silvester JA, Kurada S, Sz wajcer A, *et al.* Tests for serum transglutaminase and endomysial antibodies do not detect most patients with celiac disease and persistent villous atrophy on gluten-free diets: a meta-analysis. *Gastroenterology* 2017;153:689–701.
 - 75 Murray JA, Rubio-Tapia A, Van Dyke CT, *et al.* Mucosal atrophy in celiac disease: extent of involvement, correlation with clinical presentation, and response to treatment. *Clin Gastroenterol Hepatol* 2008;6:186–93.
 - 76 Hopper AD, Hadjivassiliou M, Hurlstone DP, *et al.* What is the role of serologic testing in celiac disease? A prospective, biopsy-confirmed study with economic analysis. *Clin Gastroenterol Hepatol* 2008;6:314–20.
 - 77 Leffler D, Schuppan D, Pallav K, *et al.* Kinetics of the histological, serological and symptomatic responses to gluten challenge in adults with coeliac disease. *Gut* 2013;62:996–1004.
 - 78 Sama VK, Skodje GI, Reims HM, *et al.* HLA-DQ:gluten tetramer test in blood gives better detection of coeliac patients than biopsy after 14-day gluten challenge. *Gut* 2017;gutjnl-2017-314461.
 - 79 Matysiak-Budnik T, Malamut G, de Serre NP, *et al.* Long-term follow-up of 61 coeliac patients diagnosed in childhood: evolution toward latency is possible on a normal diet. *Gut* 2007;56:1379–86.
 - 80 Leffler D, Kupfer SS, Leibold B, *et al.* Development of celiac disease therapeutics: report of the third gastroenterology regulatory endpoints and advancement of therapeutics workshop. *Gastroenterology* 2016;151:407–11.
 - 81 Forsberg G, Hernell O, Melgar S, *et al.* Paradoxical coexpression of proinflammatory and down-regulatory cytokines in intestinal T cells in childhood celiac disease. *Gastroenterology* 2002;123:667–78.
 - 82 Brottveit M, Beitnes AC, Tollefsen S, *et al.* Mucosal cytokine response after short-term gluten challenge in celiac disease and non-celiac gluten sensitivity. *Am J Gastroenterol* 2013;108:842–50.
 - 83 Kontakou M, Przemioslo RT, Sturgess RP, *et al.* Cytokine mRNA expression in the mucosa of treated coeliac patients after wheat peptide challenge. *Gut* 1995;37:52–7.
 - 84 Goel G, King T, Daveson AJ, *et al.* Epitope-specific immunotherapy targeting CD4-positive T cells in coeliac disease: two randomised, double-blind, placebo-controlled phase 1 studies. *Lancet Gastroenterol Hepatol* 2017;2:479–93.
 - 85 Ontiveros N, Tye-Din JA, Hardy MY, *et al.* Ex-vivo whole blood secretion of interferon (IFN)- γ and IFN- γ -inducible protein-10 measured by enzyme-linked immunosorbent assay are as sensitive as IFN- γ enzyme-linked immunospot for the detection of gluten-reactive T cells in human leucocyte antigen (HLA)-DQ2-5(+)-associated coeliac disease. *Clin Exp Immunol* 2014;175:305–15.
 - 86 Manavalan JS, Hernandez L, Shah JG, *et al.* Serum cytokine elevations in celiac disease: association with disease presentation. *Hum Immunol* 2010;71:50–7.
 - 87 Lundin KE, Scott H, Hansen T, *et al.* Gliadin-specific, HLA-DQ(alpha 1*0501,beta 1*0201) restricted T cells isolated from the small intestinal mucosa of celiac disease patients. *J Exp Med* 1993;178:187–96.
 - 88 Lundin KE, Scott H, Fausa O, *et al.* T cells from the small intestinal mucosa of a DR4, DQ7/DR4, DQ8 celiac disease patient preferentially recognize gliadin when presented by DQ8. *Hum Immunol* 1994;41:285–91.
 - 89 van de Wal Y, Kooy YM, van Veelen PA, *et al.* Small intestinal T cells of celiac disease patients recognize a natural pepsin fragment of gliadin. *Proc Natl Acad Sci U S A* 1998;95:10050–4.
 - 90 Jabri B, Sollid LM. T cells in celiac disease. *J Immunol* 2017;198:3005–14.
 - 91 Anderson RP, Degano P, Godkin AJ, *et al.* In vivo antigen challenge in celiac disease identifies a single transglutaminase-modified peptide as the dominant A-gliadin T-cell epitope. *Nat Med* 2000;6:337–42.
 - 92 Anderson RP, van Heel DA, Tye-Din JA, *et al.* T cells in peripheral blood after gluten challenge in coeliac disease. *Gut* 2005;54:1217–23.

- 93 Quarsten H, McAdam SN, Jensen T, *et al.* Staining of celiac disease-relevant T cells by peptide-DQ2 multimers. *J Immunol* 2001;167:4861–8.
- 94 Råki M, Fallang LE, Brotteit M, *et al.* Tetramer visualization of gut-homing gluten-specific T cells in the peripheral blood of celiac disease patients. *Proc Natl Acad Sci U S A* 2007;104:2831–6.
- 95 Brotteit M, Råki M, Bergseng E, *et al.* Assessing possible celiac disease by an HLA-DQ2-gliadin tetramer test. *Am J Gastroenterol* 2011;106:1318–24.
- 96 Christophersen A, Råki M, Bergseng E, *et al.* Tetramer-visualized gluten-specific CD4+ T cells in blood as a potential diagnostic marker for coeliac disease without oral gluten challenge. *United European Gastroenterol J* 2014;2:268–78.
- 97 Dahal-Koirala S, Risnes LF, Christophersen A, *et al.* TCR sequencing of single cells reactive to DQ2.5-gli-2 and DQ2.5-gli-2 reveals clonal expansion and epitope-specific V-gene usage. *Mucosal Immunol* 2016;9:587–96.
- 98 Garber ME, Saldanha A, Parker JS, *et al.* A B-cell gene signature correlates with the extent of gluten-induced intestinal injury in celiac disease. *Cell Mol Gastroenterol Hepatol* 2017;4:1–17.
- 99 Lähdeaho ML, Kaukinen K, Laurila K, *et al.* Glutenase ALV003 attenuates gluten-induced mucosal injury in patients with celiac disease. *Gastroenterology* 2014;146:1649–58.
- 100 Moreno ML, Rodríguez-Herrera A, Sousa C, *et al.* Biomarkers to monitor gluten-free diet compliance in celiac patients. *Nutrients* 2017;9:46.
- 101 Shan L, Qiao SW, Arentz-Hansen H, *et al.* Identification and analysis of multivalent proteolytically resistant peptides from gluten: implications for celiac sprue. *J Proteome Res* 2005;4:1732–41.
- 102 Shan L, Molberg Ø, Parrot I, *et al.* Structural basis for gluten intolerance in celiac sprue. *Science* 2002;297:2275–9.
- 103 Comino I, Real A, Vivas S, *et al.* Monitoring of gluten-free diet compliance in celiac patients by assessment of gliadin 33-mer equivalent epitopes in feces. *Am J Clin Nutr* 2012;95:670–7.
- 104 Comino I, Fernández-Bañares F, Esteve M, *et al.* Fecal gluten peptides reveal limitations of serological tests and food questionnaires for monitoring gluten-free diet in celiac disease patients. *Am J Gastroenterol* 2016;111:1456–65.
- 105 Moreno ML, Cebolla Á, Muñoz-Suano A, *et al.* Detection of gluten immunogenic peptides in the urine of patients with coeliac disease reveals transgressions in the gluten-free diet and incomplete mucosal healing. *Gut* 2017;66:250–7.
- 106 Sulic AM, Kurppa K, Rauhavirta T, *et al.* Transglutaminase as a therapeutic target for celiac disease. *Expert Opin Ther Targets* 2015;19:335–48.
- 107 *Guidance for industry and FDA staff qualification process for drug development tools*. US: U.S. Department of Health and Human Services FaDA, Center for Drug Evaluation and Research, 2014.
- 108 Riaz MS, Atreja A. Personalized technologies in chronic gastrointestinal disorders: self-monitoring and remote sensor technologies. *Clin Gastroenterol Hepatol* 2016;14:1697–705.
- 109 *Claims GfIP-ROMUiMPDtSL*. US: U.S. Department of Health and Human Services FaDA, Center for Drug Evaluation and Research, Center for Biologics Evaluation and Research, Health CfDaR, 2009.
- 110 Kelly CP, Green PH, Murray JA, *et al.* Larazotide acetate in patients with coeliac disease undergoing a gluten challenge: a randomised placebo-controlled study. *Aliment Pharmacol Ther* 2013;37:252–62.
- 111 Lohiniemi S, Mäki M, Kaukinen K, *et al.* Gastrointestinal symptoms rating scale in coeliac disease patients on wheat starch-based gluten-free diets. *Scand J Gastroenterol* 2000;35:947–9.
- 112 Murray JA, Kelly CP, Green PH, *et al.* No difference between latiglutenase and placebo in reducing villous atrophy or improving symptoms in patients with symptomatic celiac disease. *Gastroenterology* 2017;152:787–98.
- 113 Crocker H, Jenkinson C, Churchman D, *et al.* The Coeliac Disease Assessment Questionnaire (Cdaq): development of a patient-reported outcome measure. *Value in Health* 2016;19:A595.
- 114 Zanini B, Basché R, Ferraresi A, *et al.* Randomised clinical study: gluten challenge induces symptom recurrence in only a minority of patients who meet clinical criteria for non-coeliac gluten sensitivity. *Aliment Pharmacol Ther* 2015;42:968–76.
- 115 Hallert C, Grännö C, Grant C, *et al.* Quality of life of adult coeliac patients treated for 10 years. *Scand J Gastroenterol* 1998;33:977–8.
- 116 Kurppa K, Paavola A, Collin P, *et al.* Benefits of a gluten-free diet for asymptomatic patients with serologic markers of celiac disease. *Gastroenterology* 2014;147:610–7.
- 117 Svedlund J, Sjödin I, Dotevall G. GSRs—a clinical rating scale for gastrointestinal symptoms in patients with irritable bowel syndrome and peptic ulcer disease. *Dig Dis Sci* 1988;33:129–34.
- 118 Hendrickx GF, Somers K, Vandenplas Y. Lane-Hamilton syndrome: case report and review of the literature. *Eur J Pediatr* 2011;170:1597–602.
- 119 Silvester JA, Graff LA, Rigaux L, *et al.* Symptomatic suspected gluten exposure is common among patients with coeliac disease on a gluten-free diet. *Aliment Pharmacol Ther* 2016;44:612–9.
- 120 Mahadev S, Simpson S, Lebowohl B, *et al.* Is dietitian use associated with celiac disease outcomes? *Nutrients* 2013;5:1585–94.
- 121 Crocker HPM, Jenkinson C. The Coeliac Disease Assessment Questionnaire (CDAQ). 2016.
- 122 Lahiff C, Safaie P, Awais A, *et al.* The Crohn's disease activity index (CDAI) is similarly elevated in patients with Crohn's disease and in patients with irritable bowel syndrome. *Aliment Pharmacol Ther* 2013;37:786–94.
- 123 Mahadev S, Murray JA, Wu TT, *et al.* Factors associated with villus atrophy in symptomatic coeliac disease patients on a gluten-free diet. *Aliment Pharmacol Ther* 2017;45:1084–93.
- 124 Fritscher-Ravens A, Schuppan D, Ellrichmann M, *et al.* Confocal endomicroscopy shows food-associated changes in the intestinal mucosa of patients with irritable bowel syndrome. *Gastroenterology* 2014;147:1012–20.
- 125 Zanini B, Caselani F, Magni A, *et al.* Celiac disease with mild enteropathy is not mild disease. *Clin Gastroenterol Hepatol* 2013;11:253–8.
- 126 Lanzini A, Lanzarotto F, Villanacci V, *et al.* Complete recovery of intestinal mucosa occurs very rarely in adult coeliac patients despite adherence to gluten-free diet. *Aliment Pharmacol Ther* 2009;29:1299–308.
- 127 Mahadev S, Gardner R, Lewis SK, *et al.* Quality of life in screen-detected celiac disease patients in the United States. *J Clin Gastroenterol* 2016;50:393–7.
- 128 Nachman F, Mauriño E, Vázquez H, *et al.* Quality of life in celiac disease patients: prospective analysis on the importance of clinical severity at diagnosis and the impact of treatment. *Dig Liver Dis* 2009;41:15–25.
- 129 Burger JPW, de Brouwer B, Int'Hout J, *et al.* Systematic review with meta-analysis: dietary adherence influences normalization of health-related quality of life in coeliac disease. *Clin Nutr* 2017;36:399–406.
- 130 Paavola A, Kurppa K, Ukkola A, *et al.* Gastrointestinal symptoms and quality of life in screen-detected celiac disease. *Dig Liver Dis* 2012;44:814–8.
- 131 Mustalhti K, Lohiniemi S, Collin P, *et al.* Gluten-free diet and quality of life in patients with screen-detected celiac disease. *Eff Clin Pract* 2002;5:105–13.
- 132 Ukkola A, Mäki M, Kurppa K, *et al.* Diet improves perception of health and well-being in symptomatic, but not asymptomatic, patients with celiac disease. *Clin Gastroenterol Hepatol* 2011;9:118–23.
- 133 Fera T, Cascio B, Angelini G, *et al.* Affective disorders and quality of life in adult coeliac disease patients on a gluten-free diet. *Eur J Gastroenterol Hepatol* 2003;15:1287–92.
- 134 Addolorato G, Capristo E, Ghittoni G, *et al.* Anxiety but not depression decreases in coeliac patients after one-year gluten-free diet: a longitudinal study. *Scand J Gastroenterol* 2001;36:502–6.
- 135 Simsek S, Baysog G, Gencoglan S, *et al.* Effects of gluten-free diet on quality of life and depression in children with celiac disease. *J Pediatr Gastroenterol Nutr* 2015;61:303–6.
- 136 Moher D, Hopewell S, Schulz KF, *et al.* Consort 2010 explanation and elaboration: updated guidelines for reporting parallel group randomised trials. *BMJ* 2010;340:c869.
- 137 Tio M, Cox MR, Eslick GD. Meta-analysis: coeliac disease and the risk of all-cause mortality, any malignancy and lymphoid malignancy. *Aliment Pharmacol Ther* 2012;35:540–51.
- 138 Biagi F, Corazza GR. Mortality in celiac disease. *Nat Rev Gastroenterol Hepatol* 2010;7:158–62.
- 139 Heikkilä K, Pearce J, Mäki M, *et al.* Celiac disease and bone fractures: a systematic review and meta-analysis. *J Clin Endocrinol Metab* 2015;100:25–34.
- 140 Lebowohl B, Michaëlsson K, Green PH, *et al.* Persistent mucosal damage and risk of fracture in celiac disease. *J Clin Endocrinol Metab* 2014;99:609–16.
- 141 Lichtwark IT, Newnham ED, Robinson SR, *et al.* Cognitive impairment in coeliac disease improves on a gluten-free diet and correlates with histological and serological indices of disease severity. *Aliment Pharmacol Ther* 2014;40:160–70.
- 142 Hadjivassiliou M, Rao DG, Grinewald RA, *et al.* Neurological dysfunction in coeliac disease and non-coeliac gluten sensitivity. *Am J Gastroenterol* 2016;111:561–7.
- 143 Catassi C, Fasano A. Celiac disease as a cause of growth retardation in childhood. *Curr Opin Pediatr* 2004;16:445–9.
- 144 Tack GJ, van de Water JM, Bruins MJ, *et al.* Consumption of gluten with gluten-degrading enzyme by celiac patients: a pilot-study. *World J Gastroenterol* 2013;19:5837–47.
- 145 Catassi C, Fabiani E, Iacono G, *et al.* A prospective, double-blind, placebo-controlled trial to establish a safe gluten threshold for patients with celiac disease. *Am J Clin Nutr* 2007;85:160–6.
- 146 Holm K, Mäki M, Vuolteenaho N, *et al.* Oats in the treatment of childhood coeliac disease: a 2-year controlled trial and a long-term clinical follow-up study. *Aliment Pharmacol Ther* 2006;23:1463–72.
- 147 Catassi C, Rossini M, Rättsch IM, *et al.* Dose dependent effects of protracted ingestion of small amounts of gliadin in celiac disease children: a clinical and jejunal morphometric study. *Gut* 1993;34:1515–9.
- 148 Leffler DA, Kelly CP, Abdallah HZ, *et al.* A randomized, double-blind study of larazotide acetate to prevent the activation of celiac disease during gluten challenge. *Am J Gastroenterol* 2012;107:1554–62.
- 149 Zung WW. A self-rating depression scale. *Arch Gen Psychiatry* 1965;12:63–70.
- 150 Ciacchi C, D'Agate C, De Rosa A, *et al.* Self-rated quality of life in celiac disease. *Dig Dis Sci* 2003;48:2216–20.
- 151 Quintana JM, Cabriada J, López de Tejada I, *et al.* Translation and validation of the gastrointestinal Quality of Life Index (GIQLI). *Rev Esp Enferm Dig* 2001;93:693–706.

- 152 Casellas F, Rodrigo L, Vivancos JL, *et al.* Factors that impact health-related quality of life in adults with coeliac disease: a multicenter study. *World J Gastroenterol* 2008;14:46–52.
- 153 Carver CS. You want to measure coping but your protocol's too long: consider the brief COPE. *Int J Behav Med* 1997;4:92–100.
- 154 Smith MM, Goodfellow L. The relationship between quality of life and coping strategies of adults with coeliac disease adhering to a gluten-free diet. *Gastroenterol Nurs* 2011;34:460–8.
- 155 Sainsbury K, Mullan B, Sharpe L. Reduced quality of life in coeliac disease is more strongly associated with depression than gastrointestinal symptoms. *J Psychosom Res* 2013;75:135–41.
- 156 Brooks RG, Rabin R, De Charro F. *The measurement and valuation of health status using EQ-5D: a European perspective: evidence from the EuroQol BIOMED Research Programme*. Dordrecht; Boston: Kluwer Academic Pub, 2003.
- 157 Gray AM, Papanicolaou IN. Impact of symptoms on quality of life before and after diagnosis of coeliac disease: results from a UK population survey. *BMC Health Serv Res* 2010;10:105.
- 158 Norström F, Lindholm L, Sandström O, *et al.* Delay to coeliac disease diagnosis and its implications for health-related quality of life. *BMC Gastroenterol* 2011;11:118.
- 159 Casellas F, Rodrigo L, Lucendo AJ, *et al.* Benefit on health-related quality of life of adherence to gluten-free diet in adult patients with coeliac disease. *Rev Esp Enferm Dig* 2015;107:196–201.
- 160 Ramírez-Cervantes KL, Remes-Troche JM, Del Pilar Milke-García M, *et al.* Characteristics and factors related to quality of life in Mexican Mestizo patients with coeliac disease. *BMC Gastroenterol* 2015;15:4.
- 161 Upton J. Beck Depression Inventory (BDI). In: Gellman MD, Turner JR, eds. *Encyclopedia of behavioral medicine*. New York, NY: Springer New York, 2013:178–9.
- 162 Borghini R, Di Tola M, Salvi E, *et al.* Impact of gluten-free diet on quality of life in coeliac patients. *Acta Gastroenterol Belg* 2016;79:447–53.
- 163 Nachman F, del Campo MP, González A, *et al.* Long-term deterioration of quality of life in adult patients with coeliac disease is associated with treatment noncompliance. *Dig Liver Dis* 2010;42:685–91.
- 164 Dupuy H, *et al.* The Psychological General Wellbeing (PGWB) Index. In: Wenger N, Mattson M, Furberg C, Elinson J, eds. *Assessment of quality of life in clinical trials of cardiovascular therapies*:170–83.
- 165 Ford S, Howard R, Oyebo J. Psychosocial aspects of coeliac disease: a cross-sectional survey of a UK population. *Br J Health Psychol* 2012;17:743–57.
- 166 Peräaho M, Kaukinen K, Mustalahti K, *et al.* Effect of an oats-containing gluten-free diet on symptoms and quality of life in coeliac disease. A randomized study. *Scand J Gastroenterol* 2004;39:27–31.
- 167 Viljamaa M, Collin P, Huhtala H, *et al.* Is coeliac disease screening in risk groups justified? A four-year follow-up with special focus on compliance and quality of life. *Aliment Pharmacol Ther* 2005;22:317–24.
- 168 Paarlahti P, Kurppa K, Ukkola A, *et al.* Predictors of persistent symptoms and reduced quality of life in treated coeliac disease patients: a large cross-sectional study. *BMC Gastroenterol* 2013;13:75.
- 169 Zigmund AS, Snaith RP. The hospital anxiety and depression scale. *Acta Psychiatr Scand* 1983;67:361–70.
- 170 Häuser W, Stallmach A, Caspary WF, *et al.* Predictors of reduced health-related quality of life in adults with coeliac disease. *Aliment Pharmacol Ther* 2007;25:569–78.
- 171 Häuser W, Gold J, Stein J, *et al.* Health-related quality of life in adult coeliac disease in Germany: results of a national survey. *Eur J Gastroenterol Hepatol* 2006;18:747–54.
- 172 Barratt SM, Leeds JS, Robinson K, *et al.* Reflux and irritable bowel syndrome are negative predictors of quality of life in coeliac disease and inflammatory bowel disease. *Eur J Gastroenterol Hepatol* 2011;23:159–65.
- 173 Barratt SM, Leeds JS, Sanders DS. Quality of life in coeliac disease is determined by perceived degree of difficulty adhering to a gluten-free diet, not the level of dietary adherence ultimately achieved. *J Gastrointest Liver Dis* 2011;20:241–5.
- 174 Ware JE, Sherbourne CD. The MOS 36-item short-form health survey (SF-36). I. Conceptual framework and item selection. *Med Care* 1992;30:473–83.
- 175 Bakker SF, Pouwer F, Tushuizen ME, *et al.* Compromised quality of life in patients with both Type 1 diabetes mellitus and coeliac disease. *Diabet Med* 2013;30:835–9.
- 176 Usai P, Minerba L, Marini B, *et al.* Case control study on health-related quality of life in adult coeliac disease. *Dig Liver Dis* 2002;34:547–52.
- 177 Usai P, Manca R, Cuomo R, *et al.* Effect of gluten-free diet and co-morbidity of irritable bowel syndrome-type symptoms on health-related quality of life in adult coeliac patients. *Dig Liver Dis* 2007;39:824–8.
- 178 Johnston SD, Rodgers C, Watson RG. Quality of life in screen-detected and typical coeliac disease and the effect of excluding dietary gluten. *Eur J Gastroenterol Hepatol* 2004;16:1281–6.
- 179 Hopman EG, Koopman HM, Wit JM, *et al.* Dietary compliance and health-related quality of life in patients with coeliac disease. *Eur J Gastroenterol Hepatol* 2009;21:1056–61.
- 180 Tontini GE, Rondonotti E, Saladino V, *et al.* Impact of gluten withdrawal on health-related quality of life in coeliac subjects: an observational case-control study. *Digestion* 2010;82:221–8.
- 181 Zampieron A, Daicampi C, Martin A, *et al.* Quality of life in adult coeliac disease in a mountain area of northeast Italy. *Gastroenterol Nurs* 2011;34:313–9.
- 182 Marchese A, Klersy C, Biagi F, *et al.* Quality of life in coeliac patients: Italian validation of a coeliac questionnaire. *Eur J Intern Med* 2013;24:87–91.
- 183 Pouchot J, Despujol C, Malamut G, *et al.* Validation of a French version of the quality of life "coeliac disease questionnaire". *PLoS One* 2014;9:e96346.
- 184 Zingone F, Iavarone A, Tortora R, *et al.* The Italian translation of the coeliac disease-specific quality of life scale in coeliac patients on gluten free diet. *Dig Liver Dis* 2013;45:115–8.
- 185 Casellas F, Rodrigo L, Molina-Infante J, *et al.* Transcultural adaptation and validation of the Coeliac Disease Quality of Life (CD-QoL) Survey, a specific questionnaire to measure quality of life in patients with coeliac disease. *Rev Esp Enferm Dig* 2013;105:585–93.
- 186 Lee AR, Wolf R, Contento I, *et al.* Coeliac disease: the association between quality of life and social support network participation. *J Hum Nutr Diet* 2016;29:383–90.
- 187 Rodríguez Almagro J, Hernández Martínez A, Lucendo AJ, *et al.* Health-related quality of life and determinant factors in coeliac disease. A population-based analysis of adult patients in Spain. *Rev Esp Enferm Dig* 2016;108:181–9.
- 188 Russo F, Chimienti G, Clemente C, *et al.* A possible role for ghrelin, leptin, brain-derived neurotrophic factor and docosahexaenoic acid in reducing the quality of life of coeliac disease patients following a gluten-free diet. *Eur J Nutr* 2017;56:807–18.
- 189 Castilhos AC, Gonçalves BC, Silva MM, *et al.* Quality of life evaluation in coeliac patients from southern Brazil. *Arq Gastroenterol* 2015;52:171–5.
- 190 Dowd AJ, Jung ME. Self-compassion directly and indirectly predicts dietary adherence and quality of life among adults with coeliac disease. *Appetite* 2017;113:293–300.
- 191 Wessely S, Powell R. Fatigue syndromes: a comparison of chronic "postviral" fatigue with neuromuscular and affective disorders. *J Neurol Neurosurg Psychiatry* 1989;52:940–8.
- 192 Siniscalchi M, Iovino P, Tortora R, *et al.* Fatigue in adult coeliac disease. *Aliment Pharmacol Ther* 2005;22:489–94.
- 193 Krupp LB, LaRocca NG, Muir-Nash J, *et al.* The fatigue severity scale. Application to patients with multiple sclerosis and systemic lupus erythematosus. *Arch Neurol* 1989;46:1121–3.
- 194 Fisk JD, Doble SE. Construction and validation of a fatigue impact scale for daily administration (D-FIS). *Qual Life Res* 2002;11:263–72.
- 195 Jordá FC, López Vivancos J. Fatigue as a determinant of health in patients with coeliac disease. *J Clin Gastroenterol* 2010;44:423–7.
- 196 Gormally J, Black S, Daston S, *et al.* The assessment of binge eating severity among obese persons. *Addict Behav* 1982;7:47–55.
- 197 Passananti V, Siniscalchi M, Zingone F, *et al.* Prevalence of eating disorders in adults with coeliac disease. *Gastroenterol Res Pract* 2013;2013:1–7.
- 198 Garner DM, Olmsted MP, Bohr Y, *et al.* The eating attitudes test: psychometric features and clinical correlates. *Psychol Med* 1982;12:871–8.
- 199 Garner DM, Olmstead MP, Polivy J. Development and validation of a multidimensional eating disorder inventory for anorexia nervosa and bulimia. *Int J Eat Disord* 1983;2:15–34.
- 200 Wagner G, Zeiler M, Berger G, *et al.* Eating disorders in adolescents with coeliac disease: influence of personality characteristics and coping. *Eur Eat Disord Rev* 2015;23:361–70.
- 201 Karwautz A, Wagner G, Berger G, *et al.* Eating pathology in adolescents with coeliac disease. *Psychosomatics* 2008;49:399–406.
- 202 Fairburn CG. *Cognitive behavior therapy and eating disorders*. New York, NY: Guilford Press, 2008.
- 203 Luce KH, Crowther JH, Pole M. Eating disorder examination questionnaire (EDE-Q): norms for undergraduate women. *Int J Eat Disord* 2008;41:273–6.
- 204 Buysse DJ, Reynolds CF, Monk TH, *et al.* The pittsburgh sleep quality index: a new instrument for psychiatric practice and research. *Psychiatry Res* 1989;28:193–213.
- 205 Zingone F, Siniscalchi M, Capone P, *et al.* The quality of sleep in patients with coeliac disease. *Aliment Pharmacol Ther* 2010;32:1031–6.
- 206 Vogels T, Verrips GH, Verloove-Vanhorick SP, *et al.* Measuring health-related quality of life in children: the development of the TACQOL parent form. *Qual Life Res* 1998;7:457–65.
- 207 Kamphuis R, Theunissen N, Witb J. Short communication health-related quality of life measure for children-the TACQOL. *J Appl Therapeut* 1997;1:357–60.
- 208 Kolsteren MM, Koopman HM, Schalekamp G, *et al.* Health-related quality of life in children with coeliac disease. *J Pediatr* 2001;138:593–5.
- 209 van Koppen EJ, Schweizer JJ, Cszizmadia CG, *et al.* Long-term health and quality-of-life consequences of mass screening for childhood coeliac disease: a 10-year follow-up study. *Pediatrics* 2009;123:e582–588.
- 210 Koopman H, Theunissen N, Vogels A, *et al.* The DUC-25: a short-form questionnaire for measuring health related quality of life of children with a chronic illness. *Quality of Life Research* 1998;6:19–19.
- 211 Matthejat F, Remschmidt H. The inventory of life quality in children and adolescents (ILC). *Z Kinder-Jugendpsychiatrie* 1998;26:183–96.

- 212 Wagner G, Berger G, Sinnreich U, *et al.* Quality of life in adolescents with treated coeliac disease: influence of compliance and age at diagnosis. *J Pediatr Gastroenterol Nutr* 2008;47:555–61.
- 213 Grob A, Lüthi R, Kaiser FG, *et al.* Berner fragebogen zum wohlfinden jugendlicher (BFW). *Diagnostica* 1991;37:66–75.
- 214 Varni JW, Seid M, Rode CA. The PedsQL: measurement model for the pediatric quality of life inventory. *Med Care* 1999;37:126–39.
- 215 Sud S, Marcon M, Assor E, *et al.* Quality of life in children with diabetes and coeliac disease: minimal impact of the 'double diagnosis'. *Pediatr Diabetes* 2012;13:163–9.
- 216 Biagetti C, Gesuita R, Gatti S, *et al.* Quality of life in children with coeliac disease: a paediatric cross-sectional study. *Dig Liver Dis* 2015;47:927–32.
- 217 Sevinç E, Çetin FH, Coşkun BD. Psychopathology, quality of life, and related factors in children with coeliac disease. *J Pediatr* 2017;93:267–73.
- 218 Anon. Development of the World Health Organization WHOQOL-BREF quality of life assessment. The WHOQOL Group. *Psychol Med* 1998;28:551–8.
- 219 de Lorenzo CM, Xikota JC, Wayhs MC, *et al.* Evaluation of the quality of life of children with coeliac disease and their parents: a case-control study. *Qual Life Res* 2012;21:77–85.
- 220 Manificat S, Dazord A, Cochat P, *et al.* [Evaluation of the quality of life in pediatrics: how to collect the point of view of children]. *Arch Pediatr* 1997;4:1238–46.
- 221 Nordyke K, Norström F, Lindholm L, *et al.* Health-related quality-of-life in children with coeliac disease, measured prior to receiving their diagnosis through screening. *J Med Screen* 2011;18:187–92.
- 222 Nordyke K, Norström F, Lindholm L, *et al.* Health-related quality of life in adolescents with screening-detected coeliac disease, before and one year after diagnosis and initiation of gluten-free diet, a prospective nested case-referent study. *BMC Public Health* 2013;13:142.
- 223 Baars RM, Atherton CI, Koopman HM, *et al.* The European DISABKIDS project: development of seven condition-specific modules to measure health related quality of life in children and adolescents. *Health Qual Life Outcomes* 2005;3:70.
- 224 Byström IM, Hollén E, Fälth-Magnusson K, *et al.* Health-related quality of life in children and adolescents with coeliac disease: from the perspectives of children and parents. *Gastroenterol Res Pract* 2012;2012:1–6.
- 225 Turner-Bowker D, Hogue SJ. Short form 12 health survey (sf-12). Michalos AC, ed. *Encyclopedia of quality of life and well-being research*. Dordrecht: Springer Netherlands, 2014:5954–7.
- 226 Altobelli E, Paduano R, Gentile T, *et al.* Health-related quality of life in children and adolescents with coeliac disease: survey of a population from central Italy. *Health Qual Life Outcomes* 2013;11:204.
- 227 Hoare P, Russell M. The quality of life of children with chronic epilepsy and their families: preliminary findings with a new assessment measure. *Dev Med Child Neurol* 1995;37:689–96.
- 228 Di Filippo T, Orlando MF, Concialdi G, *et al.* The quality of life in developing age children with coeliac disease. *Minerva Pediatr* 2013;65:599–608.
- 229 Ravens-Sieberer U, Gosch A, Rajmil L, *et al.* KIDSCREEN-52 quality-of-life measure for children and adolescents. *Expert Rev Pharmacoecon Outcomes Res* 2005;5:353–64.
- 230 Myléus A, Petersen S, Carlsson A, *et al.* Health-related quality of life is not impaired in children with undetected as well as diagnosed coeliac disease: a large population based cross-sectional study. *BMC Public Health* 2014;14:425.
- 231 Kovacs M. The Children's Depression, Inventory (CDI). *Psychopharmacol Bull* 1985;21:995–8.
- 232 Eser E, Yüksel H, Baydur H, *et al.* [The psychometric properties of the new Turkish generic health-related quality of life questionnaire for children (Kid-KINDL)]. *Turk Psikiyatri Derg* 2008;19:409–17.
- 233 van Doorn RK, Winkler LM, Zwinderman KH, *et al.* CDDUX: a disease-specific health-related quality-of-life questionnaire for children with coeliac disease. *J Pediatr Gastroenterol Nutr* 2008;47:147–52.
- 234 Pico M, Spirito MF. Implementation of a health-related quality of life questionnaire for children and adolescents with coeliac disease. *Arch Argent Pediatr* 2014;112:19–25.
- 235 Torres JB, Román E, Cilleruelo M, *et al.* Health-related quality of life in spanish children with coeliac disease. *J Pediatr Gastroenterol Nutr* 2016;62:603–8.
- 236 Lins MTC, Tassitano RM, Brandt KG, *et al.* Translation, cultural adaptation, and validation of the coeliac disease DUX (CDDUX). *J Pediatr* 2015;91:448–54.
- 237 Vriezinga SL, Farih N, van der Meulen-de Jong AE, *et al.* Comparison of patients' and doctors' reports on health-related quality of life in coeliac disease. *J Pediatr Gastroenterol Nutr* 2017;64:737–41.
- 238 Meyer S, Rosenblum S. Children with coeliac disease: health-related quality of life and leisure participation. *Am J Occup Ther* 2016;70:7006220010p1.
- 239 Jordan NE, Li Y, Magrini D, *et al.* Development and validation of a coeliac disease quality of life instrument for North American children. *J Pediatr Gastroenterol Nutr* 2013;57:477–86.