OPEN ACCESS

ORIGINAL ARTICLE

Colesevelam attenuates cholestatic liver and bile duct injury in *Mdr2*^{-/-} mice by modulating composition, signalling and excretion of faecal bile acids

Claudia Daniela Fuchs,¹ Gustav Paumgartner,¹ Veronika Mlitz,¹ Victoria Kunczer,¹ Emina Halilbasic,¹ Nadja Leditznig,¹ Annika Wahlström,² Marcus Ståhlman,² Andrea Thüringer,³ Karl Kashofer,³ Tatjana Stojakovic,⁴ Hanns-Ulrich Marschall,² Michael Trauner¹

► Additional material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/gutjnl-2017-314553>).

¹Hans Popper Laboratory of Molecular Hepatology, Division of Gastroenterology and Hepatology, Department of Internal Medicine III, Medical University of Vienna, Vienna, Austria

²Department of Molecular and Clinical Medicine, Sahlgrenska Academy, University of Gothenburg, Gothenburg, Sweden

³Institute of Pathology, Medical University of Graz, Graz, Austria

⁴Clinical Institute of Medical and Chemical Laboratory Diagnostics, Medical University of Graz, Graz, Austria

Correspondence to

Professor Michael Trauner, Division of Gastroenterology and Hepatology, Department of Internal Medicine III, Medical University of Vienna, Vienna 1090, Austria; michael.trauner@meduniwien.ac.at

Received 24 May 2017

Revised 20 February 2018

Accepted 3 March 2018

Published Online First

10 April 2018

► <http://dx.doi.org/10.1136/gutjnl-2018-316375>

To cite: Fuchs CD, Paumgartner G, Mlitz V, et al. *Gut* 2018;**67**:1683–1691.

ABSTRACT

Background and aims Interruption of the enterohepatic circulation of bile acids (BAs) may protect against BA-mediated cholestatic liver and bile duct injury. BA sequestrants are established to treat cholestatic pruritus, but their impact on the underlying cholestasis is still unclear. We aimed to explore the therapeutic effects and mechanisms of the BA sequesterant colesevelam in a mouse model of sclerosing cholangitis.

Methods *Mdr2*^{-/-} mice received colesevelam for 8 weeks. Gene expression profiles of BA homeostasis, inflammation and fibrosis were explored in liver, intestine and colon. Hepatic and faecal BA profiles and gut microbiome were analysed. Glucagon-like peptide 1 (GLP-1) levels in portal blood were measured by ELISA. Furthermore, *Mdr2*^{-/-} mice as well as wild-type 3,5-diethoxy-carbonyl-1,4-dihydrocollidine-fed mice were treated with GLP-1-receptor agonist exendin-4 for 2 weeks prior to analysis.

Results Colesevelam reduced serum liver enzymes, BAs and expression of proinflammatory and profibrogenic markers. Faecal BA profiling revealed increased levels of secondary BAs after resin treatment, while hepatic and biliary BA composition showed a shift towards more hydrophilic BAs. Colonic GLP-1 secretion, portal venous GLP-1 levels and intestinal messenger RNA expression of gut hormone *Proglucagon* were increased, while ileal *Fgf15* expression was abolished by colesevelam. Exendin-4 treatment increased bile duct mass without promoting a reactive cholangiocyte phenotype in mouse models of sclerosing cholangitis. Microbiota analysis showed an increase of the phylum δ -Proteobacteria after colesevelam treatment and a shift within the phyla Firmicutes from *Clostridiales* to *Lactobacillus*.

Conclusion Colesevelam increases faecal BA excretion and enhances BA conversion towards secondary BAs, thereby stimulating secretion of GLP-1 from enteroendocrine L-cells and attenuates liver and bile duct injury in *Mdr2*^{-/-} mice.

INTRODUCTION

Bile acid (BA) sequestrants such as colesevelam are anion exchange resins that complex BAs in the intestinal lumen, thus decreasing BA reuptake and enterohepatic circulation. BA sequestrants have

Significance of this study

What is already known on this subject?

- Bile acid (BA) sequestrants (resins) are used to treat cholestatic pruritus; in single cases, improvement of cholestasis was observed.
- Resin-bound BAs stimulate the G-protein-coupled BA receptor Takeda G protein-coupled receptor 5 (TGR5) which increases glucagon-like peptide 1 (GLP-1) secretion.
- GLP-1 signalling was shown to stimulate cholangiocyte proliferation and to protect bile duct epithelia against apoptosis.

What are the new findings?

- Colesevelam treatment improved hepatic inflammation, fibrosis and ductular proliferation in the *Mdr2*^{-/-} mouse model of sclerosing cholangitis.
- Colesevelam feeding reduced colonic farnesoid X receptor messenger RNA and protein expression in *Mdr2*^{-/-} mice.
- Colesevelam feeding result in detoxification of the hepatic as well as the biliary BA composition.
- Colesevelam feeding changed faecal BA profiles towards TGR5 signalling favouring increased colonic expression of GLP-1 and increasing its levels in portal blood, potentially contributing to the improvement of cholestatic liver injury in *Mdr2*^{-/-} mice.

How might it impact on clinical practice in the foreseeable future?

- Beyond their current use in pruritus BA sequestrants (with a longstanding and excellent safety track record) may represent a cheap and readily available treatment strategy for underlying cholestatic liver injury.

been primarily used for lowering serum cholesterol and treatment of cholestasis-associated pruritus¹ as well as BA-induced diarrhoea.² Interestingly, colesevelam has been reported to improve serum BA levels³ and colestyramine to improve cholestatic liver enzymes in patients with primary sclerosing

cholangitis (PSC),⁴ suggesting potential anticholestatic and bile duct-protective properties. Resins stimulate BA signalling via the Takeda G protein-coupled receptor 5 (TGR5)⁵ while repressing colonic farnesoid X receptor (FXR),⁶ thereby promoting secretion of glucagon-like peptide 1 (GLP-1) from enteroendocrine L-cells.⁶ Notably, the GLP-1 analogue exendin-4 has also cholangioprotective effects in primary cholangiocytes as well as in bile duct ligated rats,^{7,8} in addition to its beneficial metabolic effects.⁹ Although both inhibitions of apical sodium-dependent bile salt transporter (ASBT) as well as administration of BA sequestrants result in interruption of enterohepatic circulation of BAs and improvement of cholestatic features,^{4,10,11} their mechanism of action may differ. In contrast to complex binding of BAs through resins, the recently developed inhibitors of ASBT enhance the excretion of unbound BAs into intestine, where they can signal intracellularly via FXR, thereby inhibiting GLP-1 expression/secretion.⁶ We hypothesise that BA resin treatment results in reduced liver and bile duct injury in the *Mdr2*^{-/-} mouse (as a model of sclerosing cholangitis) by modulating faecal as well as biliary BA content and composition, favouring formation of endogenous BAs that stimulate TGR5, which in turn induces cholangioprotective GLP-1 production.

MATERIALS AND METHODS

Animals

FVB/N wild-type (WT) and *Mdr2*^{-/-} mice obtained from Jackson Laboratory (Bar Harbor, Maine, USA) were housed in a 12 hours light/dark house facility with water and standard chow diet (SSNIFF, Soest, Germany) ad libitum.

Feeding protocols

Colesevelam (BA sequestrant) was a generous gift from Dr Alan F Hofmann. Eight weeks old male *Mdr2*^{-/-} mice received either control diet or a diet supplemented with 2% (w/w) colesevelam for 8 weeks.

In another series, *Mdr2*^{-/-} mice were injected once daily with exendin-4 (Sigma-Aldrich, Vienna, Austria) (0.1 µg/kg) for 2 weeks. Eight-week-old male FVB/N WT mice receiving 0.1% 3,5-diethoxycarbonyl-1,4-dihydrocollidine (DDC) for 17 days were injected with exendin-4 (0.1 µg/kg) once daily. The experimental protocols were approved by the local Animal Care and Use Committee (BMFW-66.009/0315-WF/3b/2014).

Routine serum biochemistry and histology

Serum biochemistry and histological staining was performed as described previously.¹⁰

Immunohistochemistry

Detection of hepatic cytokeratin (CK) 19, F4/80, Ki67, vascular cell adhesion protein 1 (VCAM-1) and osteopontin (OPN) was performed as described previously.^{12,13} Immunohistochemistry (IHC) was also performed for GLP-1 in colon slices. In *Mdr2*^{-/-} mice, CK19, VCAM-1, OPN and GLP-1 IHC positive areas were quantified by ImageJ software V.1.47v (National Institute of Health). In WT DDC-fed mice, gene expression profiles were used for quantification of these markers since DDC feeding results in accumulation of protoporphyrin plaques in bile ducts that give false positive areas with computational quantification.

Figure 1 Colesevelam feeding improves liver injury in *Mdr2*^{-/-} mice. (A) Representative H&E images (10× magnification) with markedly improved liver histology with reduced pericholangitis and onion skin type of biliary fibrosis in colesevelam fed *Mdr2*^{-/-} mice. (B) Serum biochemistry reflects improved levels of transaminases (ALT, AST) as well as AP and bile acids (BAs). (C) Liver to body weight ratio (LW/BW) improved due to colesevelam feeding. *Significant difference from *Mdr2*^{-/-} control group; *p*<0.05. ALT, alanine aminotransferase; AP, alkaline phosphatase; AST, aspartate aminotransferase; BA, bile acid.

Figure 2 Ductular proliferation is improved by colessevelam treatment. (A) Representative CK19 immunohistochemistry pictures (10× magnification) show reduced cholangiocyte proliferation in colessevelam-treated *Mdr2*^{-/-} mice. (B) Real-time PCR was used to assess the mRNA expression of cholangiocyte proliferation marker *CK19* which was reduced by colessevelam feeding. Expression values were normalised against *18sRNA* levels and are shown relative to expression level in *Mdr2*^{-/-} control mice. *Significant difference from *Mdr2*^{-/-} control group; *p*<0.05. mRNA, messenger RNA.

Ki67 positive cells were counted manually in five different power fields in all experimental settings.

Hepatic hydroxyproline content

To quantify liver fibrosis, hepatic hydroxyproline was measured from a standardised liver lobe as described previously.¹⁴

Hepatic, biliary and faecal bile acid analysis

Hepatic, biliary and faecal BA profiles were acquired using ultra-performance liquid chromatography tandem mass spectrometry as described previously.¹⁵

16s rRNA microbiome analysis

16s rRNA microbiome analysis was performed as described previously.¹⁶

Messenger RNA analysis and PCR

RNA isolation from liver and intestine, complementary DNA synthesis and real-time PCR were performed as described previously.¹⁷ Oligonucleotide sequences are available on request.

Western blot analysis

Protein expression was quantified as described previously.¹⁸

GLP-1 ELISA

Blood was collected from portal vein. The GLP-1 (Active 7–36) ELISA (Catalogue Number: 43-GP1HU-E01, ALPCO) was conducted according to the manufacturer's instructions.

Statistical analysis

Results were evaluated using SPSS V.20.0. Statistical analysis was performed using student's *t*-test. Data were reported as means of five to seven animals per group ±SD. A *p* value ≤0.05 was considered significant.

RESULTS

Colessevelam treatment improves cholestatic liver injury in *Mdr2*^{-/-} mice

Colessevelam was well tolerated in *Mdr2*^{-/-} mice without affecting animal behaviour, food intake or body weight after 8 weeks of administration (see online supplementary figure 1). Importantly, mice did not develop diarrhoea. Colessevelam treatment even reduced the daily stool volume from 1.52 g/mouse/day to

1.03 g/mouse/day (cumulative output). Liver histology revealed complete reversion of bile duct injury (pericholangitis and onion skin type of biliary fibrosis) in colessevelam-treated *Mdr2*^{-/-} mice (figure 1A, all mice shown in online supplementary figure 3). Additional experimental series confirmed that colessevelam treatment in *Mdr2*^{-/-} mice led to complete reversion of the liver phenotype similar to that of WT mice (online supplementary figure 2). Accordingly, serum parameters ALT, AST and BAs as well as liver to body weight ratios of treated *Mdr2*^{-/-} mice were in the same range as in WT mice (figure 1 and online supplementary figure 2). Importantly, resin treatment reduced bile duct proliferation as determined by immunohistochemistry (IHC) (figure 2A) and messenger RNA (mRNA) expression of CK 19 (figure 2B). Together, these findings implicate a beneficial role of BA sequestrant feeding on cholestatic liver and bile duct injury in a mouse model of sclerosing cholangitis (figures 1 and 2).

Colessevelam treatment attenuates hepatic inflammation and fibrosis in *Mdr2*^{-/-} mice

Because the reactive cholangiocyte phenotype in cholangiopathies is associated with development of hepatic inflammation and fibrosis, markers for both key processes in the progression of liver injury were investigated (figures 3 and 4). Representative images of F4/80 IHC displayed reduced numbers of macrophages in livers of *Mdr2*^{-/-} mice fed with colessevelam (figure 3A). Accordingly, *F4/80* mRNA levels were reduced in the treatment group in comparison to *Mdr2*^{-/-} control mice (figure 3B). Gene expression profiles of *Mcp1* and *Vcam-1* (two proinflammatory markers of liver injury in *Mdr2*^{-/-} mice) also revealed profound reduction of these genes in colessevelam fed mice (figure 3B). Sirius red staining (figure 4A) displayed a marked reduction of fibrosis in *Mdr2*^{-/-} mice subjected to colessevelam. Accordingly, transcription of profibrogenic genes such as *Col1a1* and *Col1a2* were considerably reduced in the treatment group (figure 4B). In line, hepatic hydroxyproline content as well as α-SMA protein levels were attenuated in the colessevelam fed *Mdr2*^{-/-} mice (figure 4C and online supporting figure 4A).

Colessevelam feeding alters bile acid homeostasis in *Mdr2*^{-/-} mice

In line with effective BA depletion, colessevelam feeding reduced hepatic FXR protein expression (figure 5A) and increased mRNA expression of *Cyp7a1* (the rate limiting enzyme in BA synthesis)

Figure 3 Colesevelam feeding improves hepatic inflammation in *Mdr2*^{-/-} mice. (A) Representative F4/80 immunohistochemistry images (10× magnification) showing reduced numbers of macrophages in the livers of *Mdr2*^{-/-} mice after colesevelam treatment. (B) Real-time PCR was used to assess the mRNA expression of *F4/80*, *Mcp1* and *Vcam-1* as markers of inflammation which were reduced due to treatment with colesevelam. Expression values were normalised against *18 sRNA* levels and are shown relative to expression level in *Mdr2*^{-/-} controls. * Significant difference from *Mdr2*^{-/-} control group; $p < 0.05$.

sevenfold in *Mdr2*^{-/-} mice (figure 5B). Notably, the sinusoidal BA uptake machinery Na^+ /taurocholate cotransporting polypeptide was increased at protein level by colesevelam feeding while expression of the canalicular BA export pump bile salt export pump did not differ between the groups (see online supporting figure 4B). Ileal mRNA expression of *Fxra* (figure 5C), its target hormone *Fgf15* and *Proglucagon* was also assessed (figure 5D). Ileal *Fxra* mRNA expression was reduced by colesevelam and in line, *Fgf15* mRNA was completely abolished under resin treatment, while *Proglucagon* was induced fivefold (figure 5D), indicating that colesevelam feeding may have differential impact on FXR and TGR5-dependent regulation of gene expression. Notably, gene expression of the inflammatory marker *Mcp1* was reduced in the terminal ileum of colesevelam fed *Mdr2*^{-/-} mice (figure 5D). Of particular interest, *Proglucagon* mRNA expression is also increased in the colon of colesevelam-treated *Mdr2*^{-/-} mice (figure 5E).

Bile flow as well as biliary HCO_3^- and BA output were not influenced by colesevelam feeding (figure 6A-C). Rather, hepatic and biliary BA concentrations increased, 8-fold and 1.4-fold (data not shown), respectively, while their profiles showed a shift towards a more hydrophilic BA composition with taurine-conjugated muricholic acids (MCAs) consisting 51% and 56%, respectively (figure 6D and E).

Colesevelam treatment modifies faecal BA output and composition

Resin feeding profoundly increased faecal BA output (1276.8 $\mu\text{mol/day}$ in *Mdr2*^{-/-} control mice and 6656.3 $\mu\text{mol/day}$ in *Mdr2*^{-/-} colesevelam-fed mice; cumulative faeces output). Importantly, in spite of a fivefold increase in faecal BA output *Mdr2*^{-/-} mice subjected to colesevelam did not develop diarrhoea. At the same time, the resin significantly altered faecal BA composition by increasing concentrations of primary BAs α/β MCA, cholic acid (CA) and chenodeoxycholic acid as well as of secondary BAs lithocholic acid (LCA) and deoxycholic acid (DCA) (table 1). Of note, the two most potent endogenous TGR5 ligands, LCA and DCA,¹⁹ represented 51% of total BAs in the faeces of colesevelam fed *Mdr2*^{-/-} mice, compared with 19% in the untreated group. Moreover, colesevelam-fed *Mdr2*^{-/-} mice had 91% unconjugated BAs in the faeces compared with 83% in *Mdr2*^{-/-} control mice (table 1).

Colesevelam treatment modifies gut microbiome

16sRNA microbiome analysis revealed an increase in the phylum Proteobacteria in *Mdr2*^{-/-} mice treated with colesevelam (9% vs 5% in the control group) (see online supplementary figure 5A). More specifically, this increase resulted from elevated abundance

Figure 4 Biliary fibrosis is reduced in *Mdr2*^{-/-} mice by colesesevelam treatment. (A) Representative sirius red stainings (10 $\times$ magnification) show improved biliary fibrosis in *Mdr2*^{-/-} mice fed with colesesevelam. (B) Real-time PCR was used to assess the mRNA expression of fibrotic markers *Col1a1* and *Col1a2* which were reduced by colesesevelam feeding. Expression values were normalised against *18sRNA* levels and are shown relative to expression level in *Mdr2*^{-/-} controls. (C) OH-proline levels were assessed and turned out to be improved in *Mdr2*^{-/-} mice subjected to colesesevelam treatment. *Significant difference from *Mdr2*^{-/-} control group; $p < 0.05$.

of the class δ -Proteobacteria (order Desulfobacterales, family Desulfobacteraceae) (see online supplementary figure 5B). Despite unchanged Firmicutes levels (*Mdr2*^{-/-} control group 38%; *Mdr2*^{-/-} colesesevelam group 36%), in the colesesevelam fed group a shift within the phylum Firmicutes was seen, from the class Clostridiales to the class Bacilli, more specifically from the order Clostridiales to the order Lactobacillales, known to consist several species exerting bile salt hydrolase (BSH) activity, such as *Lactobacillus*²⁰ (see online supplementary figure 5B). Also the phylum Bacteroidetes remained unchanged in *Mdr2*^{-/-} control and colesesevelam fed mice (56% vs 54%) (see online supplementary figure 5A). Of particular interest, although we see increased levels of secondary BAs in the faeces of colesesevelam-fed *Mdr2*^{-/-} mice, the class Clostridia (containing bacteria exhibiting 7 α dehydroxylase activity) was predominant in untreated *Mdr2*^{-/-} control mice (see online supplementary figure 5B).

Colesesevelam treatment increases colonic GLP-1 signalling in *Mdr2*^{-/-} mice

Since increased mRNA levels of small intestinal and colonic *Proglucagon* and elevated secondary BAs LCA and DCA in colesesevelam-fed *Mdr2*^{-/-} mice point towards an augmented

TGR5–GLP-1 signalling, colonic GLP-1 expression was assessed by IHC. *Mdr2*^{-/-} mice fed with colesesevelam exhibited increased numbers of GLP-1 secreting enteroendocrine L-cells (figure 7A). Accordingly, GLP-1 concentrations in portal blood of BA resin fed animals tended to be increased (figure 7B). Importantly (given the differential impact of FXR vs TGR5 on GLP-1 secretion), mRNA (figure 7C) as well as protein expression levels of (proximal) colonic FXR were reduced in colesesevelam-fed *Mdr2*^{-/-} mice compared with controls (figure 7D). Also, the expression of the FXR downstream target *Ibafp* was reduced at mRNA level (data not shown).

GLP-1 receptor agonist exendin-4 promotes bile duct formation

To further address a potential role of GLP-1 in improvement of liver and biliary injury,^{7,8} we treated *Mdr2*^{-/-} mice and WT mice fed with DDC¹³ (as mouse models of sclerosing cholangitis) with exendin-4 as GLP-1 receptor agonist.⁷ In both models, exendin-4 increased expression of CK19 and Ki67 (see online supplementary figures 6 and 8), while at the same time VCAM-1 and OPN remained unchanged (see online supplementary figures 7 and 9).

Figure 5 Colesevelam feeding impacts on BA homeostasis in *Mdr2*^{-/-} mice. (A) Hepatic FXR protein expression is reduced in colesevelam fed *Mdr2*^{-/-} mice (Histone 3 was used as internal control), while (B) *Cyp7a1* (key enzyme for BA synthesis) mRNA expression is profoundly increased. (C) Ileal *Fxrα* mRNA expression is reduced in colesevelam-fed *Mdr2*^{-/-} mice. (D) Ileal gene expression of *Fgf15* was completely abolished in colesevelam-fed mice, while *Proglucagon* levels were induced fivefold. Expression of inflammatory marker *Mcp1* is reduced in the terminal ileum of colesevelam-fed *Mdr2*^{-/-} mice. (E) Colonic gene expression of *Proglucagon* was increased due to colesevelam feeding. Expression values were normalised against *18sRNA* (liver) and *Gapdh* (ileum) and *18sRNA* (colon) levels and are shown relative to expression level in *Mdr2*^{-/-} controls. *Significant difference from *Mdr2*^{-/-} control group; *p* < 0.05. BA, bile acid; FXR, farnesoid X receptor.

DISCUSSION

In this study, we uncover that the BA binding resin colesevelam reverses cholestatic liver and bile duct injury in the *Mdr2*^{-/-} mouse model of sclerosing cholangitis (figure 1 and online supporting figures 2 and 3). Our findings provide strong evidence that resins such as colesevelam may have important implications as potential therapeutic strategies for cholestatic liver disease beyond their role in the treatment of pruritus. In this study, we now provide a potential mechanistic explanation for anecdotal clinical observations that resins may in addition to itch also improve cholestasis.¹ Importantly, these mechanisms appear to be different from more costly and side effect prone ASBT inhibitors that recently were propagated for the treatment of cholestatic pruritus.^{21 22}

The improvement of the cholangiopathy in the *Mdr2*^{-/-} mouse subjected to colesevelam may at least in part be explained by interruption of the enterohepatic circulation with reduced output of potentially toxic BAs.^{10 11} Importantly, faecal BA analysis showed profound increases in BA excretion due to colesevelam treatment and enhanced conversion of primary to secondary BAs (table 1) with subsequent effects on BA signalling in the gut. As such, we observed markedly increased GLP-1 expression in enteroendocrine L-cells (figure 7A) and elevated GLP-1 levels in portal blood of colesevelam fed *Mdr2*^{-/-} mice (figure 7B), pointing towards increased TGR5 activity as a result of increased formation of secondary BAs. Notably,

enteroendocrine L-cell FXR activity and TGR5-mediated GLP-1 secretion appear to be inversely related.^{5 6} Mice lacking FXR showed elevated levels of *Proglucagon* mRNA and elevated serum GLP-1 levels.⁶ Moreover, colesevelam treatment of *ob/ob* mice was found to inhibit *Fgf15* mRNA expression in ileum (further arguing for reduced FXR activity) and increased colonic *Proglucagon* expression.⁶ In line, in our study small intestinal and colonic FXR mRNA and protein expression was reduced in *Mdr2*^{-/-} mice subjected to colesevelam feeding while at the same time colonic GLP-1 expression and portal blood levels of this hormone were increased (figures 5 and 7).

GLP-1 released from the colon, entering the liver via portal blood, may bind to its receptor at cholangiocytes and modulate their adaptive response to cholestasis, by reducing cholangiocyte apoptosis while at the same time favouring their proliferation, thereby preventing ductopenia, a unifying event in cholangiopathies.^{7 8 23} This phenomenon would be predicted to protect cholangiocytes from BA-induced stress. GLP-1 docking to its receptor increases cAMP production, which results in an anti-inflammatory response due to the inhibition of NFκB and STAT1 expression, thereby reducing expression of inflammatory cytokines and chemokines as well as adhesion molecules such as *Tnfa*, *Vcam-1* and *Mcp1*,²⁴ known to contribute the reactive cholangiocyte phenotype.²⁵ This hypothesis is at least partially supported by our findings in *Mdr2*^{-/-} mice as well as in DDC-fed WT mice which were injected with the GLP-1 receptor agonist

Figure 6 Colesevelam treatment does not influence bile flow but changes bile composition. (A) Hepatobiliary bile flow (B) biliary HCO_3^- output and (C) biliary BA output remain unchanged under colesevelam treatment. (D) Hepatic and (E) biliary BA profile showed a shift towards a more hydrophilic BA composition with taurine-conjugated MCAs being the most prominent species. BA, bile acid; MCA, muricholic acid; TCA, taurine-conjugated cholic acid; TCDCA, taurine-conjugated deoxycholic acid; T α MCA, tauro-alpha-muricholic acid; T β MCA, tauro-beta-muricholic acid; T ω MCA, tauro-omega-muricholic acid; TUDCA, taurine-conjugated ursodeoxycholic acid.

extendin-4. In both mouse models, bile duct mass (CK19 expression and staining) as well as cell proliferation (Ki67 staining) were increased due to extendin-4 injection while at the same time,

markers for the reactive (proinflammatory/profibrogenic) cholangiocyte phenotype (VCAM-1 and OPN) remained unchanged (see online supporting figures 6-9). Stimulation of cholangiocyte proliferation without promotion of a reactive cholangiocyte phenotype could be viewed as a potentially protective mechanistic counteracting ductopaenia.

In addition to increased GLP-1 signalling, also changes in hepatic and biliary BA composition due to colesevelam feeding may add to the overall beneficial effect seen in livers of resin-treated $Mdr2^{-/-}$ mice. As such, colesevelam feeding led to a 'hydrophilisation' (thereby detoxification) of hepatic as well as biliary BA composition with predominance of taurine-conjugated α -MCAs, β -MCAs and ω -MCAs (figure 6). Interestingly, this 'hydrophilisation' was not seen in $Mdr2^{-/-}$ mice treated with an ASBT inhibitor,¹⁰ as reflected by the negligible concentration of biliary hydrophilic muricholic acid species while concentration of hydrophobic TDCA was increased.¹⁰ In addition to differences of GLP-1 signalling between ASBT inhibition and resin feeding, these divergent effects on biliary BA composition may significantly contribute to the mechanistic differences between these two treatment strategies.

The divergent effects on GLP-1 secretion might be explained by the fact that in ASBT inhibition, largely unmodified BAs enter the colon which can be reabsorbed and then activate intracellular FXR-dependent signalling pathways reported to suppress GLP-1 expression.⁶ Resin-bound BAs cannot be reabsorbed to the same extent and thus are still able to signal via the G protein-coupled receptor TGR5, thereby stimulating GLP-1 secretion. In line with this hypothesis, we found reduced colonic FXR expression in colesevelam-treated $Mdr2^{-/-}$ mice (figure 7C,D), which could contribute to increased GLP-1 expression,⁶ in addition

Table 1 Faecal bile acid levels

	<i>Mdr2</i> ^{-/-} control (pmol/mg)	<i>Mdr2</i> ^{-/-} colesevelam (pmol/mg)
α MCA	21.4 $\pm$ 11.6	399.8 $\pm$ 34.2*
β MCA	125.3 $\pm$ 39.8	450 $\pm$ 82.2*
ω MCA	334.4 $\pm$ 192.3	300.1 $\pm$ 59.1
CA	47.7 $\pm$ 32.2	1141.9 $\pm$ 371.9*
CDCA	4.6 $\pm$ 3.6	238.5 $\pm$ 65.1*
DCA	142.7 $\pm$ 63.3	3072.7 $\pm$ 914.8*
UDCA	2.2 $\pm$ 1.1	19.1 $\pm$ 3.8*
LCA	15.1 $\pm$ 5.9	265.8 $\pm$ 107.0*
T α MCA	2.2 $\pm$ 1.2	42.0 $\pm$ 40.9*
T β MCA	58.4 $\pm$ 23.2	79.3 $\pm$ 40.9
T ω MCA	10.0 $\pm$ 4.5	3.5 $\pm$ 1.6*
TCA	51.4 $\pm$ 15.9	226.2 $\pm$ 153.2
TCDCA	4.2 $\pm$ 3.7	135.2 $\pm$ 76.2*
TDCA	0.8 $\pm$ 0.8	49.4 $\pm$ 40.9*
TUDCA	0.8 $\pm$ 0.6	3.3 $\pm$ 1.7*
TLCA	nd	3.5 $\pm$ 2.2*
Total BAs	839.9 $\pm$ 280.3	6475.6 $\pm$ 1202.6*

*Significant difference from *Mdr2*^{-/-} control; p<0.05.

BA, bile acid; CA, cholic acid; DCA, deoxycholic acid; MCA, mucholic acid; nd, not detectable; TCA, taurine-conjugated cholic acid; TCDCA, taurine-conjugated deoxycholic acid; TDCA, taurine-conjugated deoxycholic acid; TCDCA, taurine-conjugated chenodeoxycholic acid; TLCA, taurine-conjugated lithocholic acid; T α MCA, tauro-alpha-muricholic acid; T β MCA, tauro-beta-muricholic acid; T ω MCA, tauro-omega-muricholic acid; TUDCA, taurine-conjugated ursodeoxycholic acid.

to stimulation via TGR5.²⁶ Therefore, in the context of cholestatic liver disease, inhibiting FXR and stimulating TGR5 in the colon seems to have beneficial effects. Interestingly, colonic FXR expression remained unaffected in *Mdr2*^{-/-} mice treated with the ASBT inhibitor A4250 (data not shown). Together, these observations suggest that changes in colonic BA signalling are specific for the BA resin therapy in *Mdr2*^{-/-} mice and explained at least in part the more pronounced improvement of cholestatic liver injury in *Mdr2*^{-/-} mice in comparison to reported effects of ileal ASBT inhibitors.^{10 11}

Since BAs are potential regulators of microbiota, we performed a 16S rRNA analysis to determine microbiota composition. Although elevated concentrations of secondary BAs may implicate a shift towards the phylum Firmicutes (since they include bacteria responsible for secondary BA production²⁷), the abundance of Firmicutes remained unchanged under colestevam treatment (see online supplementary figure 5A) and the class Clostridia (which contains bacteria exhibiting 7 α -dehydroxylase activity) was even predominant in the untreated *Mdr2*^{-/-} control group (see online supplementary figure 5B). However, 7 α -dehydroxylation is restricted to taurine-deamidated BAs. Indeed, primary BAs need to be deconjugated before transformation to secondary BAs.²⁸ In line, *Mdr2*^{-/-} mice fed with colestevam have more unconjugated BAs in faeces than *Mdr2*^{-/-} control mice (*Mdr2*^{-/-} control mice 83% versus *Mdr2*^{-/-} mice fed with colestevam 91%). Interestingly, the family *Lactobacillaceae*, containing the genus *Lactobacillus* which is known to exert BSH activity²⁰ was predominant in colestevam-treated *Mdr2*^{-/-} mice (see online supporting figure 5B). Furthermore, increased BA deconjugation may also explain the elevated amount of the class δ -Proteobacteria which consists of bacteria having desulfatation

properties, since taurine contains a sulfonic moiety,²⁰ which might represent a growth factor for these bacteria. In this context, δ -Proteobacteria produce hydrogen sulfide which was shown to have anti-inflammatory properties since it is relevant for the release of trefoil factor 3, which may play a key role in mucosal regeneration and repair processes,²⁹ although this is discussed controversially.³⁰ Another reason for increased concentration of faecal secondary BAs could be a prolonged transit time of resin-bound BAs in the colon, which would provide the bacteria more time for metabolism of BAs.

Extending the use of BA sequestrants that have a long-standing and excellent safety track record may represent a cheap and readily available approach in the treatment of cholestatic liver injury beyond pruritus. Cholestyramine has been shown to improve liver enzymes in PSC patients.⁴ Furthermore, BA sequestrants are the first-line treatment for BA-induced diarrhoea,³¹ a side effect that was seen both in phase I and phase II clinical trials²² with ASBT inhibitors.²¹ While 'older resins' formulated as powders were poorly tolerated because of considerably large doses and poor taste, the newer compounds can be formulated as pills or gels which may improve patients acceptance and compliance.

In conclusion, our study demonstrates that interruption of the enterohepatic circulation of BAs with colestevam improves sclerosing cholangitis in the *Mdr2*^{-/-} mouse by altering hepatic and biliary BA composition towards a more hydrophilic configuration. Moreover increasing faecal BA excretion and modulating the faecal BA composition towards more abundant TGR5 ligands results in increased colonic levels of GLP-1. Therefore, our results suggest that reduction of ileal BA reuptake by BA sequestrants may open new

Figure 7 Colestevam treatment modulates intestinal GLP-1 secretion. (A) Representative GLP-1 IHC images (10 $\times$ magnification) reflect increased numbers of GLP-1 secreting enteroendocrine L-cells in the colon of *Mdr2*^{-/-} mice fed with colestevam. (B) GLP-1 levels in portal blood were measured by ELISA as described in material and method section. GLP-1 concentration tended to be increased in the colestevam-fed animals. (C) Colonic *Fxr* mRNA expression is significantly reduced in colestevam-treated *Mdr2*^{-/-} mice. (D) Colonic protein expression of FXR was reduced in *Mdr2*^{-/-} mice by colestevam treatment. *Significant difference from *Mdr2*^{-/-} control group; $p < 0.05$. GLP-1, glucagon-like peptide 1; mRNA, messenger RNA.

perspectives for pharmacological treatment of patients with cholangiopathies such as PSC.

Contributors CDF: study concept and design, writing of the manuscript, data collection, statistical analysis and interpretation of data. EH, VM, VK, AW, MS, NL, TS, AT: data collection, critical revision of the manuscript for important intellectual content. GP, KK, HUM: critical revision of the manuscript for important intellectual content. MT: study concept and design, interpretation of data, outlining and revising the manuscript; study oversight and acquisition of project funding.

Funding This study was funded by Austrian Science Fund. Grant number: F3517-B20.

Competing interests MT served as a consultant for Albireo, Falk, Genfit, Gilead, Intercept, MSD, Novartis and Phenex and is a member of the speakers' bureau of Falk, Gilead, MSD and Roche. He further received travel grants from Falk, Roche and Gilead and unrestricted research grants from Albireo, Falk, Gilead, Intercept, MSD and Takeda. He is also coinventor of a patent on the medical use of norUDCA. H-UM served as consultant for Albireo, AstraZeneca and Intercept and received travel grants from Falk. CDF received travel grants from Gilead, Roche, Falk, Merck, Vifor, Abbvie and Böhringer Ingelheim. EH received travel grants from Falk, Roche and MSD. She also served as a consultant for Novartis and Intercept. AW consulted for Intercept and received travel grants from Falk.

Patient consent Not required.

Provenance and peer review Not commissioned; externally peer reviewed.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

© Article author(s) (or their employer(s) unless otherwise stated in the text of the article) 2018. All rights reserved. No commercial use is permitted unless otherwise expressly granted.

REFERENCES

- Bunchorntavakul C, Reddy KR. Pruritus in chronic cholestatic liver disease. *Clin Liver Dis* 2012;16:331–46.
- Garg A, Grundy SM. Cholestyramine therapy for dyslipidemia in non-insulin-dependent diabetes mellitus. A short-term, double-blind, crossover trial. *Ann Intern Med* 1994;121:416–22.
- Kuiper EM, van Erpecum KJ, Beuers U, et al. The potent bile acid sequestrant colestevlam is not effective in cholestatic pruritus: results of a double-blind, randomized, placebo-controlled trial. *Hepatology* 2010;52:1334–40.
- Polter DE, Gruhl V, Eigenbrodt EH, et al. Beneficial effect of cholestyramine in sclerosing cholangitis. *Gastroenterology* 1980;79:326–33.
- Potthoff MJ, Potts A, He T, et al. Colesevelam suppresses hepatic glycogenolysis by TGR5-mediated induction of GLP-1 action in DIO mice. *Am J Physiol Gastrointest Liver Physiol* 2013;304:G371–G380.
- Trabelsi MS, Daoudi M, Prawitt J, et al. Farnesoid X receptor inhibits glucagon-like peptide-1 production by enteroendocrine L cells. *Nat Commun* 2015;6:7629.
- Marzoni M, Alpini G, Saccomanno S, et al. Glucagon-like peptide-1 and its receptor agonist exendin-4 modulate cholangiocyte adaptive response to cholestasis. *Gastroenterology* 2007;133:244–55.
- Marzoni M, Alpini G, Saccomanno S, et al. Exendin-4, a glucagon-like peptide 1 receptor agonist, protects cholangiocytes from apoptosis. *Gut* 2009;58:990–7.
- Sonne DP, Hansen M, Knop FK. Bile acid sequestrants in type 2 diabetes: potential effects on GLP1 secretion. *Eur J Endocrinol* 2014;171:R47–65.
- Baghdasaryan A, Fuchs CD, Österreicher CH, et al. Inhibition of intestinal bile acid absorption improves cholestatic liver and bile duct injury in a mouse model of sclerosing cholangitis. *J Hepatol* 2016;64:674–81.
- Miethe AG, Zhang W, Simmons J, et al. Pharmacological inhibition of apical sodium-dependent bile acid transporter changes bile composition and blocks progression of sclerosing cholangitis in multidrug resistance 2 knockout mice. *Hepatology* 2016;63:512–23.
- Baghdasaryan A, Claudel T, Kusters A, et al. Curcumin improves sclerosing cholangitis in Mdr2^{-/-} mice by inhibition of cholangiocyte inflammatory response and portal myofibroblast proliferation. *Gut* 2010;59:521–30.
- Fickert P, Stöger U, Fuchsbichler A, et al. A new xenobiotic-induced mouse model of sclerosing cholangitis and biliary fibrosis. *Am J Pathol* 2007;171:525–36.
- Fickert P, Wagner M, Marschall HU, et al. 24-norUrsodeoxycholic acid is superior to ursodeoxycholic acid in the treatment of sclerosing cholangitis in Mdr2 (Abcb4) knockout mice. *Gastroenterology* 2006;130:465–81.
- Tremaroli V, Karlsson F, Werling M, et al. Roux-en-Y gastric bypass and vertical banded gastroplasty induce long-term changes on the human gut microbiome contributing to fat mass regulation. *Cell Metab* 2015;22:228–38.
- Fröhlich EE, Farzi A, Mayerhofer R, et al. Cognitive impairment by antibiotic-induced gut dysbiosis: Analysis of gut microbiota-brain communication. *Brain Behav Immun* 2016;56:140–55.
- Wagner M, Fickert P, Zollner G, et al. Role of farnesoid X receptor in determining hepatic ABC transporter expression and liver injury in bile duct-ligated mice. *Gastroenterology* 2003;125:825–38.
- Trauner M, Arrese M, Soroka CJ, et al. The rat canalicular conjugate export pump (Mrp2) is down-regulated in intrahepatic and obstructive cholestasis. *Gastroenterology* 1997;113:255–64.
- Kawamata Y, Fujii R, Hosoya M, et al. A G protein-coupled receptor responsive to bile acids. *J Biol Chem* 2003;278:9435–40.
- Gérard P. Metabolism of cholesterol and bile acids by the gut microbiota. *Pathogens* 2013;3:14–24.
- Graffner H, Gillberg PG, Rikner L, et al. The ileal bile acid transporter inhibitor A4250 decreases serum bile acids by interrupting the enterohepatic circulation. *Aliment Pharmacol Ther* 2016;43:303–10.
- Hegade VS, Kendrick SF, Dobbins RL, et al. Effect of ileal bile acid transporter inhibitor GSK2330672 on pruritus in primary biliary cholangitis: a double-blind, randomised, placebo-controlled, crossover, phase 2a study. *Lancet* 2017;389:1114–23.
- Beuers U, Göke B. GLP-1 analogues: a new therapeutic approach to prevent ductopenia in cholangiopathies? *Gut* 2009;58:902–3.
- Lee YS, Jun HS. Anti-inflammatory effects of GLP-1-based therapies beyond glucose control. *Mediators Inflamm* 2016;2016:1–11.
- Fickert P, Fuchsbichler A, Wagner M, et al. Regurgitation of bile acids from leaky bile ducts causes sclerosing cholangitis in Mdr2 (Abcb4) knockout mice. *Gastroenterology* 2004;127:261–74.
- Harach T, Pols TW, Nomura M, et al. TGR5 potentiates GLP-1 secretion in response to anionic exchange resins. *Sci Rep* 2012;2:430.
- Ridlon JM, Alves JM, Hylemon PB, et al. Cirrhosis, bile acids and gut microbiota: unraveling a complex relationship. *Gut Microbes* 2013;4:382–7.
- Ridlon JM, Kang DJ, Hylemon PB. Bile salt biotransformations by human intestinal bacteria. *J Lipid Res* 2006;47:241–59.
- Albert TK, Laubinger W, Müller S, et al. Human intestinal TFF3 forms disulfide-linked heteromers with the mucus-associated FCGBP protein and is released by hydrogen sulfide. *J Proteome Res* 2010;9:3108–17.
- Linden DR. Hydrogen sulfide signaling in the gastrointestinal tract. *Antioxid Redox Signal* 2014;20:818–30.
- Camilleri M. Bile acid diarrhea: prevalence, pathogenesis, and therapy. *Gut Liver* 2015;9:332–9.