challenging epithelial and subepithelial neoplasms that are not amenable to conventional resection techniques.

Methods This was a retrospective case series study of patients underwent two methods of endoscopic full-thickness resection (EFTR), either of which was deep resection using ESD knives and post-resection closure with OTSC (Group 1), the other was pre-resection closure with OTSC and second EFTR with snare (Group 2).

Results Of all 21 patients, 11 cases were in Group 1 and 10 in Group 2. The mean time of EFTR procedure was 76.83 ± 34.97 min in Group 1 which was significantly longer than that of Group 2 ($p=0.0128$). The mean time of OTSC closure and length of hospital stay of Group 1 were also longer compared to Group 2, but the difference was not significant. Both of complete resection (R0) and technical success rate of Group 1 were 83.3% and were both 100% for Group 2.

Conclusions EFTR for pre-resection closure is potentially faster compared with the concept of applying closure after EFTR. Larger prospective controlled studies comparing those two techniques are warranted in the future.

IDDF2018-ABS-0102 RELATIONSHIP BETWEEN HISTOLOGICAL CHARACTERISTICS AND BRAF MUTATION, P53, KI67 IMMUNOEXPRESSION IN PATIENTS WITH COLORECTAL POLYP IN THAI NGUYEN

Luu Thi Binh*, Doan Anh Thang, Nguyen Trong Hieu, Tran Duc Quy, Dong Duc Hoang. *Thai Nguyen University of Medicine and Pharmacy, Vietnam*

10.1136/gutjnl-2018-IDDFabstracts.110

Background Changes in immune expression of BRAF, P53, and Ki67 proteins are part of the colorectal carcinogenic mechanisms associated with colorectal polyps.

Aims To analyse the relationship between histological characteristics and BRAF mutation, P53, Ki67 immunoexpression in patients with a colorectal polyp.

Methods A total of 81 non-cancerous colorectal polyp patients were randomised to the study. Patients undergoing endoscopic, histopathological and immunohistochemical analyses.

Results BRAF mutation in the dysplasia area showed an average of 1 ± 0.69 , non-dysplasia 0.33 ± 0.49 , $p=0.001$; P53 expression in dysplasia was found to be 1.74 ± 1.38 , non-dysplasia 1.22 ± 1.05 , $p=0.001$. Ki67 expression in dysplasia was found to be 1.33 ± 1.18 , non-dysplasia 1.20 ± 0.97 , $p=0.002$. The BRAF mutation in dysplasia area detected 50% of the traditional serrated polyps. BRAF mutation in dysplasia had 25% level ++, in pervasive samples, $p<0.001$. P53 expression in dysplasia was 68.6% level +++ in adenoma polyps group, in non-dysplasia area rate 33.3% level +++ in non-neoplastic polyp, $p<0.001$. P53 expression were mainly found in the bottom samples, the dysplasia area was 44.3% level +++, non-dysplasia 15.7% level +++, $p=0.02$. Ki67 expression in dysplasia area was 35.3% in the level of +++ adenoma polyps, non-dysplasia at 20% level +++ non neoplastic polyps, $p<0.001$. Ki67 expression of dysplasia 28.6% level +++ was mainly found in the bottom samples, $p<0.001$.

Conclusions BRAF mutation is more common in serrated polyps, P53, Ki67 expression are more common in adenomatous polyps. Immunohistochemical changes of these proteins are more common in the dysplasia area and there are differences between cell layers.

IDDF2018-ABS-0103 RELATIONSHIP BETWEEN ENDOSCOPIC CHARACTERISTICS AND BRAF MUTATION, P53, KI67 IMMUNOEXPRESSION IN PATIENTS WITH COLORECTAL POLYP IN THAI NGUYEN

Luu Thi Binh*, Doan Anh Thang, Nguyen Trong Hieu, Tran Duc Quy, Dong Duc Hoang. *Thai Nguyen University of Medicine and Pharmacy, Vietnam*

10.1136/gutjnl-2018-IDDFabstracts.111

Background Co-results of endoscopy and BRAF mutation, P53, Ki67 immunoexpression increase sensitivity and specificity in the early diagnosis of colorectal cancer.

Aims To analyse the relationship between endoscopic characteristics and BRAF mutation, P53, Ki67 immunoexpression in patients with a colorectal polyp.

Methods A total of 81 non-cancerous colorectal polyp patients were randomised to the study. Patients undergoing endoscopic, histopathological, and immunohistochemical analyses.

Results Gender ratio: male 69.1%, female 30.9%. Highest age group 41–60 was 58%. There are a total of 175 polyps, an average number per patient 2.16 ± 1.14 . The proportion of multiple polyp patients was 61.7%, single polyp 38.3%. Sessile polyp was highest: 38.3%. The size of polyps <1 cm was the highest rate, with 46.9%, >2 cm with only 7.4%. Colon polyps accounted for 67.9%, rectal 32.1%. BRAF mutation in single polyp accounted for 22.8%, multiple polyps 22%. BRAF mutation in flat-elevated polyp were lowest with 20%, pedunculated polyps had the highest rate of 26.9%. Mutation of BRAF polyp size >2 cm accounted for 50% as the highest. P53 expression was similar in the single and multiple polyp. P53 expression level +++ in pedunculated polyp with 57.7%, subpedunculated 57.1%, sessile 32.3%. In terms of size, 100% polyps >2 cm express P53, $p=0.03$. Ki67 expression was similar in the single and multiple polyp. Polyp size >2 cm accounted for the highest Ki67 expression rate of 66.6%. Pedunculated polyp had the highest rate of Ki67 expression in the dysplasia of level +++ with 42.3%, subpedunculated and sessile polyp 16.1%, $p=0.04$.

Conclusions P53 immunoexpression are usually manifested in large pedunculated and subpedunculated polyps. Ki67 immunoexpression in the dysplasia area, manifested in pedunculated and subpedunculated polyps.

IDDF2018-ABS-0115 THE ASSOCIATION BETWEEN ELECTRONIC CIGARETTE SMOKING AND GASTROESOPHAGEAL REFLUX DISEASE IN ADULT URBAN POPULATION

¹Ahmad Fariz Malvi Zamzam Zein*, ²Catur Setiya Sulistiyana, ³Murdani Abdullah. ¹Department of Internal Medicine, Faculty of Medicine, Swadaya Gunung Jati University, Indonesia; ²Department of Medical Education, Faculty of Medicine, Swadaya Gunung Jati University, Indonesia; ³Division of Gastroenterology, Department of Internal Medicine, Cipto Mangunkusumo Hospital, Indonesia

10.1136/gutjnl-2018-IDDFabstracts.112

Background This study was aimed to investigate the association between electronic cigarette (e-cig) smoking and gastroesophageal reflux disease (GERD) among the adult urban population.

Methods A cross-sectional study was conducted among 267 adult people in Cirebon City, West Java, Indonesia. A self-administered questionnaire based on earlier validated GERD