OPEN ACCESS

ORIGINAL ARTICLE

Identification of TMAO-producer phenotype and host–diet–gut dysbiosis by carnitine challenge test in human and germ-free mice

Wei-Kai Wu,^{1,2} Chieh-Chang Chen,³ Po-Yu Liu,^{4,5} Suraphan Panyod,^{2,6} Ben-Yang Liao,⁷ Pei-Chen Chen,² Hsien-Li Kao,³ Han-Chun Kuo,^{8,9} Ching-Hua Kuo,^{8,9} Tina H T Chiu,¹⁰ Rou-An Chen,² Hsiao-Li Chuang,¹¹ Yen-Te Huang,¹¹ Hsin-Bai Zou,¹² Cheng-Chih Hsu,¹² Ting-Yan Chang,⁷ Chin-Lon Lin,¹³ Chi-Tang Ho,¹⁴ Hon-Tsen Yu,^{4,5} Lee-Yan Sheen,² Ming-Shiang Wu^{3,6}

► Additional material is published online only. To view please visit the journal online (<http://dx.doi.org/10.1136/gutjnl-2018-317155>).

For numbered affiliations see end of article.

Correspondence to

Professor Ming-Shiang Wu, Department of Internal Medicine, National Taiwan University Hospital, Taipei 100, Taiwan; mingshiang@ntu.edu.tw and Professor Lee-Yan Sheen, Institute of Food Science and Technology, National Taiwan University, Taipei 106, Taiwan; lysheen@ntu.edu.tw

Received 9 July 2018

Revised 8 October 2018

Accepted 11 October 2018

Published Online First

30 October 2018

Watch Video
gut.bmj.com

Check for updates

© Author(s) (or their employer(s)) 2019. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Wu W-K, Chen C-C, Liu P-Y, et al. *Gut* 2019;**68**:1439–1449.

ABSTRACT

Objective The gut microbiota-derived metabolite, trimethylamine N-oxide (TMAO) plays an important role in cardiovascular disease (CVD). The fasting plasma TMAO was shown as a prognostic indicator of CVD incident in patients and raised the interest of intervention targeting gut microbiota. Here we develop a clinically applicable method called oral carnitine challenge test (OCCT) for TMAO-related therapeutic drug efforts assessment and personalising dietary guidance.

Design A pharmacokinetic study was performed to verify the design of OCCT protocol. The OCCT was conducted in 23 vegetarians and 34 omnivores to validate gut microbiota TMAO production capacity. The OCCT survey was integrated with gut microbiome, host genotypes, dietary records and serum biochemistry. A humanised gnotobiotic mice study was performed for translational validation.

Results The OCCT showed better efficacy than fasting plasma TMAO to identify TMAO producer phenotype. The omnivores exhibited a 10-fold higher OR to be high TMAO producer than vegetarians. The TMAO-associated taxa found by OCCT in this study were consistent with previous animal studies. The TMAO producer phenotypes were also reproduced in humanised gnotobiotic mice model. Besides, we found the faecal *CntA* gene was not associated with TMAO production; therefore, other key relevant microbial genes might be involved. Finally, we demonstrated the urine TMAO exhibited a strong positive correlation with plasma TMAO ($r=0.92$, $p<0.0001$) and improved the feasibility of OCCT.

Conclusion The OCCT can be used to identify TMAO-producer phenotype of gut microbiota and may serve as a personal guidance in CVD prevention and treatment.

Trial registration number NCT02838732; Results.

INTRODUCTION

The gut microbiota plays a critical role in the development of cardiovascular disease (CVD), and studies have shown that some micro-organism-derived metabolites produced in the intestine are involved in CVD pathogenesis.^{1–4} Intestinal microbial metabolites, such as trimethylamine N-oxide (TMAO), short-chain fatty

Significance of this study**What is already known on this subject?**

- The gut microbiota dependent metabolite, trimethylamine N-oxide (TMAO) was proved to contribute to the progression of atherosclerosis.
- Although the fasting plasma TMAO was shown as a prognostic indicator in patients with cardiovascular disease (CVD), it might underestimate the TMAO producing function of gut microbiota because of host's efficient renal clearance of plasma TMAO.
- Omnivores and vegetarians have different functions of gut microbiota, but the function of TMAO production capacity has not been addressed clearly.
- The microbial Rieske-type *cntA/B* gene is reported to be in part responsible for aerobic carnitine metabolism to trimethylamine from in silico and in vitro studies but has not been evaluated in an in vivo human study.

What are the new findings?

- This is the first study designed for the clinical assessment of individual TMAO production capacity using an oral carnitine challenge test (OCCT) with verified challenge dose and sampling times.
- The OCCT exhibits better efficacy than fasting plasma TMAO in identifying TMAO-producer phenotype of human gut microbiota.
- Omnivores have higher risk to be TMAO producer than vegetarians, but some long-term vegetarians can still demonstrate substantial TMAO production from oral carnitine by their gut microbiota.
- Faecal *CntA* may not be a useful biomarker in identifying TMAO producer phenotype according to our in vivo human study. There might be other potential microbial genes responsible for carnitine metabolism to TMAO in the faeces.

Significance of this study

How might it impact on clinical practice in the foreseeable future?

- The OCCT reveals the TMAO-producer phenotypes and may serve as a personalised dietary guidance for patients with CVD.
- The OCCT use in human study may serve as a benchmark for assessing therapeutic drug development efforts for the TMAO pathway.
- The OCCT can simulate a postprandial TMAO and provide a benchmark for future investigation of diet-induced enhanced thrombotic potential.

acids and secondary bile acids, act as mediators in CVD-related chronic disorders such as atherosclerosis, obesity and type 2 diabetes.^{5 6} Among these gut microbiota-derived metabolites, TMAO is mainly produced from the metabolism of dietary carnitine and choline by gut bacteria. Studies have identified TMAO as an essential proatherogenic and prothrombotic compound that drives the pathogenesis of CVD.^{7–9}

L-carnitine is an abundant nutrient in meat products (especially in red meat) and functions by carrying fatty acids across the mitochondrial membrane for beta-oxidation.¹⁰ However, carnitine may be metabolised by gut microbiota and thereby increase the risk of cardiovascular events in patients with coronary artery disease.¹¹ In the intestine, gut microbiota convert unabsorbed carnitine into trimethylamine (TMA), which is subsequently oxidised to TMAO by flavin monooxygenase (FMO3) in the liver.¹ TMAO then enters circulation and is efficiently excreted by the kidneys.¹² Prospective epidemiology studies have suggested a positive correlation between increased plasma TMAO levels and CVD progression.^{12 13 14} In a cohort (n=2595) of patients with coronary artery disease, high carnitine levels were associated with incident risks of myocardial infarction, stroke and death within 3 years, but only significantly in patients with concomitantly high TMAO levels.¹¹ These findings may serve as a reasonable explanation for the relationship between red meat consumption and increased CVD risk.¹⁵ In animal studies, chronic dietary carnitine exposure increased the plasma level of TMAO, the trimethylamine (TMA) synthesis capacity of gut microbiota and the progression of atherosclerosis.^{11 16} The mechanism underlying the contribution of TMAO to atherosclerosis involves increased foam cell formation, decreased reverse cholesterol transport, and enhanced platelet aggregation.^{1 11 17} The microbial effects of TMAO production can be reduced by broad-spectrum antibiotics but the issue of antibiotic resistance remains to be challenging.² A recently discovered choline analogue named 3,3-dimethyl-1-butanol has demonstrated promising efficacy to inhibit microbial conversion of choline/carnitine into TMA.^{18 19} Other natural bioactive compounds such as resveratrol and allicin were also reported in animal study to reduce TMAO production.^{20 21}

Recently, the fasting plasma TMAO was also proved as a prognostic marker for both short-term and long-term incident cardiovascular events among subjects with acute coronary syndrome or stroke.^{14 22} Nevertheless, there remained no clear cut-off value of plasma TMAO for unfavourable outcomes. In a dose-response meta-analysis of clinical studies, the risk of

Table 1 Characteristics of study participants

	Vegetarian (n=23)	Omnivore (n=34)	P values
Female (n (%))	14 (65)	24 (71)	
Age (years)	34.13±1.70	30.18±1.30	0.0665
BMI (kg/m ²)	22.40±0.55	21.81±0.57	0.4793
Genotype FMO3-SNP (n (%))			
Glu158Lys – AAG	6 (26.1)	8 (23.5)	
Glu308Gly – GGG	6 (26.1)	8 (23.5)	
Plasma			
Glucose-AC (mg/dL)	69.39±2.20	75.41±1.91	0.0459
AST (U/L)	11.30±1.036	15.29±1.19	0.0211
ALT (U/L)	7.22±0.94	12.91±2.38	0.0627
BUN (mg/dL)	9.57±0.65	11.63±0.50	0.0145
Creatinine (mg/dL)	0.55±0.03	0.60±0.02	0.2041
T-cholesterol (mg/dL)	140.50±4.27	174.90±5.23	<0.0001
Triglyceride (mg/dL)	89.13±11.31	88.32±10.07	0.9584
LDL-C (mg/dL)	75.13±4.25	97.44±4.64	0.0014
hsCRP (mg/dL)	0.06±0.01	0.18±0.09	0.2766
TMAO (μM)	1.87±0.21	3.54±0.96	0.1618
Carnitine (μM)	34.72±3.07	63.46±5.08	<0.0001
Urine			
TMAO (nmol/mmol Cr)	33.98±3.49	68.13±20.70	0.1843
Carnitine (nmol/mmol Cr)	2.52±0.64	15.37±6.35	0.1035

*The p value is from the comparison between vegetarian and omnivore volunteers using Student's t-test. Values are the mean±SEM.

AST, aspartate aminotransferase; ALT, alanine aminotransferase; BMI, body mass index; BUN, blood urea nitrogen; Cr, creatinine; FMO3, flavin monooxygenase; hsCRP, high-sensitivity C reactive protein; LDL-C, low-density lipoprotein cholesterol; TMAO, trimethylamine N-oxide.

all-cause mortality increased by 7.6% per each 10 μM increment of plasma TMAO.²³ In recent animal and human platelet aggregation studies, the concentration of plasma TMAO greater than 10–30 μM significantly enhance the thrombosis potential.^{17 19 24} However, the fasting plasma TMAO level in subjects with normal renal function seldomly exceed 10 μM for both healthy and diseased populations.^{2 25} Accordingly, a postprandial plasma TMAO might theoretically reflect the pathophysiological level of TMAO. Thus, we established an oral carnitine challenge test (OCCT) to simulate the postprandial plasma TMAO as well as functionally measure the TMAO synthesis capacity from host-diet-microbiota interactions.

RESULTS

Omnivores exhibited remarkably higher levels of plasma carnitine and cholesterol than did vegetarians

A comparison of demographic data and biochemical values revealed significant differences in several metabolic parameters, namely, plasma carnitine, serum total cholesterol, low-density lipoprotein, urea nitrogen, fasting glucose and aspartate transaminase levels, between the omnivores and vegetarians (table 1). Of the compared parameters, carnitine and cholesterol levels between the omnivores and vegetarians exhibited the most significant differences (p<0.0001). However, the urine carnitine level between the omnivores and vegetarians did not exhibit a significant difference. In general, the urine TMAO level was remarkably higher than the plasma TMAO level, whereas carnitine was more abundant in plasma samples. This finding may indicate that the human body tends to preserve carnitine as a nutrient and excrete TMAO as waste through the urinary system.^{12 26}

Figure 1 The divergence of dietary patterns between vegetarian and omnivore contributed no significant differences for gut microbiome composition and diversity. (A) Heatmap of dietary micronutrients in omnivores versus vegetarians (q value < 0.1) with clustering nutrients colour labelled by six nutrient categories. Red=higher abundance, blue=lower abundance. (B) The carnitine and cholesterol consumption levels of vegetarians and omnivores exhibited highly significant differences. (C) The principle component analysis of FFQ nutrients data indicated significantly divergent patterns between omnivores and vegetarians (permutational multivariate analysis of variance (PERMANOVA): $p < 0.001$). (D) Compositional profiling of gut microbiota in vegetarians and omnivores revealed no significant difference (PERMANOVA: $p = 0.3528$) demonstrated by principle coordinate analysis calculated using Bray-Curtis distance. (E) Comparison of alpha diversity index in vegetarians versus omnivores. (F) The Firmicutes/Bacteroidetes ratio in omnivores versus vegetarians exhibited no significant difference. Data in all bar plots are expressed as mean $\pm$ SEM. All statistics in bar plots and box plots were analysed by Student's t -test. FFQ, food frequency questionnaire.

Fasting TMAO levels in blood and urine were both higher in the omnivores (blood: $3.54 \pm 0.96 \mu\text{M}$; urine: $68.13 \pm 20.70 \text{ nmol/mmol}$) than in the vegetarians (blood: $1.87 \pm 0.21 \mu\text{M}$; urine: $33.98 \pm 3.49 \text{ nmol/mmol}$); however, the differences did not reach statistical significance (table 1).

Markedly different dietary patterns between omnivores and vegetarians contributed to minor difference in gut microbiome

Among 87 nutrient levels calculated using the food frequency questionnaire (FFQ), 43 differed significantly between the omnivores and vegetarians ($p < 0.05$) with FDR < 0.1 (online supplementary table S1). The omnivores consumed more cholesterol, carnitine, saturated fat and animal protein than did the vegetarians (figure 1A,B). Thus, the omnivores exhibited higher serum cholesterol, carnitine, LDL and urea nitrogen levels (table 1). The results of the principle component analysis of all the nutrients calculated using the FFQ also showed distinct dietary patterns between the omnivores and vegetarians (figure 1C). However, gut microbiome profiles and indicators between the omnivores and vegetarians did not considerably differ according to the results of the principle coordinate analysis, alpha diversity and Firmicutes/Bacteroidetes (F/B) ratio (figure 1D-F), although moderately significant differences in minor gut microbiome communities ($< 2\%$ species) were identified (online supplementary figure S1A). The results of the linear discriminant analysis effect size (LEfSe) analysis revealed that Prevotellaceae was prevalent in the vegetarians' gut microbiome, whereas the

gut microbiome of the omnivores featured characteristic taxa of Clostridiaceae, Bacteroidales S24 and Eubacterium (online supplementary figure S1B). These findings are consistent with those of previous studies.^{27 28}

Plasma TMAO peaked in the interval between 24 hours and 48 hours of the OCCT

We assumed that TMAO formation in blood occurred at approximately 24 hours of the OCCT based on the time required for bowel transit, microbial catabolism and the enzymatic reaction of hepatic FMO3 to convert the ingested carnitine into plasma TMAO. We performed a pharmacokinetic study of the OCCT to confirm this hypothesis (figure 2). In the study, 13 participants received seven blood tests at indicated time points after ingestion of three tablets of L-carnitine fumarate (General Nutrition Centers (GNC), Pennsylvania, USA). We verified the carnitine amount of each tablet through high-performance liquid chromatography (HPLC) to ensure that the dose of L-carnitine in each test was consistent. After the OCCT, plasma TMAO levels remained stable at trough levels for 12 hours and peaks started to present after 24 hours (figure 2A). The plasma carnitine level peaked at 4 hours after OCCT while TMA and γ -butyrobetaine in the plasma remained at very low concentration (online supplementary figure S2A,B). Among the 13 participants, 6 participated in an additional pharmacokinetic study 3 months later, and the original trends and values for each individual were reproduced in the results (figure 2B,C). Thus, the OCCT was simplified

Figure 2 Pharmacokinetic study of oral carnitine challenge test (OCCT). (A) Thirteen volunteers were recruited for a pharmacokinetic (PK) study of the OCCT. Each participant received three tablets of GNC L-carnitine (approximately 1200 mg L-carnitine) and blood drawings at 4th hour, 8th hour, 12th hour, 24th hour, 36th hour and 48th hour. (B) The bar plots of AUC in OCCT for different volunteers and the same volunteer with different PK studies (six volunteers received the second PK study 3 months later). (C) Normalised dissimilarity of AUC of different pharmacokinetic studies in the same and different individuals (defined as $|AUC1 - AUC2| / [AUC1 + AUC2]$) AUC1: AUCs of 1st PK study; AUC2: AUCs of second PK study. These data suggested the trend of TMAO production capacity is reproducible in the same individual periodically. (D) Validation and simplification of sample collection time points for the OCCT according to the results of pharmacokinetic studies. AUC, area under the curve; CCT, carnitine challenge test; TMAO, trimethylamine N-oxide.

to involve three sample collection times: the baseline (fasting plasma TMAO), 24 hours and 48 hours (figure 2D).

OCCT effectively differentiated gut microbiota-mediated TMAO production capacity between the omnivore and vegetarian groups

Differences in fasting plasma TMAO levels between the omnivores and vegetarians were moderate. Therefore, the proposed OCCT was developed to determine individual TMAO production capacity as mediated by gut microbiota. All the omnivore and vegetarian volunteers participated in a simplified OCCT with blood and urine collection at baseline, 24 hours and 48 hours. In the OCCT curve, plasma TMAO levels in the omnivores versus vegetarians diverged at 24 hours and 48 hours, but fasting TMAO levels remained similar (figure 3A,B). In contrast to the values of fasting TMAO, those of the area under the curve (AUC), and the maximum values of the OCCT were significantly higher in the omnivores than in the vegetarians (figure 3C).

Omnivores were more likely to be high TMAO producers than the vegetarians

To define the functional phenotype of gut microbiota for TMAO production, we plotted the AUC values from the OCCT results

for all the 57 participants and grouped the values into quartiles. The 14 participants with Q1-level AUC values were labelled as low TMAO producers, and the 14 participants in Q4 were labelled as high TMAO producers (figure 3D). The participants with levels in between Q1 and Q4 were labelled as intermediate producers. Among the omnivore participants, 35.3% were grouped as high TMAO producers and 14.7% were grouped as low TMAO producers. By contrast, only 8.7% of the vegetarians were grouped as high TMAO producers and 39.1% were grouped as low TMAO producers (figure 3E). With regard to TMAO production phenotypes defined by the OCCT, the participants in the omnivore group exhibited a 10-fold OR (OR: 10.8, 95% CI 1.69 to 68.94) of being a high TMAO producer compared with the vegetarian group (figure 3F). However, despite long-term adherence to a strict vegetarian diet (>10 years), some vegetarians exhibited considerable TMAO production when high doses of carnitine were consumed.

Gut microbiome profiles become distinctive after functional phenotyping of TMAO production capacity

In this study, the functional phenotypes of the high TMAO producers (n=14) and low TMAO producers (n=14) identified using the OCCT corresponded with distinctive OCCT

Figure 3 Omnivores and vegetarians exhibited different levels of ability to transform L-carnitine into TMAO in the body. (A) An oral carnitine challenge test (OCCT) was administered to 23 vegetarians and 34 omnivores, and plasma TMAO levels were measured at indicated times after the OCCT. The differences in plasma TMAO levels between the vegetarians and omnivores appeared at 24 hours and 48 hours compared with baseline. Data are expressed as mean $\pm$ SEM; * $p < 0.05$. (B) No significant difference of fasting plasma TMAO levels was noted between the vegetarian and omnivore groups. (C) The AUC values and maximum values of the carnitine challenge test for the omnivores were both significantly higher than for the vegetarians. (D) The population was grouped into four quartiles according to the AUC values of the OCCT. The Q4 population was defined as high TMAO producers, the Q1 as low producers and Q2–Q3 as intermediate producers. (E) A percentage of 35.3 of the omnivores were grouped as high TMAO producers compared with 8.7% of the vegetarians. A percentage of 14.7 of the omnivores were grouped as low producers compared with 39.1% of the vegetarians. (F) Among the high producers, 12/14 (86%) were omnivores, whereas among the low producers, 5/14 (36%) were omnivores. The ORs of omnivores versus vegetarians as being high TMAO producers is 10.8 (95% CI 1.69 to 68.94). Data in all bar plots are expressed as mean $\pm$ SEM. Plasma TMAO levels at indicated times in OCCT and plasma fasting TMAO data were analysed by Student's *t*-test. AUC of TMAO and TMAO_{max} in OCCT were analysed by Mann-Whitney U test. AUC, area under the curve; TMAO, trimethylamine N-oxide.

curves and significantly different AUC and TMAO_{max} values (figure 4A–C). However, the difference in fasting plasma TMAO levels between the high and low TMAO producers remained moderate (figure 4B). This finding further suggested that a challenge test, rather than a single blood test, should be considered in clinical assessments to determine the functional phenotype of TMAO-producing gut microbiota. In addition, the compositional patterns of gut microbiome in the high versus low TMAO producers were distinctive, as demonstrated by the heatmap and principle coordinate analysis (figure 4D,G). Other common gut microbiome functional indicators, including F/B ratio, Shannon index and Chao1 index, also indicated significant differences between the high and low TMAO producers (figure 4E,F). The higher alpha diversity of the high TMAO producers suggested higher compositional complexity of the gut microbiome for acquiring TMA synthesis functions. The higher F/B ratio in the high TMAO producers corresponded to the findings of high-risk patients with CVD in previous studies.^{9 29} Furthermore, the results of the LEfSe analysis regarding the characteristic bacterial taxa between the high and low TMAO producers were similar to the results of previous mouse experiments (figure 4H).^{11 16}

TMAO-producing phenotypes determined using the OCCT were reproduced in the humanised gnotobiotic mice model

The TMAO-producing functional phenotypes determined using the OCCT were based on complex diet–gut microbiota–host interactions. Therefore, we used a humanised gnotobiotic mouse model to reconstruct these complex interactions *in vivo*. Fecal microbiota transplantation (FMT) was performed on germ-free mice by using faeces from a high TMAO producer and a low TMAO producer from our study cohort. In the animal model, diet and host factors were well controlled. We divided eight 9-week-old male germ-free mice into two groups (n=4 in each group) and performed FMT with gastric gavage in the mice of each group using faecal samples from two human donors (a high TMAO producer and a low TMAO producer). All the mice received a carnitine-supplemented diet for 5 weeks after which they underwent d9-carnitine oral challenge tests. The results demonstrated that the transplantation of gut microbiota transmitted the human host's TMAO-producing phenotypes to the mouse hosts (figure 4I).

Previously found microbial enzyme system *cntA/B* gene may be an insufficient marker for *in vivo* human TMAO production

For the process of TMAO production in the human body, the key step is the conversion of carnitine into TMA from gut

Figure 5 Carnitine intake amount, faecal *CntA* abundances and host *FMO3* genotypes correlated unfavourably with TMAO-producing ability in the human body. (A) The *CntA* abundances (% of community) exhibited a poor correlation to fasting plasma TMAO, (B and C) AUC and maximum values of plasma TMAO of OCCT. (D) Carnitine intake amount also poorly correlated with the AUC of plasma TMAO. (E and F) Two common host flavin mono-oxygenase (*FMO*) single-nucleotide polymorphism, *FMO3*/Lys158 and *FMO3*/Gly308 mutant alleles, were detected in the study population, and no difference of TMAO production was exhibited between mutant and wild genotypes. Data in bar plots are expressed as mean±SEM. Pearson's correlation was used to calculate association between two variables. Student's t-test was used for two-group comparison. AUC, area under the curve; OCCT, oral carnitine challenge test; TMAO, trimethylamine N-oxide.

microbiota metabolism in the intestine. It was reported in an in vitro experiment that a Rieske-type oxygenase/reductase microbial enzymatic system gene, *cntA/B*, was capable of converting carnitine into TMA.³⁰ In another study, a degenerate primer was designed in attempt to measure TMA-producing capacity by quantifying *cntA* in the faecal microbiome community.³¹ We used this published method to measure the abundance of *cntA* in faecal DNA samples and compare the faecal *cntA* abundances to the TMAO values obtained from the same participants. Results indicated that faecal *cntA* amounts correlated poorly with numerous TMAO-relevant values determined using the OCCT (figure 5A–C). We further evaluated the correlation between the relative abundances of previously reported *cntA*-containing bacterial taxa (ie, *Escherichia coli*, *Klebsiella* spp and *Citrobacter* spp)^{30,31} and TMAO values, but no significantly positive correlation was revealed (online supplementary figure S3A–C). In addition, considering the difference in TMAO-producing ability between the omnivores and vegetarians, we compared the calculated carnitine intake amounts with the TMAO AUC values, and the correlation remained non-significant (figure 5D). Furthermore, we assessed the effect of host genetic factors on the TMAO concentration; two commonly occurring single-nucleotide polymorphisms of *FMO3* (Glu158Lys and Glu308Glu) were examined among the participants in this study,^{32,33} and no significant differences in TMAO values between the mutant and wild-type participants were identified (figure 5E,F). Finally, we

compared the demographic, biochemical and dietary data of the high and low TMAO producers and found no significantly different markers (table 2 and online supplementary figure S4). Collectively, these findings suggested that current microbial and host genetic biomarkers, dietary data and other features may not reflect the complex metabolic pathway of the interaction among diet, microbial community and the host for TMAO formation.

Urine sampling provides an alternative and convenient method for the OCCT

While the microbial enzyme system *cntA* genetic marker failed to correlate TMAO formation, the OCCT remains the standard for measuring TMAO production capacity in human body. However, plasma TMAO levels usually peak after 24 hours of the OCCT, and the required serial blood sampling may present challenges for clinical practice and patient compliance. Thus, we compared 171 paired blood and urine samples collected at the same times. The samples exhibited a strong significant correlation of TMAO concentrations. Moreover, the AUC values of 57 paired blood and urine samples from the participants who underwent the OCCT were compared and exhibited a highly significant correlation (figure 6A). These results suggested that urine collection may provide a substitute for blood collection in the OCCT, which may increase the feasibility of this assessment method for clinical practice.

Table 2 Characteristics of high and low TMAO producers

	Low producer (n=14)	High producer (n=14)	P values
Female (n (%))	9 (64)	11 (78)	
Age (years)	30.86±7.84	32.64±8.33	0.5644
BMI (kg/m ²)	22.49±3.58	21.02±3.25	0.2653
Genotype FMO3-SNP (n (%))			
Glu158Lys – AAG	2 (14.3)	4 (28.6)	
Glu308Gly – GGG	2 (14.3)	2 (14.3)	
Plasma			
Glucose-AC (mg/dL)	71.14±9.54	71.21±10.05	0.9848
AST (U/L)	12.64±5.40	14.50±7.71	0.4672
ALT (U/L)	8.93±6.89	14.43±20.14	0.3426
BUN (mg/dL)	10.32±3.76	11.22±1.70	0.4223
Creatinine (mg/dL)	0.55±0.17	0.57±0.09	0.6746
T-cholesterol (mg/dL)	159.07±29.13	169.43±30.15	0.3638
Triglyceride (mg/dL)	73.43±39.96	99.93±84.73	0.2996
LDL-C (mg/dL)	92.93±27.84	91.64±14.43	0.8792
hsCRP (mg/dL)	0.05±0.04	0.08±0.09	0.2924
TMAO (μM)	1.49±0.54	5.62±8.27	0.0737
Carnitine (μM)	46.53±29.31	58.64±28.42	0.2774
Urine			
TMAO (nmol/mmol Cr)	30.42±15.62	125.02±175.14	0.0546
Carnitine (nmol/mmol Cr)	6.02±8.32	15.26±46.00	0.4665

*The p value is from the comparison between the high and low producers using Student's t-test. Values are the mean±SEM.

ALT, alanine aminotransferase; AST, aspartate aminotransferase; BMI, body mass index; BUN, blood urea nitrogen; Cr, creatinine; FMO3, flavin monooxygenase; hsCRP, high-sensitivity C-reactive protein; LDL-C, low-density lipoprotein cholesterol; TMAO, trimethylamine N-oxide.

DISCUSSION

We developed a method to determine the functional status of the human gut microbiota, with regard to its TMAO production capacity. The OCCT in this study successfully distinguish between the omnivores and vegetarians according to their TMAO-producing ability. Using this approach, we characterise the gut microbiome of high and low TMAO producers as well as reproduce the TMAO producing phenotypes in germ-free mice by faecal microbiota transplantation. We also demonstrated the urine TMAO as an alternative biomarker for plasma TMAO that may facilitate the transition to clinical practice.

The OCCT may be used to identify individual TMAO-producing capacity from gut microbiota and thereby serve as a guidance of intervention to reduce TMAO production in the human body. For example, a low-carnitine diet may be suggested to high TMAO producers as a dietary instruction for patients with CVD. Antibiotics might be an option to reduce TMAO production by inhibiting TMAO-relevant gut bacteria but may be discouraged due to potential side effects and treatment failure caused by antibiotic resistance. Moreover, the OCCT can simulate a postprandial TMAO and reveal pathophysiological levels of plasma TMAO in subjects. In a recent pioneer study conducted by Zhu *et al.*,²⁴ 18 healthy volunteers receiving continuous choline supplementation for 1 month significantly increased the fasting plasma TMAO level as well as enhanced platelet hyper-responsiveness. Therefore, the OCCT may also be used in revealing the diet-induced TMAO and assessing the correlated thrombosis potential. Finally, since the OCCT could indicate the TMAO producer phenotype, it might serve as a benchmark for further TMAO-relevant microbial biomarker investigation and validation.

In previous studies, the fasting plasma TMAO level was proved to be a prognostic indicator of major adverse cardiac

event for patients with CVD.^{2 11} From literature review, there seems a significant overlap of median/IQR of fasting plasma TMAO between patients with CVD and healthy subjects.^{2 25} The fasting plasma TMAO levels are usually low and exhibit rightward-skewed distributions among both healthy and diseased populations.^{2 14 25} The phenomenon may be explained by the fluctuated feature of plasma TMAO caused by efficient urinary excretion and dietary influence.^{12 26} It also suggests the fasting plasma TMAO may not able to reveal higher and potential pathological value of a postprandial plasma TMAO. In recent studies by Zhu *et al.*, the concentration of plasma TMAO would be at least 10–30 μM to enhance thrombosis potential and arterial occlusion.^{17 19 24} Therefore, the TMAO values acquired from OCCT might be helpful in revealing the pathophysiological levels of plasma TMAO from regular diet–microbiota interactions.

Recently, Koeth *et al.* recruited five omnivores and five vegetarians to receive a carnitine challenge test of 250 mg of isotope-labelled d3-methyl-carnitine, and the vegetarians exhibited almost no TMAO production as compared with the omnivores.¹¹ However, we noticed that some vegetarians and even vegans were still capable of producing TMAO from their gut microbiota when administrating carnitine. Some of the TMAO producer vegetarians are lacto-ovo-vegetarians and may be exposed to small amount of carnitine from their diet, but the dietary carnitine intake among vegetarians calculated by FFQ is not correlated to the TMAO production in OCCT. We assumed that the bacteria capable of carnitine metabolism may still be able to thrive by nutrients other than carnitine in the gut, although the carnitine intake may provide additional carbon source to these bacteria and promote the growth. However, this assumption remains to be investigated by further studies. A previous animal study conducted showed that γ-butyrobetaine is a proatherosclerotic intermediate in gut microbial metabolism of carnitine to TMAO.¹⁶ Nevertheless, the levels of γ-butyrobetaine in our human plasma samples are quite low and had no obvious peak values. We speculated the bacteria responsible for converting carnitine to γ-butyrobetaine may not be functionally predominant in the human gut, but this should be confirmed in future study. Furthermore, although previous research had already used isotope-labelled carnitine for a carnitine challenge test,¹¹ the relatively low challenge dose of isotope-labelled carnitine may not reflect the pathophysiological status of diet–microbiota interactions. The accessibility and representativeness of an isotope-labelled compound for a challenge test may also be a barrier for clinical practice. Collectively, the OCCT in this study using regular carnitine with designed dosage and sampling times may have advantages to step forward into clinical setting.

A novel microbial Rieske-type oxygenase/reductase gene, the *cntA/B*, was found being able to convert carnitine into TMA.³⁰ The gene was reported to exist among only few species including *E. coli*, *Klebsiella* spp and *Citrobacter* spp and so on.^{30 31} However, both *cntA* gene quantification and related bacterial species abundance in our faecal samples showed no significant correlation to the TMAO values from OCCT. Therefore, there might exist other key player bacteria and genes contributing to transformation of carnitine into TMA in the colon (an obligatory anaerobic environment) since the *cntA/B* is an oxygen-dependent enzymatic reaction system. However, with regard to the error rate by using a degenerate primer for qPCR and the relatively low taxonomic resolution of 16S amplicon sequencing, further metagenome wide study may be required to address the issue more clearly.

This study has some limitations. First, although choline and carnitine are both materials from which gut microbiota produce

Figure 6 Urine TMAO levels exhibited strong correlation with plasma TMAO levels and may serve a substitute specimen for an oral carnitine challenge test (OCCT). (A) The logarithmic TMAO values of 171 plasma samples strongly correlated with the corresponding TMAO values of urine samples from the same subject and sampling times. The logarithmic AUC values of plasma TMAO in 57 participants strongly correlated with the AUC values of urine TMAO in the same OCCT. (B) The OCCT may reflect the results from crosstalk between diet, gut microbiota and the host and could be used to measure TMAO-producing ability individually. The OCCT may serve as a personalised dietary guidance or a diet-induced thrombotic risk surveillance. It could also be used to assess therapeutic efficacy of new drug development or may be served as a benchmark for investigation of TMAO-relevant biomarkers in the faeces. Pearson's correlation was used to calculate association between two variables. AUC, area under the curve; TMAO, trimethylamine N-oxide.

TMA, we only used carnitine as the challenge material in this study. We chose to develop the OCCT because, in normal human population, we have natural cohort of high carnitine and low carnitine consuming groups (ie, omnivores and vegetarians); thus, we can use this test to investigate if people consuming higher carnitine from their diet have higher capability of converting carnitine into TMA through their gut microbiota.

In contrast, it is difficult to find a cohort with subjects divided into high and low choline diets since the choline is widely distributed in many food categories, including eggs, soybeans, meat, fish, some fruits and vegetables.¹⁹ Therefore, our study could only successfully prove that people with higher carnitine consumption in the diet generally had higher TMAO production capacity from gut microbiota carnitine metabolism. Besides, the carnitine metabolising bacteria (eg, *cntA/B* containing bacteria such as *E. coli*, *Klebsiella* spp, *Citrobacter* spp and so on) are different from that responsible for choline utilisation (eg, *cutC/D* containing *Desulfovibrio alaskensis*, *Proteus mirabilis* and so on).^{30 31 34} Therefore, the capacity of TMAO production from carnitine and choline metabolism by gut flora may be more appropriate to discuss separately. Second, in our study, we noticed that the peak of TMAO in plasma occurred after 48 hours of the OCCT in some cases and that could skew the AUC of the OCCT curves. In addition, only two samples were

taken between 12 hours and 48 hours after OCCT and the low resolution of sampling in the 12–48 span may lead to a missed peak value in OCCT. Adding a sample timing of 72 hours would be helpful but may still be less than optimal.

In summary, we presented a clinically oriented OCCT that can be used to determine the proatherosclerotic TMAO-producing capacity of complex diet–host–microbial interactions. To the best of our knowledge, the proposed methods represent the most comprehensive design thus far for the assessment of TMAO-producing capacity of gut microbiota in a human study. In the era of precision medicine, the OCCT may serve as a personalised dietary guidance for patients with CVD, an assessment tool for the therapeutic efforts of new drug development, a standard method for diet-induced thrombosis risk survey as well as a potential benchmark to investigate TMAO-relevant biomarkers in the faeces (figure 6B).

MATERIALS AND METHODS

Omnivore and vegetarian study populations

Flyers advertising the study were distributed, and 57 healthy volunteers (23 vegetarians and vegans and 34 omnivores) were recruited. In this study, participants who self-reported not having eaten any meat or seafood products for prior 2 years or longer

were defined as vegetarians. All participants were screened using a health history questionnaire. Inclusion criteria were as follows: (1) age ≥ 20 years and (2) no exposure to antibiotics, probiotics or carnitine supplements within the previous month. Participants were excluded from the study if they reported recent gastrointestinal discomfort (such as abdominal pain or diarrhoea) or a history of chronic diseases including myasthenia gravis, diabetes mellitus, chronic renal disease, hyperparathyroidism, epilepsy and severe anaemia. Each participant completed a FFQ for dietary assessment and signed a waiver of informed consent provided by the Institutional Review Board of National Taiwan University Hospital. This study was funded by the Ministry of Science and Technology.

Dietary assessments

The participants' diets were assessed using a semiquantitative FFQ. The FFQ used in this study was validated in a previous study; the questionnaire exhibited reliability and validity for identifying major nutrients in the diets of Taiwanese vegetarians and omnivores.³⁵ In this study, the nutrient analysis programme used to calculate the results of the FFQ was based on Taiwan's Food Composition Database. Carnitine intake was calculated on the basis of values published previously.³⁶ The intake of nutrients and food groups were standardised to z-scores by using linear regression adjusted for total calorie intake with residual values centred.

Oral carnitine challenge test

All the participants fasted overnight (>8 hours) before the OCCT. Fasting blood and spot urine samples were collected as baseline, and three tablets of L-carnitine fumarate (GNC) were then administered orally to the participants. Subsequently, the participants underwent time-series blood drawings with concurrent spot urine collections at 24 hours and 48 hours after the oral carnitine challenge. The participants were requested to provide urine samples within 2 hours of blood sample collections. All the participants were asked to avoid red meat, seafood and any medication during the period of the carnitine challenge test. The plasma and urine samples were aliquoted after centrifugation at 3000 rpm and stored in a freezer at -20°C . Ten carnitine tablets (GNC) were examined through HPLC to verify the amount of L-carnitine used in the OCCT. The mean dose of carnitine fumarate in each tablet was 693.5 ($\text{SD} \pm 63.2$) mg (≈ 400 mg L-carnitine/tablet).

Pharmacokinetic study of plasma TMAO by oral administration of carnitine

Thirteen independent volunteers were recruited for a pharmacokinetic study of the OCCT. After the oral carnitine challenge, blood samples were collected at 4, 8, 12, 24, 36 and 48 hours. Six of the volunteers agreed to participate in the same pharmacokinetic test again 3 months later. All the volunteers signed informed consent waivers provided by the Institutional Review Board of National Taiwan University Hospital.

Statistical analysis

Two-tailed Student's t-test or a Wilcoxon non-parametric test were used to compare group means as considered appropriate. The OR of omnivores versus vegetarians as being high TMAO producers and corresponding 95% CI were calculated using logistic regression model. The Pearson's correlation was used to calculate association between two variables. All statistics were analysed using R software V3.4.1 or GraphPad Prism (V7).

Other methods are described in online supplementary document.

Author affiliations

- ¹Department of Internal Medicine, National Taiwan University Hospital Bei-Hu Branch, Taipei, Taiwan
- ²Institute of Food Science and Technology, National Taiwan University, Taipei, Taiwan
- ³Department of Internal Medicine, National Taiwan University Hospital, Taipei, Taiwan
- ⁴Department of Life Science, National Taiwan University, Taipei, Taiwan
- ⁵Genome and Systems Biology Degree Program, Academia Sinica, Taipei, Taiwan
- ⁶Department of Internal Medicine, College of Medicine, National Taiwan University, Taipei, Taiwan
- ⁷Institute of Population Health Sciences, National Health Research Institutes, Miaoli, Taiwan
- ⁸The Metabolomics Core Laboratory, Center of Genomic Medicine, National Taiwan University, Taipei, Taiwan
- ⁹School of Pharmacy, College of Medicine, National Taiwan University, Taipei, Taiwan
- ¹⁰Department of Nutrition Therapy, Dalin Tzu Chi Hospital, Chiayi, Taiwan
- ¹¹National Applied Research Laboratories, National Laboratory Animal Center, Taipei, Taiwan
- ¹²Department of Chemistry, National Taiwan University, Taipei, Taiwan
- ¹³Department of Internal Medicine, Dalin Tzu Chi Hospital, Taipei, Taiwan
- ¹⁴Department of Food Science, Rutgers University, New Brunswick, New Jersey, USA

Acknowledgements We would like to acknowledge the service provided by the Medical Microbiota Center of the First Core Laboratory, National Taiwan University College of Medicine. The germ-free mouse facilities used in this study were provided by the Humanized Mouse Platform: Patient Derived Tumor, Gut Microbiota Transplantation and Gene Modification Service at the National Core Facility for Biopharmaceuticals (MOST 106-2319-B-492-001), National Laboratory Animal Center, National Applied Research Laboratories, Taiwan. The instrument support from NTU Mass Spectrometry Platform was acknowledged. We are also grateful for the laboratory assistance from Yu-Ting Yang, Chang-Hao Lai and Pei-Lin Wang.

Contributors W-KW designed the animal and human experiments, assisted in statistical analyses and drafted the manuscript. C-CC and H-LK assisted human specimen collection. P-YL and SP assisted in experiment, performed bioinformatics and statistical analysis. B-YL, T-YC and H-TY revised bioinformatics and statistical analysis. THTC and C-LL assisted FFQ design and analysed dietary assessments. H-LK, C-HK, C-TH and H-BZ performed metabolomics study and assisted with mass spectrometry analysis. H-LC and Y-TH provided the humanized mouse platform. P-CC and R-AC assisted the experiment and animal study. C-TH critically reviewed the manuscript. L-YC and M-SW designed the experiments, provided the funding for the study and critically revised the manuscript.

Funding This study was funded by grants from the Ministry of Science and Technology (Taiwan) (106-2314-B-002-039-MY3 and 107-2321-B-002-017) and partially supported by a grant from the Ministry of Science and Technology (Taiwan) (106-3114-B-002-003), as well as a portion of a grant from the Program for Translational Innovation of Biopharmaceutical Development-Technology Supporting Platform Axis (106-0210-01-10-01 and 107-0210-01-19-04). This work was also partially financially supported by the 'Center of Precision Medicine' from The Featured Areas Research Center Program within the framework of the Higher Education Sprout Project by the Ministry of Education in Taiwan (Grant Number: NTU-107L9014-1) and the Ministry of Science and Technology, Executive Yuan, ROC, Taiwan (Grant Number: TCTC-TR2 106-2321-B-002-025). This work was also supported by the research grant from National Taiwan University Hospital (107-T11).

Competing interests None declared.

Patient consent Obtained.

Ethics approval Institutional Review Board of National Taiwan University Hospital.

Provenance and peer review Not commissioned; externally peer reviewed.

Data sharing statement NCBI Sequence Read Archive: 16s rDNA amplicon of human omnivore (SRX4119476 – SRX4119509) and vegetarian (SRX4119396 – SRX4119418).

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

REFERENCES

- 1 Wang Z, Klipfell E, Bennett BJ, *et al*. Gut flora metabolism of phosphatidylcholine promotes cardiovascular disease. *Nature* 2011;472:57–63.

- 2 Tang WH, Wang Z, Levison BS, *et al.* Intestinal microbial metabolism of phosphatidylcholine and cardiovascular risk. *N Engl J Med* 2013;368:1575–84.
- 3 Tang WH, Hazen SL. The contributory role of gut microbiota in cardiovascular disease. *J Clin Invest* 2014;124:4204–11.
- 4 Tilg H. A Gut Feeling about Thrombosis. *N Engl J Med* 2016;374:2494–6.
- 5 Tremaroli V, Bäckhed F. Functional interactions between the gut microbiota and host metabolism. *Nature* 2012;489:242–9.
- 6 Sonnenburg JL, Bäckhed F. Diet-microbiota interactions as moderators of human metabolism. *Nature* 2016;535:56–64.
- 7 Zeisel SH, Warrier M. Trimethylamine N-oxide, the microbiome, and heart and kidney disease. *Annu Rev Nutr* 2017;37:157–81.
- 8 Brown JM, Hazen SL. Microbial modulation of cardiovascular disease. *Nat Rev Microbiol* 2018;16:171–81.
- 9 Jonsson AL, Bäckhed F. Role of gut microbiota in atherosclerosis. *Nat Rev Cardiol* 2017;14:79–87.
- 10 Flanagan JL, Simmons PA, Vehige J, *et al.* Role of carnitine in disease. *Nutr Metab* 2010;7:30.
- 11 Koeth RA, Wang Z, Levison BS, *et al.* Intestinal microbiota metabolism of L-carnitine, a nutrient in red meat, promotes atherosclerosis. *Nat Med* 2013;19:576–85.
- 12 Evans AM, Fornasini G. Pharmacokinetics of L-carnitine. *Clin Pharmacokinet* 2003;42:941–67.
- 13 Senthong V, Wang Z, Fan Y, *et al.* Trimethylamine N-oxide and mortality risk in patients with peripheral artery disease. *J Am Heart Assoc* 2016;5:5.
- 14 Li XS, Obeid S, Klingenberg R, *et al.* Gut microbiota-dependent trimethylamine N-oxide in acute coronary syndromes: a prognostic marker for incident cardiovascular events beyond traditional risk factors. *Eur Heart J* 2017;38:814–24.
- 15 Micha R, Michas G, Mozaffarian D. Unprocessed red and processed meats and risk of coronary artery disease and type 2 diabetes—an updated review of the evidence. *Curr Atheroscler Rep* 2012;14:515–24.
- 16 Koeth RA, Levison BS, Culley MK, *et al.* γ -Butyrobetaine is a proatherogenic intermediate in gut microbial metabolism of L-carnitine to TMAO. *Cell Metab* 2014;20:799–812.
- 17 Zhu W, Gregory JC, Org E, *et al.* Gut Microbial Metabolite TMAO Enhances Platelet Hyperreactivity and Thrombosis Risk. *Cell* 2016;165:111–24.
- 18 Wang Z, Roberts AB, Buffa JA, *et al.* Non-lethal Inhibition of gut microbial trimethylamine production for the treatment of atherosclerosis. *Cell* 2015;163:1585–95.
- 19 Roberts AB, Gu X, Buffa JA, *et al.* Development of a gut microbe-targeted nonlethal therapeutic to inhibit thrombosis potential. *Nat Med* 2018;24:1407–17.
- 20 Chen ML, Yi L, Zhang Y, *et al.* Resveratrol Attenuates Trimethylamine-N-Oxide (TMAO)-induced atherosclerosis by regulating TMAO Synthesis and Bile Acid Metabolism via Remodeling of the Gut Microbiota. *MBio* 2016;7:e02210–5.
- 21 Wk W, Panyod S, Ct H, *et al.* Dietary allixin reduces transformation of L-carnitine to TMAO through impact on gut microbiota. *J Funct Foods* 2015;15:408–17.
- 22 Haghighi A, Li XS, Liman TG, *et al.* Gut microbiota-dependent trimethylamine *n*-oxide predicts risk of cardiovascular events in patients with stroke and is related to proinflammatory monocytes. *Arterioscler Thromb Vasc Biol* 2018;38:2225–35.
- 23 Schiattarella GG, Sannino A, Toscano E, *et al.* Gut microbe-generated metabolite trimethylamine-N-oxide as cardiovascular risk biomarker: a systematic review and dose-response meta-analysis. *Eur Heart J* 2017;38:2948–56.
- 24 Zhu W, Wang Z, Tang WHW, *et al.* Gut microbe-generated trimethylamine *n*-oxide from dietary choline is prothrombotic in subjects. *Circulation* 2017;135:1671–3.
- 25 Wang Z, Levison BS, Hazen JE, *et al.* Measurement of trimethylamine-N-oxide by stable isotope dilution liquid chromatography tandem mass spectrometry. *Anal Biochem* 2014;455:35–40.
- 26 Rebouche CJ, Chenard CA. Metabolic fate of dietary carnitine in human adults: identification and quantification of urinary and fecal metabolites. *J Nutr* 1991;121:539–46.
- 27 do Rosario VA, Fernandes R, Trindade EB. Vegetarian diets and gut microbiota: important shifts in markers of metabolism and cardiovascular disease. *Nutr Rev* 2016;74:444–54.
- 28 Wu GD, Compher C, Chen EZ, *et al.* Comparative metabolomics in vegans and omnivores reveal constraints on diet-dependent gut microbiota metabolite production. *Gut* 2016;65:63–72.
- 29 Tang WH, Kitai T, Hazen SL. Gut microbiota in cardiovascular health and disease. *Circ Res* 2017;120:1183–96.
- 30 Zhu Y, Jameson E, Crosatti M, *et al.* Carnitine metabolism to trimethylamine by an unusual Rieske-type oxygenase from human microbiota. *Proc Natl Acad Sci U S A* 2014;111:4268–73.
- 31 Rath S, Heidrich B, Pieper DH, *et al.* Uncovering the trimethylamine-producing bacteria of the human gut microbiota. *Microbiome* 2017;5:5.
- 32 Cho CE, Taesuwan S, Malysheva OV, *et al.* Trimethylamine-N-oxide (TMAO) response to animal source foods varies among healthy young men and is influenced by their gut microbiota composition: a randomized controlled trial. *Mol Nutr Food Res* 2017;61:1600324.
- 33 Türkanoğlu Özçelik A, Can Demirdöğen B, Demirkaya S, *et al.* Flavin containing monooxygenase 3 genetic polymorphisms Glu158Lys and Glu308Gly and their relation to ischemic stroke. *Gene* 2013;521:116–21.
- 34 Craciun S, Balskus EP. Microbial conversion of choline to trimethylamine requires a glycyl radical enzyme. *Proc Natl Acad Sci U S A* 2012;109:21307–12.
- 35 Chiu TH, Huang HY, Chen KJ, *et al.* Relative validity and reproducibility of a quantitative FFQ for assessing nutrient intakes of vegetarians in Taiwan. *Public Health Nutr* 2014;17:1459–66.
- 36 Rebouche CJ, Engel AG. Kinetic compartmental analysis of carnitine metabolism in the dog. *Arch Biochem Biophys* 1983;220:60–70.