Results 130 patients received endoscopic resection for rectal neuroendocrine tumor (ESD, 45; EMR, 43; EMR-C, 42). The en bloc resection rate was 100% in all three groups. In the ESD, EMR, and EMR-C groups, R0 resection occurred in 93.3%, 60.5%, and 83.3%, respectively (EMR vs. EMR-C, $p = 0.019$; EMR vs. ESD, $p < 0.01$), while the median procedure times were 35.2, 7.1, and 6.9 min, respectively (ESD vs. EMR-C, $p < 0.01$; ESD vs. EMR, $p < 0.01$). Postoperative bleeding occurred after ESD in two patients, EMR-C in 1 patient, but not after EMR. One patient with tumor-positive margin after EMR underwent an additional surgery, and lymph node metastasis was seen. No local recurrences or distant metastatic lesions were identified in any of the patients during the follow-up period.

Conclusions EMR-C was a safe and effective treatment for rectal neuroendocrine tumors.

IDDF2019-ABS-0240

OVERLAPPING SYMPTOMS OF PEPTIC ULCER AND GASTROESOPHAGEAL REFLUX DISEASE OF PATIENTS PRESENTING WITH DYSPESIA SYNDROME IN KARAWACI, BANTEN, INDONESIA

¹Nata Pratama Hardjo Lugito*, ¹Andree Kurniawan, ¹Ignatius Bima Prasetya, ²Yohanes Chandra Kurniawan, ²Akhil Deepak Vatvani, ²Kalis Waren. ¹Department of Internal Medicine, Faculty of Medicine, Pelita Harapan University, Indonesia; ²Faculty of Medicine, Pelita Harapan University, Indonesia

10.1136/gutjnl-2019-IDDFabstracts.207

Background The dyspepsia symptoms including epigastric pain, throbbing pain, fullness, bloating, tenderness, colicky pain, mouth bitterness, heartburn, abdominal pain, vomit and decreased appetite. The dyspepsia symptoms of peptic ulcer and gastroesophageal reflux disease often overlap. Dyspepsia symptoms and disease subtypes in Asian countries were different compared to Western countries. The aim of this study is to discover the association between dyspepsia symptoms in Karawaci, Banten, Indonesia as one of the Asian countries and the diagnosis of peptic ulcer and gastroesophageal reflux disease based on endoscopy.

Methods This retrospective study used medical records of patients admitted with complaints of dyspepsia in Siloam General Hospital in Karawaci, Banten, Indonesia. Risk factors associated with peptic ulcer and gastroesophageal reflux disease such as epidemiological and clinical data collected were gender, age, symptoms, and results of endoscopy. Association between risk factors and peptic ulcer or gastroesophageal reflux disease were analyzed using Chi-Square Test and Fisher Test(bivariate analysis) and Logistic Regression (multivariate analysis).

Results A total of 199 patients with dyspepsia symptoms were admitted on the period of 2017–2018, which 51 patients undergoing upper gastrointestinal endoscopy. The dyspepsia symptoms were the complaints the patients reported, including epigastric pain, throbbing pain, fullness, bloating, tenderness, mouth bitterness, heartburn, abdominal pain, vomit and decreased appetite. There were 33 patients with upper gastrointestinal endoscopy findings in accordance with peptic ulcer and gastroesophageal reflux disease. Bivariate analysis showed risk factors associated with peptic ulcer disease were epigastric pain (OR=1.778, 95% CI 0.889 - 3.555, $p = 0.135$), bloating (OR=0.419, 95% CI 0.277 - 0.635, $p = 0.199$), and abdominal

pain (OR=2.560, 95% CI 0.774- 8.809, $p = 0.064$). Multivariate analysis showed a risk factor associated with peptic ulcer disease was abdominal pain (OR=5.333, 95% CI 0.885-32.156, $p = 0.068$).

Conclusions Abdominal pain was the only risk factor associated with peptic ulcer disease and it was not statistically significant. There were overlapping dyspepsia symptoms among patients with peptic ulcer or gastroesophageal reflux disease in Karawaci, Banten, Indonesia.

IDDF2019-ABS-0243

PREVALENCE OF SERRATED LESIONS IN AVERAGE-RISK POPULATION RECEIVING COLONOSCOPY SCREENING: A SYSTEMATIC REVIEW AND META-ANALYSIS

Junjie Huang*, Paul Chan, Tiffany Pang, Peter Choi, Jason Huang, Jingxuan Wang, Hanyue Ding, Colette Leung, Martin Wong. Jockey Club School of Public Health and Primary Care, Faculty of Medicine, Chinese University of Hong Kong, Hong Kong

10.1136/gutjnl-2019-IDDFabstracts.208

Background Serrated lesions are precursors of approximately one-third of colorectal cancers (CRCs). They can be classified into hyperplastic polyps, sessile serrated polyps (SSPs), and traditional serrated adenomas (TSAs). Despite increasing knowledge on the histological categorisation of serrated lesions, the detection rate was highly variable and endoscopist dependent. This study aims to perform a systematic review and meta-analysis to determine the overall prevalence of serrated lesions and their subtypes in average-risk population undergoing CRC screening.

Methods MEDLINE and Embase were searched to identify population-based studies that reported the prevalence of serrated lesions. Studies on average-risk populations using colonoscopy as a screening tool were included. The histologic classification of serrated lesions followed the criteria recommended by the World Health Organization. Metaprop was applied to model within-study variability by binomial distribution and Freeman-Tukey double arcsine transformation was adopted to stabilise the variances. The prevalence figures were presented by proportions and their 95% confidence intervals (CIs) using random-effects models.

Results This meta-analysis included 17 studies involving 129,001 individuals. The overall prevalence rates of serrated lesions (19.0%, 95% CI=15.3%-23.0%), SSPs (2.5%, 95%

Abstract IDDF2019-ABS-0243 Figure 1 Prevalence of serrated lesions in average risk population

CI=1.5%-3.8%) and TSAs (0.3%, 95% CI=0.1%-0.8%) were shown in figure 1 (figure 1). Subgroup analysis indicated their higher prevalence among male (22.0%, 95% CI=16.4%-28.2%) than female (14.0%, 95% CI=8.2%-21.0%), and Caucasian (25.9%, 95% CI=16.4%-28.2%) than Asian (14.6%, 95% CI=11.6%-18.2%). The prevalence of SSPs was also higher among Caucasian (2.9%, 95% CI=1.5%-4.7%) than Asian (0.7%, 95% CI=0.4%-1.2%). However, their prevalence were similar between male (2.9%, 95% CI=1.0%-5.7%) and female (2.4%, 95% CI=0.8%-4.9%).

Conclusions This was the first systematic review and meta-analysis to determine the overall prevalence of serrated lesions and their different subtypes. The pooled prevalence estimates can be used as indicators for establishing high-quality CRC screening programmes. Future studies should be performed to evaluate the factors independently associated with the presence of serrated lesions, which could inform clinical practice to enhance their detection rate.

IDDF2019-ABS-0249

COMPARISON OF POST ENDOSCOPIC MUCOSAL RESECTION BLEEDING BETWEEN EPINEPHRINE AND HYPERTONIC SALINE SUBMUCOSAL INJECTION FOR COLON POLYPS IN PATIENTS TAKING ANTITHROMBOTIC AGENTS

¹Daisuke Yamaguchi*, ²Yasuhiro Sakata, ¹Yuki Takeuchi, ¹Kei Ikeda, ¹Wataru Yoshioka, ¹Amane Jyubashi, ¹Tomohito Morisaki, ¹Keisuke Ario, ¹Seiji Tsunada. ¹Ureshino Medical Center, Japan; ²Department of Internal Medicine and Gastrointestinal Endoscopy, Saga Medical School, Japan

10.1136/gutjnl-2019-IDDFabstracts.209

Background The present study has investigated the comparison of the clinical outcomes of prophylactic submucosal saline-epinephrine injection and hypertonic saline injection for colon polyps in patients taking antithrombotic agents.

Methods We retrospectively investigated the medical records of 183 patients who taking antithrombotic agents among 891 patients who underwent colonic endoscopic mucosal resection (EMR) from January 2013 to March 2018. The patients were divided into two groups: group A, 92 patients with saline-epinephrine injection use; and group B, 91 patients with hypertonic saline (10%NaCl) injection use. Clinical characteristics, type of antithrombotic agents, treatment outcomes and adverse events were compared between the two groups. (figure 1)

Results The mean age was 73.5 and 74.0 years in group A and B, sex and patient characteristics were not different between the two groups. The main antithrombotic agents were low-dose aspirin (A: 33, B: 31 patients), warfarin (A: 9,

B: 22 patients), and clopidogrel (A: 20, B: 11 patients). Before colonic EMR, heparin bridge therapy (A: 20, B: 23 patients) and cilostazol replacement (A: 15, B: 11 patients) were equally performed.

There was no significant difference between the two groups in characteristics of colonic polyps. En bloc resection rate was similar successfully between the two groups (94.6% vs 97.8%). EMR procedure was performed by trainees more frequently in group A than B (73.9% vs 48.4%; $p < 0.001$). There were no significant differences in post EMR bleeding (7.6% vs 4.4%), immediate EMR bleeding (6.5% vs 3.3%) and other adverse events between the two groups.

Conclusions The post EMR bleeding in patients taking antithrombotic agents was similar between saline-epinephrine and hypertonic saline submucosal injection.

IDDF2019-ABS-0252

EFFECT OF MULTIDONOR INTENSIVE FECAL MICROBIOTA TRANSPLANTATION BY CAPSULES FOR ACTIVE UNCREATIVE COLITIS: A PROSPECTIVE TRIAL

¹Yanyun Fan*, ¹Qiongyun Chen, ²Bangzhou Zhang, ²Zhangran Chen, ¹Qingwen Huang, ¹Hongzhi Xu, ¹Jianlin Ren, ²Xiang Zhang. ¹Zhongshan Hospital Department of Gastroenterology, Xiamen University, China; ²Institute for Microbial Ecology, Medical College of Xiamen University, China

10.1136/gutjnl-2019-IDDFabstracts.210

Background Fecal microbiota transplantation (FMT) can induce remission in patients with ulcerative colitis(UC). We aim to establish the efficacy of multidonor, easy way FMT by capsules in active UC, and identify the factors associated with response to therapy.

Methods We conducted a prospective trial in Zhongshan Hospital Affiliated to Xiamen University, Xiamen, China. A total of 28 adults with active UC (Mayo score 4–11) were enrolled in this trial, followed by capsules 3 days one week. One donor's fecal microbiota was made into one capsule, and fecal microbiota transplantation was each derived from two or three unrelated donors. The primary outcome was steroid-free clinical remission with endoscopic remission or response. We performed 16S rDNA stool analysis to access associated microbial changes. Based on LC-MS/MS technology metabolomic researches were carried out to study metabolome of the serum.

Results The primary outcome with clinical remission was achieved in 16 out of 28 patients (57.14%), and 11 out of 22 patients(50%) had the mucosal healing. No adverse events were identified in these patients. The effects have no significant relationship with the age, gender, severity and extent of disease.FMT increased microbial diversity and altered composition. After FMT, unlike non-responders, the intestinal flora structure of responders was close to the donor, and the significant change began 1 week after transplantation. After FMT, responders had enrichment of *Alloprevotella* compared with non-responders and had increased levels of taurochenodeoxycholate and taurocholate. *Escherichia-Shigella* were decreased after FMT in responders, but no change in non-responders. The correlation analysis between the differential metabolites and the differential flora showed that the bile acid-related metabolites were positively correlated with the abundance of *Alloprevotella* and *Prevotella* 9.

Conclusions Intensive-dosing, multidonor, FMT by capsules induces clinical remission in active ulcerative colitis and is associated with the changes of intestinal flora. *Alloprevotella*

Abstract IDDF2019-ABS-0249 Figure 1 Flow diagram of the study