and bile acid-related metabolites were related to curative effect. The enrichment of *Alloprevotella* after FMT may be able to treat ulcerative colitis by regulating bile acid metabolism, improving mucosal permeability and regulating immunity. Future work should focus on the multi center verification and the relationship of immunity and gut microbiome.

IDDF2019-ABS-0253

EXPLAINED VARIANCE AND PREDICTABILITY OF INFLAMMATORY BOWEL DISEASES BY GENETIC RISK SCORE IN FIVE ASIAN POPULATIONS (RESULTS FROM THE INTERNATIONAL IBD GENETICS CONSORTIUM)

¹Shifteh Abedian*, ²Sunny H Wong, ³Suzanne Van Sommeren, ⁴Atsushi Takahashi, ⁵Jae Hee Cheon, ⁶Ajit sood, ⁷Homayoon Vahedi, ⁸Keiko Yamazaki, ⁹Won Ho Kim, ⁹Thelma BK, ¹⁰Nasser E Daryani, ⁸Michiaki Kubo, ¹¹Suk-Kyun Yang, ¹²Rupa Banerjee, ⁷Reza Malekzadeh, ³Rinse K Weersma, ²Siew C Ng, ¹Behrooz Z Alizadeh. ¹Department of Epidemiology, University of Groningen and University Medical Center Groningen, Groningen, Netherlands; ²Department of Medicine and Therapeutics, Institute of Digestive Disease, State Key Laboratory of Digestive Disease, LKS Institute of Health Science, the Chinese University, Hong Kong; ³Department of Gastroenterology and Hepatology, University of Groningen and University Medical Center Groningen, Groningen, Netherlands; ⁴Laboratory for Statistical Analysis, RIKEN, Yokohama, Japan; ⁵Department of Internal Medicine and Institute of Gastroenterology, Yonsei University College of Medicine, Seoul, Korea, South; ⁶Department of Medicine, Dayanand Medical college and Hospital, Ludhiana, India; ⁷Digestive Disease Research Center, Digestive Disease Research Institute, Tehran University of Medical Sciences, Tehran, Iran; ⁸Division of Genomic Epidemiology and Clinical Trials, Clinical Trials Research Center, Nihon University School of Medicine, Tokyo, Japan; ⁹Department of Genetics, University of Delhi South Campus, New Delhi, India; ¹⁰Department of Gastroenterology, Emam Hospital, Tehran, Iran; ¹¹Department of Gastroenterology and Hepatology, Asan Medical Center, University of Ulsan College of Medicine, Seoul, Korea, South; ¹²Department of Gastroenterology, Asian Institute of Gastroenterology, Hyderabad, India

10.1136/gutjnl-2019-IDDFabstracts.211

Background In the absence of properly designed studies, the clinical implication of genetic findings in Inflammatory Bowel Disease (IBD) is a matter of persistent debate especially in Asian population where the prevalence of IBD including Crohn's Disease (CD) and Ulcerative Colitis (UC) is rising. We aimed to investigate the predictability of IBD, CD, and UC by the means of Genetic Risk Score (GRS), in yet unaffected high-risk individuals from East Asia (EA) and Central Asia (CA).

Methods This present study included 9,698 subjects, consisting of 2,003 CD, 2,730 UC and 4,965 countries, age and gender-matched controls, genotyped on the Immunochip array of three EA (Japan, South-Korea and China) and two CA countries (India and Iran). We generated a multi-locus GRS for each population by combining information from up to 201 known genome-wide significant IBD associated variants to

summarize a total load of genetic risk for each phenotype. We estimated explained variance and predictability of IBD, CD, and UC by GRS. We shuffled the EA data into training set including two out of the three EA populations to build a model to calculate the odds ratio (OR) for each IBD variants, and a test set including the third population for the validation of predictive model built in the training set. For Indian and Iranian populations, we used the previously estimated ORs for the Caucasian population, to build GRS and test the predictive model in these two populations.

Results GRS of IBD could significantly explain up to 4.40% and 4.14% of IBD variance in EA and CA populations but given a prevalence of 0.01% and 0.04% for IBD it yields to a negligible predictive probability up to 8.8×10^{-4} and 5.52×10^{-4} . GRS of CD and UC could significantly explain CD and UC to a lesser extent compared to IBD given a lower prevalence of CD and UC (figure1).

Conclusions The present study shows that genetic findings based on Trans-ethnic analyses are applicable across Asian populations. GRS alone can explain a limited percentage of disease occurrence in general population (<5% of disease susceptibility) and may not predict IBD in the Asian populations.

IDDF2019-ABS-0254

FACTORS ASSOCIATED WITH READINESS TO SCREEN FOR COLORECTAL CANCER: A POPULATION-BASED STUDY USING THE STAGES OF CHANGE MODEL

Junjie Huang*, Maggie Chan, Jingxuan Wang, Hanyue Ding, Colette Leung, Martin Wong. Jockey Club School of Public Health and Primary Care, Faculty of Medicine, The Chinese University of Hong Kong, Hong Kong

10.1136/gutjnl-2019-IDDFabstracts.212

Background Colorectal cancer (CRC) screening is crucial in reducing its mortality yet its success depends on long-term participation. To promote the screening programme, there is a need for understanding the factors associated with its uptake. Nevertheless, most previous studies only evaluated the factors associated with screening adherence as a binary outcome. The Stages of Change model (SOC) provides an alternative means to categorise prospective participants into more precise groups. This study aims to investigate the social-demographic factors associated with readiness to CRC screening using the SOC.

Methods A population-based telephone survey was conducted among 2,400 individuals aged 61–70 years old. Data on the social-demographic factors (age, gender, educational level, marital status, occupation, income, smoking and self-perceived health status), past experience; current status; and future intention to receive CRC screening were collected

Abstract IDDF2019-ABS-0253 Figure 1 Odds ratio and explained variance in target populations per phenotype and per state