cefmetazole, fluoroquinolones, gentamicin, and trimethoprim-sulfamethoxazole after T14 and C10 at week 2 returned to the basal state at week 8 and 1 year. Although body weight and BMI slightly increased, there were significant improvements in metabolic parameters with a decrease in insulin resistance, triglycerides and low-density lipoprotein, and an increase in high-density lipoprotein. Overall, there was no significant change in the prevalence of metabolic syndrome at week 8 and 1 year, after T14, C10, and BQ10. (figure 1)

Conclusions Eradication of *H. pylori* infection has minimal disruption of the microbiota, no effect on antibiotic resistance of *E. coli* and some positive effects on metabolic parameters. These collectively lend support to the long-term safety of *H. pylori* eradication therapy

IDDF2019-ABS-0219

EFFICACY AND SAFETY OF ADDITIONAL AUTOLOGOUS PLATELET RICH STROMA IN TRANSANAL MUCOSAL ADVANCEMENT FLAP REPAIR OF TRANSSPHINCTERIC CRYPTOGLANDULAR PERIANAL FISTULAS

¹Jeanine Arkenbosch*, ²Oddeke van Ruler, ³Jeroen Stevens, ²Eelco de Graaf, ²Ruud Schouten, ¹Annemarie de Vries, ¹Janneke van der Woude. ¹Department of Gastroenterology and Hepatology, Erasmus Medical Center, Rotterdam, Netherlands; ²Department of Surgery, IJsselland Hospital, Capelle a/d IJssel, Netherlands; ³Department of Plastic and Reconstructive Surgery, Bergman Clinic, The Hague, Netherlands

10.1136/gutjnl-2019-IDDFAbstracts.10

Background Treatment of transsphincteric cryptoglandular perianal fistulas is challenging and associated with high recurrence rates. Transanal Advancement Flap Repair (TAFR) fails in almost one of every three patients, probably due to persistent chronic inflammation. Autologous Platelet-rich Stroma (PRS), platelet-rich plasma (PRP) combined with progenitor cells from autologous stromal vascular fraction (SVF), obtained from liposuction, could suppress chronic inflammation and improve success rates in TAFR. This study aimed to assess the feasibility, safety and efficacy of additional injection of autologous during TAFR of transsphincteric cryptoglandular fistulas. **Methods** 22 patients with transsphincteric cryptoglandular fistulas who underwent TAFR between December 2017 and October 2018 were prospectively included after informed consent. All patients underwent standardized TAFR and standardized preparation of autologous PRS. Inclusion criteria were transsphincteric fistulas with only one internal opening (or a second one very close by) and absence of pelvic abscess. Clinical healing was defined as the absence of symptoms and closure of the external opening at physical examination.

Abstract IDDF2019-ABS-0073 Figure 1 Compare model to others predictors for mortality in 6 weeks

Radiographic closure was defined as complete closure on post-operative MRI.

Results This study included 22 consecutive patients (10:12 male: female; median age 44 years (IQR 33–55). Follow-up data of at least 6 months was available for 18 of patients to date. All patients had one or more previous operations ranging from seton drainage to previous TAFR (3/18) or ligation of the intersphincteric fistula tract (LIFT; 2/18). Clinical healing was reached in 16 out of 18 (88.9%) patients after a median postoperative follow-up of 3 months (IQR 2.3–5.0). Two of the 18 patients did not show clinical healing at their last consultation at 4 months follow-up. Of the available 15 MRIs to date (1 is pending), 14 showed radiographic closure (93.3%). Postoperative complications encompassed thrombosis (n=1) and reoperation due to rebleeding (n=1).

Conclusions In 18 patients with transsphincteric cryptoglandular fistula treated with the addition of autologous PRS during TAFR, 93% (14/15) showed complete radiographic closure. The addition of autologous PRS appears to be feasible, safe and highly promising due to high success rates. More studies are needed to determine the exact impact.

Clinical Hepatology

IDDF2019-ABS-0073

A SIMPLIFIED PROGNOSTIC MODEL TO PREDICT MORTALITY IN PATIENTS WITH ACUTE VARICEAL BLEEDING: MULTICENTER STUDY RESULTS

Sakkarin Chirapongsathorn*, Kuntapon Akkarachinores, Amnart Chaipresert, Pawinee Saybungkla. Division of Gastroenterology and Hepatology, Phramongkutklao Hospital, College of Medicine, Thailand

10.1136/gutjnl-2019-IDDFAbstracts.11

Background Acute variceal bleeding (AVB) is a serious complication associated with high mortality. The aim of our study was to investigate mortality predictors and develop a new simple prognostic model using easily verified factors at admission in AVB patient.

Methods Between October 2012 and September 2018, 1,144 consecutive patients with AVB from Phramongkutklao hospital in Bangkok and Maharat Nakhon Sri Thummarat hospital in Nakhon Sri Thummarat were included. A simplified prognostic model was developed using multiple logistic regression after identifying significant predictors of 6-week mortality. Mortality prediction accuracy was assessed with area under the receiver operating characteristic (AUROC) curve. We compared the

Model to predict mortality in 6 weeks

Area under ROC curve (AUROC)= 0.9343

- Sensitivity 89.04%(95%CI : 86.48 – 91.60)
- Specificity 85.34%(95%CI : 82.44 – 88.44)

APASL 2019
MANILA