to 42.9), $p=0.001$), respectively. No enlargement of mean maximal target lesion diameter (mm) in group A (before vs. after; 61.37 ± 20.1 vs 64.25 ± 22.0 ($P = 0.099$), but significant increase in group B (50.1 ± 21.1 vs. 55.4 ± 18 ($p=0.017$), respectively). No significant difference of 6-month survival rate.

Conclusions EUS-RFA plus concurrent CMT significantly reduce morphine dosage requirement than CMT in UPC. RFA additionally stabilized the tumor maximal target diameter whereas CMT failed.

IDDF2019-ABS-0172

GUT MICROBIOTA SHIFT AND LOW FIBRE INTAKE IN POST GESTATIONAL DIABETES WOMEN

¹Zubaidah Hasain*, ²Nor Azmi Kamaruddin, ³Nor Azlin Mohamed Ismail, ⁴Tong Seng Fah, ⁵Nurul Huda Razalli, ¹Norfilza Mohd Mokhtar, ²Raja Affendi Raja Ali. ¹Department of Physiology, Faculty of Medicine, Universiti Kebangsaan Malaysia, Kuala Lumpur, Malaysia; ²Department of Medicine, Faculty of Medicine, Universiti Kebangsaan Malaysia Medical Centre, Kuala Lumpur, Malaysia; ³Department of Obstetrics and Gynaecology, Faculty of Medicine, Universiti Kebangsaan Malaysia Medical Centre, Kuala Lumpur, Malaysia; ⁴Department of Family Medicine, Faculty of Medicine, Universiti Kebangsaan Malaysia Medical Centre, Kuala Lumpur, Malaysia; ⁵Dietetic Programme, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, Kuala Lumpur, Malaysia

10.1136/gutjnl-2019-IDDFabstracts.180

Background Women with previous history of gestational diabetes mellitus (GDM) have unhealthy dietary patterns and profound gut microbiota shift. They have seven-fold higher risk to develop type 2 diabetes mellitus (T2DM) in the future. We hypothesised that probiotics intervention will modulate gut microbiota and reverse glucose intolerance (GI) in post GDM women. The aim of the study was to investigate the dietary patterns and gut microbiota composition in post GDM women. Secondly, we aimed to determine the effects of probiotics on the baseline anthropometric and biochemical markers of post GDM women.

Methods Baseline clinical characteristics including glucose tolerance assessment, anthropometric measurement and a 3-day dietary record of 45 post GDM women were obtained. Post GDM women were grouped based on glucose tolerance [normal glucose tolerance (NGT) and glucose intolerance (GI)]. 36 participants were assigned for 12 weeks of either probiotics or placebo intervention. Anthropometric and biochemical markers of pre and post-treatment were evaluated. Faecal samples were sent for 16S sequencing pre and post-treatment.

Results 42.2% of 45 post GDM women have postpartum GI and significantly obese as compared with the NGT group ($p<0.01$). Mean daily fiber intakes of post GDM women was significantly suboptimal according to the Malaysian dietary recommendation ($p<0.001$). Pre-treatment, total cholesterol and high-sensitivity C-reactive protein (hsCRP) of 36 participants were above the recommendation value (4.95 ± 1.01 mmol/L, 5.54 ± 5.29 mg/L). Post-treatment, change in body mass index (BMI), waist-hip ratio (WHR), HbA1c and hsCRP of ten participants differed significantly between probiotics and placebo group ($p<0.05$). At the phylum level, the gut composition of 12 post GDM women was enriched with an abundance of *Bacteroidetes* and a relatively small proportion

of *Firmicutes*, *Verrucomicrobia*, and *Proteobacteria*. Based on the genus level, relative abundance of *Prevotella_9* genus in post GDM women with GI was 20.1% compared to only 2.1% in NGT group.

Conclusions Post GDM women with glucose intolerance were obese, consumed suboptimal fibre and have gut microbiota shift similar to the T2DM adult. Roles of probiotics in post GDM women needs further validation.

IDDF2019-ABS-0173

SYSTEMATIC REVIEW OF CALCINEURIN INHIBITORS (CNI) AND VEDOLIZUMAB (VDZ) COMBINATION THERAPY IN ACUTE SEVERE ULCERATIVE COLITIS (ASUC)

Huiyu Lin*, Wee Chian Lim. Tan Tock Seng Hospital, Singapore

10.1136/gutjnl-2019-IDDFabstracts.181

Background Patients with acute severe ulcerative colitis (ASUC) may be refractory to treatment with steroids and anti-tumour necrosis factor agents (anti-TNF). Ciclosporin inhibitors (CNI) have been used effectively as a fast-acting bridge to slower-onset immunomodulators in thiopurine-naïve patients; concerns over toxicity limit prolonged use as maintenance. Patients who are azathioprine-exposed or anti-TNF-refractory have limited medical treatment options, often resulting in colectomy. Combination of CNI as induction and slower-acting but potentially safer vedolizumab (VDZ) has recently been used in patients with severe inflammatory bowel disease (IBD). We aim to review the utility in ASUC.

Methods A systematic bibliographic review was conducted on PubMed using the keywords "vedolizumab", "calcineurin inhibitors", "inflammatory bowel disease", "severe ulcerative colitis" within the period 2013 to October 2018.

Results There were 2 prospective observational studies ⁽¹⁻²⁾ [N= 30] and 1 retrospective study ⁽³⁾ [N=39]. Patients were refractory to conventional treatment with steroids and/or anti-TNF therapy. CNI (ciclosporin or tacrolimus) was used for induction of remission in majority of cases, or as rescue agent in those failing induction with Vedolizumab [subgroup of 1 study, N=7].

In 2 studies, IV cyclosporine or Tacrolimus was started; a week later, CNI-responsive patients were given VDZ induction/maintenance and CNIs were stopped after 8–12 weeks per protocol. In another study, VDZ was initiated on average 30 days after CNI, with average combination CNI+VDZ of 64 days

Combination CNI+VDZ showed good short-term efficacy. At 1 year, there was a respectable colectomy-free rate of 75%, comparable to other studies with infliximab/ciclosporin combined with azathioprine. In those receiving steroids at baseline, Steroid-free remission was achieved in $18/36 = 50\%$ at week 14. Serious adverse events (N=7) were attributed to CNIs; there were no deaths.

Conclusions Preliminary studies of combination CNI and VDZ in patients with ASUC appear promising. Further prospective trials are needed for the confirmation of the utility and efficacy of this treatment strategy in the management of ASUC