condition in sub Saharan Africa. Addressing the westernisation of the population (in lifestyle and diet) may not be as important as the provision of diagnostic facilities. The burden of colorectal neoplasia is likely to be much higher than currently reported, and more work on incidence and prevalence of these conditions could support further development of endoscopic services in Africa.

REFERENCES

1. *Eur J Cancer Prev.* 1998 Oct;7(5):38–1
2. *Niger Med J.* 2017 May-Jun; 58(3): 87–91

PWE-053 META-ANALYSIS OF TUMOUR MICROSATELLITE-INSTABILITY, AS A PREDICTOR OF RESPONSE TO FLUOROURACIL-BASED ADJUVANT CHEMOTHERAPY FOR COLON CANCER

¹Alberto Quaglia*, ²Nikhil Aggarwa, ³Mark McPhail, Kevin Monahan. ¹Royal Free Hospital London, London, UK; ²Imperial College London, London, UK; ³King's College London, London, UK

10.1136/gutjnl-2019-BSGAbstracts.377

Introduction Controversy remain on the use of 5-FU-based regimens in treating microsatellite instability-related colorectal cancer, particularly in relation to the effect of patient's age and of combining 5-FU with other drugs. The aim of this study was to carry out a meta-analysis of current literature to investigate the relationship between tumour microsatellite status and response to 5-FU based adjuvant chemotherapy including patient stratification by staging.

Methods A systematic literature review of PubMed was conducted. Studies were included/excluded based on pre-specified criteria. CRC with low MSI (MSI-L) were clustered together with MSS CRC. Overall survival at 5 years was estimated from Kaplan-Meier curves. Publication bias was investigated using funnel plots and Egger's test. Statistical analyses were conducted using the R program (version 3.2.4).

Results Out of a total of 1807 studies identified in the PubMed database until December 2018, 326 were reviews or were not written in English and were excluded. Of the 1481 remaining studies, 17 fulfilled the screening criteria. Eight were excluded because they did not include overall survival data at 5 years. Nine studies were therefore used for the meta-analysis. The quality of all studies was considered fair according to pre-defined criteria and Egger's test revealed no publication bias. There was however considerable variation between studies in terms of study design, tumour site, tumour staging, and molecular techniques used for the identification of MSI. According to the meta-analysis result 5-FU treated individuals who died at 5 years were 0.69 times more likely to have MSI than those who were alive, but this did not reach statistical significance. There were no sufficient studies to carry out subgroup analysis by stage.

Conclusions Meta-analysis based on data extracted from published literature is marred by the heterogeneity and/or paucity of clinical, pathological and molecular data in published studies. Similarly to two previous meta-analysis studies there was no significant difference in the overall survival of patients with MSI-H and MSS/MSI-L CRC treated with adjuvant 5-FU. More studies are necessary to clarify whether patients with MSI-H CRC, and in particular those at a relatively early stage, should be offered 5-FU based adjuvant chemotherapy.

PWE-054 ENDOSCOPIC INTERVENTIONS AVERT PERMANENT END OSTOMY IN HIGH-RISK FOR ADVERSE OUTCOME(HRAO) PATIENTS

^{1,2}Prabh Singh*, ¹Ikwinder Kaur, ¹Alyssa Meyer, ¹Zaheer Rizvi, ¹Victor Chedid, ¹Kevin Quinn, ¹Navtej Buttar. ¹Mayo Clinic, Rochester, USA; ²Government Medical College and Hospital, Chandigarh, India

10.1136/gutjnl-2019-BSGAbstracts.378

Introduction Anastomotic stenosis, fistulas or leaks distal to the loop ostomy either require a complex redo pelvic surgery or a potential conversion to end ostomy, both of which are associated with significant morbidity or mortality. Our aim was to assess minimally invasive endoscopic approaches as primary or rescue intervention in high-risk for adverse outcome (HRAO) patients who were otherwise destined to a permanent end ostomy.

Method All ostomy patients with endoscopic stent placement between May 2012 and July 2018 were included. Relevant demographic and clinical data were abstracted from the electronic medical record. Data was presented using descriptive methods.

Results A total of 12 (mean age 47.9y; 70% male) HRAO patients underwent minimally invasive endoscopic interventions: Eight patients had stenosis (66%), 2 patients had leak/fistula and 2 patients had stenosis as well as a fistula. The ostomy was performed for IBD in 6 (50%) patients, colorectal carcinoma in two, diverticulitis in two, rectal prolapse in one and recurrent sigmoid volvulus in one patient. Majority of the patients had ileostomy except one with colostomy and another with Altemeier procedure. 42% patients were on steroids, 58.3% were on immunosuppressants and 25% patients were smokers. 50% of the procedures were rescue interventions after reversal. Axios stents were used in 66% and Viabil in 34%. Stent diameter ranged from –5 mm and length from –0 mm. Two patients required a rendezvous antegrade-retrograde procedure. Adjunct therapies such as endoscopic suturing, transanal drainage, biocompatible plugs or biodegradable plugs were used in 33.3% of patients. On follow up, 10 patients (83%) had durable endoscopic response without the need for further interventions but two required surgery including one with permanent end ostomy. Both failures were in the patients who had strictures complicated by fistulas. One patient acutely presented with stent obstruction. In four patients stents spontaneously migrated without residual symptoms and in five, stents were removed after symptom resolution. One patient still has stent in place 8 months after endoscopic stent placement.

Conclusion Minimally invasive endoscopic management in high-risk for adverse outcome (HRAO) patients who were otherwise destined to a permanent end ostomy is a safe & effective management option. A consideration should be given to endoscopic intervention as it has potential to avert redo surgery or permanent end ostomy.

PWE-055 IRON SUPPLEMENTATION, MICROBIOME RELATED METHANOGENESIS AND CONSTIPATION – NOVEL MODEL TO EXPLAIN AN AGE-OLD PROBLEM

¹Sarah Smith*, ¹Jordan Haworth, ²Sam Treadway, ²Anthony Hobson. ¹Functional Gut Diagnostics, Manchester, UK; ²The Functional Gut Clinic, London, UK

10.1136/gutjnl-2019-BSGAbstracts.379

Introduction Oral iron supplementation is given in a broad range of conditions such as iron-deficiency anaemia but can cause significant gastrointestinal side effects in up to 70% of patients. Constipation and bloating are amongst the commonest side effects. Recent research has shown that increased methane production by archaea in the gut microbiome is related to the slowing of gut transit and constipation, via inhibition of smooth muscle contractility (1). Geobiological research has shown that iron is the most abundantly required metal in the archaeal enzymatic pathway for methanogenesis (2). We hypothesized that oral iron consumption resulted in increased methane production.

Methods A retrospective study of 396 patients who attended The Functional Gut Clinic for a lactulose or glucose hydrogen and methane breath test between June 2018 and January 2019 was carried out. Prior to the test, patients followed a 24hr low fermentable diet and 12hr fast. After providing a baseline breath sample, 10g lactulose or 75g glucose was ingested with 200mL water. Breath samples were taken every 15-minute for 135 minutes. Methane was detected using a sensitive gas chromatography technique and methane producers defined as those producing ≥ 10 ppm at any point during the study. The relationship between iron supplementation and the prevalence of breath methane was analysed via a Chi-squared test and cumulative methane production was analysed via a two-tailed T-test. Due to lack of information about iron supplementation, 59 patients were excluded.

Results Of the patients that took iron supplements, 32% produced methane, versus 17.5% of non-iron takers. A significant relationship was identified between iron supplementation and breath methane ($\chi^2(1, N=337)=4.3885, p<0.05$). Mean cumulative methane production between iron and non-iron takers (536ppm and 376ppm respectively) was statistically significant ($p<0.05$). No adverse events were reported during the test.

Conclusions This preliminary, observational study provides the basis for a new model linking a dietary intervention (ingestion of oral iron supplementation) to a gut microbiome metabolic process (methanogenesis) and a clinical outcome (constipation) which is a common, unexplained side effect of the intervention. This model can be used in prospective studies to determine risk factors for developing GI side effects and test potential therapeutic interventions aimed at disrupting methanogenesis and improving tolerability. These advances would have significant clinical and cost saving potential in a broad group of patients.

REFERENCES

1. Pimentel, M, et al. - *Am J Physiol Gastrointest Liver Physiol* 2006:290
2. Goswami, R, et al. - *Biotech*, 2016: 6:72

PWE-056 ROSIGLITAZONE: A POTENTIAL NEW STROMA TARGETED THERAPEUTIC AGENT IN COLON CANCER

¹Maria Tabuso, ²Raghu Adya, ²Mark Christian, ^{1,2}Ramesh P Arasaradnam. ¹UHCW, Coventry, UK; ²University of Warwick, Warwick, U K

10.1136/gutjnl-2019-BSGAbstracts.380

Introduction The impact of tumour associated stroma on cancer cell invasion and metastasis is an attractive emerging field. Using Nanostring Pancancer pathway we have previously identified pro-fibrotic genes collagen type 1, alpha 1 (COL1A1), fibroblast growth factor 7 (FGF7) and secreted frizzled related

protein 2 (SFRP2) significantly up regulated in human colon cancer peritumour adipose tissue compared to distal adipose tissue. Aim of this study was to evaluate the effect of Rosiglitazone, a peroxisome proliferator-activated receptor gamma (PPARG) agonist known to have anti fibrogenic properties, on COL1A1, FGF7 and SFRP2.

Methods The effect of Rosiglitazone on COL1A1 and SFRP2 was evaluated employing in vitro co-culture systems. 3T3 L1 murine adipocytes were co-cultured with HCT 116 colon cancer cells with and without 1 nM Rosiglitazone for 72 hours. COL1A1, FGF7 and SFRP2 expression were evaluated using quantitative real time-polymerase chain reaction (qRT-PCR). Statistical analysis was performed using paired Student's t-test with a cut-off of p-value <0.05 .

Results Both COL1A1 and SFRP2 resulted significantly down-regulated in co-cultured 3T3 L1 adipocytes treated with Rosiglitazone compared to untreated co-cultured 3T3 L1 adipocytes (20 fold change, $p=0.02$; 3.3 fold change, $p=0.01$ respectively). We did not observe any significant effect of Rosiglitazone on FGF7 expression in treated 3T3 L1 adipocytes compared to untreated adipocytes.

Conclusion Our results have demonstrated significant down regulation of pro-fibrotic genes COL1A1 and SFRP2 in mature adipocytes induced by Rosiglitazone. Rosiglitazone may represent a novel therapeutic agent in stromal targeted therapy in colon cancer.

PWE-057 COLON CANCER CELLS INDUCE 3T3-L1 ADIPOCYTES DE-DIFFERENTIATION THROUGH DOWN REGULATION OF PPARG

¹Maria Tabuso*, ²Mark Christian, ^{1,2}Ramesh P Arasaradnam. ¹UHCW, Coventry, UK; ²University of Warwick, Warwick, UK

10.1136/gutjnl-2019-BSGAbstracts.381

Introduction Adipocyte de-differentiation at the tumour invasive front has been demonstrated in breast, ovarian and colon cancers, although signalling mechanisms are not known. Aim of this study was to evaluate the effect of colon cancer cells on adipocyte de-differentiation.

Methods We investigated interactions between HCT 116 colon cancer cells and murine 3T3-L1 mature adipocytes using an in vitro co-culture system. HCT-116 colon cancer cells were co-cultured with 3T3-L1 adipocytes with and without 1 nM Rosiglitazone, a potent peroxisome proliferator-activated receptor gamma (PPARG) agonist that induces adipocyte differentiation, for 72 hours. Functional and morphological modifications were assessed. Fatty acid binding protein 4 (FABP4), a mature adipocyte marker, and PPARG were evaluated by quantitative real-time polymerase chain reaction (qRT-PCR). Lipid accumulation was evaluated with Oil Red O staining. Statistical analysis was performed using Student's t-test with a cut-off of p-value <0.05 .

Results 3T3-L1 mature adipocytes in co-culture with HCT-116 colon cancer cells exhibited significant down-regulation of FABP4 (5 fold change, $p<0.05$) and PPARG (2 fold change, $p<0.05$). Treatment with Rosiglitazone of 3T3-L1 adipocytes in co-culture with HCT 116 colon cancer cells could not rescue FABP4 expression compared to treated non-cultured adipocytes (5 fold change, $P<0.01$). Red Oil O staining showed marked delipidation of co-cultured adipocytes, with appearance of fibroblast-like cells.