assistance from a physician in order to benefit their patient. However, studies support that in only 23% of consultations was the patient allowed to complete their opening statement of concern, and in 69% of cases the physician interrupted the patient's statement to re-direct the enquiry to the physician's agenda. This IBD service recognised the paradigm in its approach to patient consultation. An innovative patient centred agenda sheet was developed. In order to redesign the consultation model to allow patients the opportunity to put forward their concerns to be addressed during the consultation.

Methods A patient agenda sheet was developed. Page one included; an objective, rapid assessment tool, utilising the Bristol stool chart alongside an amalgamation of the Harvey Bradshaw index and Simple Clinical Colitis Activity Index. This allows the clinician to quickly establish disease activity and severity.

Page two is a free text section which includes suggestions for discussion points during the consultation which prompts the patient to address their concerns.

Patient agenda sheets were distributed when patients arrived in clinic to be completed in the waiting room. The sheets were used by the clinicians as a prompt during the consultation.

An audit of 100 clinic letters and patient agenda sheets was undertaken to establish emerging themes. The clinician perspective regarding the use of the patient agenda sheet was also analysed.

Results Uptake was positive with a 94% completion rate.

There were six main themes identified included; medication issues, managing flare ups, fertility, travel, surveillance colonoscopy and general check-ups.

The agenda sheet reduced clinician time spent establishing disease activity and focused the consultation on addressing the patient's agenda.

Conclusion We successfully changed the consultation model from a physician's agenda to a patient centred approach. As well as empowering patients to discuss the issues that concern them, the patient agenda sheet has allowed clinicians to identify topics within the consultation which may not have previously addressed such as pregnancy and fertility. Furthermore it has facilitated rapid signposting to appropriate clinics such as Guided Self- management.

PTH-091 EFFECTS OF VEDOLIZUMAB ON HRQOL AND WORK PRODUCTIVITY IN CROHN'S DISEASE: RESULTS FROM VERSIFY

¹S Danese, ²S Adsul, ²D Lindner, ³S Jones, ⁴H Patel, ⁵JF Colombel. ¹Humanitas University, Milan, Italy; ²Takeda Pharmaceuticals International AG, Switzerland; ³Takeda Development Centre Europe Ltd, London, UK; ⁴Takeda International, Deerfield, USA; ⁵Icahn, School of Medicine at Mount Sinai, USA

10.1136/gutjnl-2019-BSGAbstracts.150

Introduction The open-label, phase 3b, single-arm VERSIFY trial demonstrated that intravenous (IV) vedolizumab (VDZ) induced endoscopic healing in patients (pts) with moderately to severely active Crohn's disease (CD). We evaluated the effects of IV VDZ on quality of life (QOL) and work productivity over a 52-week study period.

Methods 56 patients were enrolled into an IV vedolizumab 52-week substudy. QOL was assessed using Inflammatory Bowel Disease Questionnaire (IBDQ) and Euro Quality of

Life-5D (EQ-5D) utility index and visual analogue scale (VAS), and work productivity using Work Productivity and Activity Impairment (WPAI-CD). For the 52-week substudy population (n=56), changes over 52 weeks were evaluated. IBDQ remission was considered as a total IBDQ score of ≥ 170 points, with an improvement of ≥ 16 points considered clinically meaningful. Outcomes were examined by endoscopic remission status and by prior anti-tumour necrosis factor-alpha (anti-TNF α) use.

Results Mean pt age was 39.6 years, 54% were male, 43% had prior anti-TNF α treatment and 29% achieved endoscopic remission at any ileocolonoscopy visit up to Week 52. Improvements in IBDQ total score were observed as early as Week 14 and were sustained up to Week 52. At week 52 improvements were greater in pts with endoscopic remission (183 vs 164, table 1) and in pts with no prior anti-TNF α use (178 vs 157, table 1). Similar trends of greater improvements in EQ-5D utility index (0.91 vs 0.83) and VAS (79 vs 68) were observed at Week 52 in pts with endoscopic remission. At Week 52, EQ-5D utility scores improved equally regardless of prior anti-TNF α use, whereas EQ-5D VAS scores were slightly higher in pts naïve to anti-TNF α vs those who had previously failed anti-TNF α treatment. Improvements in WPAI-CD subscores were consistently higher in pts with endoscopic remission; overall work impairment and daily activities impairment were substantially improved in pts naïve to anti-TNF α .

Conclusions Overall, IV VDZ treatment was associated with substantial improvements in both QOL instruments and work productivity measures. The improvements in QOL and work productivity were greater among pts who achieved endoscopic remission and pts who had no prior anti-TNF α treatment.

Abstract PTH-091 Table 1 Changes in IBDQ from baseline to Week 52

Instrument	Baseline (Week 0)	Week 52
IBDQ Total, mean (SD) [N]		
Overall	127.2 (34.1) [55]	169.2 (35.3) [56]
Endoscopic remission - Yes	120.1 (35.2) [16]	182.8 (29.1) [16]
Endoscopic remission - No	130.1 (33.7) [39]	163.7 (36.5) [40]
Prior anti-TNF α use -Yes	128.9 (35.5) [24]	157.1 (39.1) [24]
Prior anti-TNF α use - No	125.8 (33.6) [31]	178.3 (29.7) [32]

PTH-092 EARLY 'REAL WORLD' EXPERIENCE WITH TOFACITINIB FOR MODERATE TO SEVERE ULCERATIVE COLITIS

Sailish Honap*, Esha Sharma, Shuvra Ray, Georgina Cunningham, Aravind Tamilarasan, Joel Mawdsley, Simon Anderson, Jeremy Sanderson, Peter Irving. Guy's and St. Thomas' Hospitals NHS Foundation Trust, London, UK

10.1136/gutjnl-2019-BSGAbstracts.151

Introduction Tofacitinib is an oral, small molecule Janus kinase inhibitor, which recently received NICE approval for the treatment of moderate to severe treatment refractory ulcerative colitis. We present early clinical and biochemical outcome data for a small group of new starters in a tertiary IBD referral centre.