

## COVID-19 infection in Crohn's disease under treatment with adalimumab

Jin *et al* described recently 74 cases of people having COVID-19 and experiencing gastrointestinal symptoms.<sup>1</sup> In Italy, we are managing a dramatically increasing number of people infected with severe acute respiratory syndrome corona virus 2, the virus

causing COVID-19.<sup>2,3</sup> Despite the Italian Government has implemented extraordinary measures to restrict viral spread, including significant decrease of air and train traffic within the country, cases are increasing (>75 000 people until 26 March 2020) also in other regions than Lombardy, which is the main Italian region affected from COVID-19 (>35 000 people until 26 March 2020).<sup>4</sup> This means that a lot of people under immunosuppressive-immunomodulating therapies could be infected by this virus, and that the immunosuppressive status could influence both the status of the basic disease and both the course of pulmonary disease.

A 30-year-old male was admitted on emergency department of the Bari University Hospital, Puglia Region, Southern Italy, on 12 March 2020 (with a low number of cases until that date) due to sudden occurrence of fever (38°C) associated with chest pain during breaths. He underwent to oral and nasal specimens, which resulted positive for COVID-19, and he was admitted to infectious disease division. He was affected from ileal Crohn's disease (CD), and was under treatment with mesalazine 3 g/day and adalimumab 40 mg subcutaneously every other week for 5 years due to steroid dependency. He was under sustained remission for 2 years, and no other comorbidities was reported. At entry, he needed just oxygen support; thoracic CT scan showed mild interstitial inflammation and laboratory examination (including erythrocyte sedimentation rate and C reactive protein) was normal except for a mild thrombocytopenia ( $164^3/\mu\text{L}$ , n.v.  $179\text{--}373^3/\mu\text{L}$ ). No abdominal pain was reported, and bowel movements were normal. Any specific therapy was started except for suspension of adalimumab, and he had a rapid clinical improvement: fever and chest pain disappeared within 24 hours, and at the fifth day of hospital stay he was asymptomatic (an Rx of the chest did not find sign of inflammation at that time). Significantly, no occurrence of abdominal pain or changes in bowel habits were recorded, and also faecal calprotectin assessment was normal (38 mg/kg, n.v. <50 mg/kg).

This case is of interest for two main reason. Although this infection is characterised by typical respiratory symptoms, Jin *et al*<sup>1</sup> and Wang *et al*<sup>5</sup> reported less common features like diarrhoea, nausea, vomiting and abdominal discomfort. Moreover, COVID-19-RNA was found recently in the stool of 53.42% of people tested positive for COVID-19.<sup>6</sup> If COVID-19 colonises gastrointestinal tract (unfortunately, we did not check COVID-19-RNA in stools of our patient), it could cause a worsening of a pre-existing intestinal disease as CD. Our case seems to exclude this. Since covid-19 binds ACE 2,<sup>7</sup> a possible explanation of our case could be because the patient was in clinical remission with no active inflammation, so less risk of having ACE 2 expression upregulation in the gut.<sup>8</sup>

Another interesting feature is that pulmonary disease recovered quickly despite our patient was under immunosuppressive treatment with an antitumour necrosis factor  $\alpha$  (TNF $\alpha$ ). Of course, this could be explained by the young age of our patient, because in Italy younger patients seem to have a more favourable course.<sup>4</sup> Another hypothesis is that not all immunosuppressive treatment can cause clinical worsening, probably due to different mechanisms in suppressing immune system activity. We know that TNF $\alpha$  is overexpressed also in severe respiratory distress syndrome.<sup>9</sup> In this way, current treatment with adalimumab not only could not be a risk factor for both severity of pulmonary disease and recurrence of the intestinal disease in patients with CD infected by COVID-19, but also could reinforce the hypothesis to use adalimumab in treating at least some of the infected people.<sup>2</sup> Of course, the questions raised by this case have to be addressed by the large epidemiological studies currently ongoing.

Antonio Tursi <sup>1</sup>, Gioacchino Angarano,<sup>2</sup>  
Laura Monno,<sup>2</sup> Annalisa Saracino,<sup>2</sup>  
Fabio Signorile,<sup>2</sup> Aurelia Ricciardi,<sup>2</sup> Alfredo Papa<sup>3</sup>

<sup>1</sup>Gastroenterology Service, Azienda Sanitaria Locale Barletta Andria Trani, Andria, Puglia, Italy

<sup>2</sup>Division of Infectious Diseases, Bari University Hospital, University of Bari, Bari, Puglia, Italy

<sup>3</sup>Division of Internal Medicine and Gastroenterology, Complesso Integrato "Columbus", Catholic University of the Sacred Heart Faculty of Medicine and Surgery, Roma, Lazio, Italy

**Correspondence to** Professor Antonio Tursi, Gastroenterology Service, Azienda Sanitaria Locale Barletta Andria Trani, 76123 Andria BT, Italy; antotursi@tiscali.it

**Contributors** Acquisition, analysis and interpretation of data: AT, GA, LM, AS, FS, AR and AP; drafting the work or revising it critically for important intellectual content: AT and AP; final approval of the version published, agreement to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved: AT, GA, LM, AS, FS, AR and AP.

**Funding** The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

**Competing interests** None declared.

**Patient and public involvement** Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

**Patient consent for publication** Obtained.

**Provenance and peer review** Not commissioned; internally peer reviewed.

This article is made freely available for use in accordance with BMJ's website terms and conditions for the duration of the COVID-19 pandemic or until otherwise determined by BMJ. You may use, download and print the article for any lawful, non-commercial purpose (including text and data mining) provided that all copyright notices and trade marks are retained.

© Author(s) (or their employer(s)) 2020. No commercial re-use. See rights and permissions. Published by BMJ.

**To cite** Tursi A, Angarano G, Monno L, *et al.* *Gut* 2020;**69**:1364–1365.

Received 27 March 2020

Revised 2 April 2020

Accepted 6 April 2020

Published Online First 20 April 2020

*Gut* 2020;**69**:1364–1365. doi:10.1136/gutjnl-2020-321240

#### ORCID iD

Antonio Tursi <http://orcid.org/0000-0001-5767-5541>

#### REFERENCES

- Jin X, Lian J-S, Hu J-H, *et al.* Epidemiological, clinical and virological characteristics of 74 cases of coronavirus-infected disease 2019 (COVID-19) with gastrointestinal symptoms. *Gut* 2020. doi:10.1136/gutjnl-2020-320926. [Epub ahead of print: 24 Mar 2020].
- Wu Z, McGoogan JM. Characteristics of and important lessons from the coronavirus disease 2019 (COVID-19) outbreak in China. *JAMA* 2019;2020:25–8.
- WHO. Statement on the second meeting of the International health regulations (2005) emergency Committee regarding the outbreak of novel coronavirus (2019-nCoV), 2020. Available: [https://www.who.int/news-room/detail/30-01-2020-statement-on-the-second-meeting-of-the-international-health-regulations-\(2005\)-emergency-committee-regarding-the-outbreak-of-novel-coronavirus-\(2019-ncov\)](https://www.who.int/news-room/detail/30-01-2020-statement-on-the-second-meeting-of-the-international-health-regulations-(2005)-emergency-committee-regarding-the-outbreak-of-novel-coronavirus-(2019-ncov)) [Accessed 16 Mar 2020].
- Protezione Civile proseguono. Coronavirus: sono 62.013 I positivi, 2020. Available: <http://www1.protezionecivile.gov.it/media-comunicazione/comunicati-stampa/-/content-view/view/1237989>
- Wang D, Hu B, Hu C, *et al.* Clinical characteristics of 138 hospitalized patients with 2019 novel Coronavirus–Infected pneumonia in Wuhan, China. *JAMA* 2020;323:1061.
- Xiao F, Tang M, Zheng X, *et al.* Evidence for gastrointestinal infection of SARS-CoV-2. *Gastroenterology* 2020. doi:10.1053/j.gastro.2020.02.055. [Epub ahead of print: 03 Mar 2020].
- Ciulla MM. Coronavirus uses as binding site in humans angiotensin-converting enzyme 2 functional receptor that is involved in arterial blood pressure control and fibrotic response to damage and is a drug target in cardiovascular disease. Is this just a phylogenetic coincidence? *J Med Virol* 2020. doi:10.1002/jmv.25774. [Epub ahead of print: 26 Mar 2020].
- Garg M, Royce SG, Tikellis C, *et al.* Imbalance of the renin-angiotensin system may contribute to inflammation and fibrosis in IBD: a novel therapeutic target? *Gut* 2020;69:841–51.
- Malaviya R, Laskin JD, Laskin DL. Anti-TNF $\alpha$  therapy in inflammatory lung diseases. *Pharmacol Ther* 2017;180:90–8.

