Results The average age of patients was 46 years, with men accounted for 69% of the total. After treatment, normalization of ALT 71.26%, the viral response of 90.23%, HBV DNA below the detection level was 66.3%. Liver fibrosis evaluated by FibroScan before and after 6 months treatment were 7.15 ± 1.56 kPa, and 3.58 ± 1.19 kPa evaluated by FibroScan.

Conclusions TDF was effective for patients after treatment on liver fibrosis assessed by FibroScan in chronic hepatitis B patients.

IDDF2020-ABS-0199

HISTOPATHOLOGICAL AND IMMUNOHISTOCHEMICAL CHARACTERISTICS OF HEPATOCELLULAR CARCINOMA WITH PORTAL VEIN THROMBOSIS

¹Le Thi Thu Hien*, ²Le Quoc Tuan, ³Nguyen Quang Duat. ¹Department of Internal Medicine, Thai Nguyen University of Medicine and Pharmacy, Vietnam; ²Department of Internal Medicine, Thanh Ba Hospital, Vietnam; ³Department of Internal Medicine, 103 Military Hospital, Vietnam

10.1136/gutjnl-2020-IDDF.168

Background Detection of portal vein thrombosis (PVT), especially benign or neoplastic PVT in new diagnosed hepatocellular carcinoma (HCC) has key meaning for prognosis as well as making the choice of treatment methods.

Objectives To analysis of histopathological and immunohistochemical characteristics of hepatocellular carcinoma with portal vein thrombosis

(1)To the analysis of histopathological and immunohistochemical characteristics of hepatocellular carcinoma with portal vein thrombosis

(2)To analyse the histopathological and immunohistochemical characteristics of hepatocellular carcinoma with portal vein thrombosis

Methods We conducted a prospective study of 50 patients with HCC have PVT at Clinic 103 Cam Khe from June 2017 to August 2019. The PVT specimens were collected by biopsy through the skin according to ultrasound guidance. Immunohistochemistry test: the dyeing was performed by BondMax automatic dyeing machine (Leica Biosystems - Australia).

Results The location of HCC tumors associated with the location of PVT ($p < 0.01$). Thromboses in the trunk of the portal vein of 56%, thromboses in the branches of the portal vein of 44%. The degree of thrombosis significantly related to the size of HCC tumor ($p < 0.01$). All of the PVT in patients with HCC were malignant thromboses with moderate differentiation of 58%, poor differentiation of 36%, high differentiation of 6%. Neoangiogenesis in the thromboses by

using immunohistochemistry: low level of 6%, moderate level of 46% and a high level of 48%. The neoangiogenesis in the thromboses significantly related to cancer cell differentiation ($p < 0.01$).

Conclusions The more angiogenesis was, the lower the grade of cell differentiation.

IDDF2020-ABS-0210

STATIN USE ASSOCIATED WITH REDUCED RISK OF ALL-CAUSE MORTALITY IN HEPATOCELLULAR CARCINOMA PATIENTS FOLLOWING LIVER RESECTION: A SYSTEMATIC REVIEW AND META-ANALYSIS

Philip Chun Yeung*, Siu Tim Cheung, Kelvin Kwok-Chai Ng, Paul Bo-San Lai, Charing Ching-Ning Chong. The Chinese University of Hong Kong, Hong Kong

10.1136/gutjnl-2020-IDDF.169

Background The benefit of statins in overall survival of hepatocellular carcinoma (HCC) patients after curative liver resection has been controversial. Some retrospective studies identified perioperative use of statins was associated with significantly reduction in all-cause mortality in HCC patients after liver resection, among patients with or without chronic hepatitis viral infection in multivariate survival analysis. However, there are also several studies conducted in Japan and United States against the claim. To evaluate the effect of statins on the risk of all-cause mortality in HCC patients after curative liver resections, we performed a systematic review and meta-analysis on this topic.

Methods A systematic search of Medline, Embase, Cochrane Library and Web of Science was conducted through August 2020. Studies were included if they evaluated perioperative exposure to statins, reported the all-cause mortality of HCC patients after curative resection and reported adjusted hazard ratios (HR) of multivariable analysis by Cox proportional hazards model. Summary HR estimates with 95% confidence intervals (CI) were calculated using the random-effects model.

Results The analysis included 4 studies reporting the all-cause mortality of 3,762 HCC patients after liver resection, in which 384 of them received perioperative statins treatments. A meta-analysis of the studies showed a significant (48%) reduction in the risk of all-cause mortality among HCC patients who had perioperative statins use (adjusted HR, 0.52; 95% CI, 0.34–0.82), with moderate heterogeneity among studies (Cochran's Q test, $P = 0.12$, $I^2 = 49\%$) (figure 1) with no publication bias.

Abstract IDDF2020-ABS-0210 Figure 1 Forest plot for the meta-analysis of the association between statin use and overall survival in HCC patients after liver resection

Conclusions In conclusion, perioperative statins use may reduce the risk of all-cause mortality of HCC patients after curative liver resection. Further prospective studies and larger randomized controlled trials (RCTs) are warranted to validate the conclusion.

IDDF2020-ABS-0218

THE VALUES OF ACOUSTIC RADIATION FORCE IMPULSE ELASTOGRAPHY OF LIVER IN EVALUATING THE DEGREE OF LIVER FIBROSIS IN ALCOHOLIC LIVER DISEASE

¹Ha Phuong Linh*, ²Tran Huu Ngoc. ¹Department of Internal Medicine, 103 Cam Khe Clinic, Vietnam; ²Department of Internal Medicine, Hung Vuong Hospital, Vietnam

10.1136/gutjnl-2020-IDDF.170

Background Prevalence of alcoholic liver disease (ALD) is increasingly elevated all over the world. The degree of liver fibrosis (LF) is an important factor that affects the treatment and prognosis. Acoustic radiation force impulse (ARFI) elastography is a noninvasive and a new way of assessing LF.

Aims To evaluate the diagnostic performance of ARFI elastography for assessing LF in ALD.

Methods This prospective study included 40 patients with ALD at Thai Nguyen National Hospital from June 2019 to February 2020. These patients had been diagnosed with ALD according to the guideline of the AASL 2010. Collecting samples for laboratory testing. Liver stiffness measurement was done using ARFI elastography. Liver biopsy was graded using Metavir classification.

Results Majority were in age group of 41–50 years (63%). The mean ARFI elastometry showed 2.71 ± 0.59 m/s (range 0.86–4.53) in 40 male patients. Shear wave velocity (SWV) significantly correlated with the fibrosis stage. The area under the ROC curves (AUROCs) for diagnosis of $\geq F2$ and $\geq F3$ were 0.87 (95% CI: 0.78–0.95) and 0.89 (95% CI: 0.81–0.97), respectively. The cut-off values of SWV for $\geq F2$ and $\geq F3$ were 1.98 m/s (Se 78.25%, Sp 88.37%, PPV 85.6% and NPV 82.4%) and ≥ 2.31 m/s (Se 95.68%, Sp 85.58%, PPV 71.6%, NPV 97.8%), respectively. APRI significantly correlated with the fibrosis stage. AUROCs for diagnosis of $\geq F2$ and $\geq F3$ were 0.8 (95% CI: 0.58–0.76) and 0.78 (95% CI: 0.84–0.82), respectively. The cut-off values of APRI for diagnosis of $\geq F2$ and $\geq F3$ were ≥ 0.62 (Se 51.1%, Sp 87.13%, PPV 72.3% and NPV 68.1%) and ≥ 1.171 (Se 42%, Sp 95.52%, PPV 81%, NPV 83.1%), respectively.

Conclusions Increasing SWV correlates with a higher degree of LF. ARFI elastography is a noninvasive, reliable, and repeatable diagnostic test in evaluating the degree of LF.

IDDF2020-ABS-0219

NEUTROPHIL LYMPHOCYTE RATIO AND PLATELET LYMPHOCYTE RATIO ARE EXCELLENT MARKERS FOR PREDICTING SURVIVAL AND SEVERITY OF HEPATOCELLULAR CARCINOMA

Sagar Walinjar*, Anil Arora, Ashish Kumar, Praveen Sharma, Naresh Bansal. Institute of Liver, Gastroenterology and Pancreatobiliary Sciences, Sir Ganga Ram Hospital, India

10.1136/gutjnl-2020-IDDF.171

Background Hepatocellular carcinoma (HCC) is the 5th most common malignancy worldwide. Its clinical presentation in terms of its severity is quite variable, and survival in its advanced stages is poor. Neutrophil lymphocyte ratio (NLR) and Platelet lymphocyte ratio (PLR) are the markers which have been proven useful in prognostication of many malignancies. We aimed to evaluate the predictive value of NLR and PLR in prognosticating patients with HCC and to study its association with severity of HCC.

Methods Patients with newly diagnosed HCC getting admitted between Aug 2018 and Jan 2020 were included in the study. The imaging features, ECOG status and liver functional status were analyzed, and BCLC staging was done. Patients were given appropriate treatment according to the BCLC stage. Pre-treatment NLR and PLR were calculated from differential leukocyte count. Repeat imaging was done at 1 and 3 months post-treatment. Patients were followed up for 6 months for survival. Optimum cut off values of NLR and PLR were calculated by ROC curve analysis. The OS at 6 months was compared on Kaplan Meier curve using high and low NLR, PLR values. Association between NLR and PLR was also tested with BCLC stages, recurrence-free and progression-free survival.

OVERALL SURVIVAL AT 6 MONTHS

Abstract IDDF2020-ABS-0219 Figure 1 Kaplan meier curves for overall survival at 6 months according to NLR and PLR