# ROTER

In cases of dyspepsia with or without a radiologically proven ulcer Roter will normally provide prompt symptomatic relief and lead to rapid healing of the lesion.

The reputation of Roter is world wide. It is a complete treatment in itself and requires no additional antispasmodics or anticholinergics.




FORMULA: Each Roter tablet contains: Bismuth Subnitrate
Roter (350 mg.), Magnesium Carbonate (400 mg.), Sodium
Bicarbonate (200 mg.) and Cortex Rhamni Frangulae (25 mg.).

PACKINGS OF ROTER TABLETS: Tins of 40 and 120, also dispensing sizes, 360 and 720 (P.T. exempt).

## ROTER IN PEPTIC ULCER

## No nourishment here . . .

. . . unless an adequate supply of pancreatic enzymes is available to complete digestion.




Pancrex V is the most effective preparation for replacing pancreatic secretion in the gut. It is indicated in cystic fibrosis of the pancreas, chronic pancreatitis, pancreatectomy, post-cholecystectomy and in all digestive disorders where a relative deficiency of pancreatic enzymes


is a contributory factor. Pancrex V is activated, whole dried pancreas


having a tryptic activity approximately equal to 5 times that of Pancreatin B.P.

## Pancrex V

References: Diseases of Children (1964), Blackwell, Oxford. Diseases of Infancy and Childhood. 8th Edn. (1962), Churchill, London. Lancet (1960), 1, 365. Brit. Med. J. (1958), 2, 1039.







## new Maalox gives 3 outstanding benefits

## Greater, more lasting antacid effect

Maalox has a greater acid neutralising capacity than aluminium hydroxide gel, and its action is more prolonged.

## No constipation

Because of the presence of magnesium hydroxide in Maalox, there is no danger of constipation.

## More palatable—for longer

Maalox has a pleasant taste and smooth consistency. It remains acceptable long after other antacids have grown wearisome.



These three benefits give Maalox a special value in patients who must take antacids frequently or for long periods. Maalox is a creamy white colloidal suspension of magnesium and aluminium hydroxide gels. It is presented in bottles of 16 fl. oz.

Please write for full prescribing information to

Rorer-Berk Limited
Catteshall Lane Godalming Surrey







in Sonne Dysentery

# KANTREX\* CAPSULES

STOP THE SPREAD OF INFECTION

How do you select the most effective therapy?

**FACT** "During the last few years the most popular form of treatment has been a combination of sulphonamide and oral streptomycin."

**FACT** "Treatment... has become more complicated since the appearance of drug-resistant strains of *Shigella sonnei.*"

**FACT** "...87% of *S. sonnei* were found to be resistant to sulphadiazine...for the antibiotics...the percentages of resistant strains were: potassium penicillin G, 93%; streptomycin, 17%; tetracycline,

12%; chloramphenicol, 11%; ampicillin, 6%; sodium colistimethate, 2%; kanamycin, 1%; neomycin, 1%."<sup>2</sup>

**FACT** Unlike neomycin, kanamycin does not irritate the gut wall, nor does it give rise to reactive diarrhoea.

**FACT** KANTREX CAPSULES ARE YOUR LOGICAL CHOICE FOR TREATING SONNE DYSENTERY broad-spectrum coverage...no side effects

PRESENTATION AND PRICE Basic NHS Cost: 52/4 for 30


capsules; 172/- for 100 capsules

REFERENCES 1. Leading Article, (1965) Brit. med. J. 2:893 2. Practical Notes, (1965) Practitioner, 195:714

 ${\bf *Trade\ Mark, Bristol\ Laboratories\ Limited, authorized\ user.}$ 



## A remarkably simple treatment in pancreatic enzyme replacement



Hitherto there has not been any reliable method of determining lipase activity in pancreatic extracts—none that ensured adequate control of fat digestion. This is now possible with Cotazym.

Cotazym (concentrated hog pancreas extract) enables known amounts of lipase to be administered easily and control to be established rapidly and accurately.

Cotazym is effective at lower dosage than has previously been possible and has, moreover, greater therapeutic activity. It has no offensive odour or taste and this, together with the simple method of its administration, makes it extremely acceptable to patients.

INDICATIONS

Fibrocystic disease
Pancreatic deficiency states
Post-gastrectomy malabsorption
Steatorrhoea


## COTAZYM




Each Cotazym capsule contains:

Lipase — sufficient to digest 17G. dietary fat Trypsin — sufficient to digest 34G. dietary protein Amylase — sufficient to digest 40G. dietary starch

Bottles of 100 capsules — Basic N.H.S. Price 33/4d Dosage — 1-3 capsules with each meal or as directed by the physician.



Organon Laboratories Ltd., Crown House, London Road, Morden, Surrey



# In Ulcerative Colitis

available as a 0.5 g. tablet, either plain or enteric coated

### FIRST DEMONSTRATION . . . RESULTS SIGNIFICANT

"This is the first demonstration in a formal trial that any treatment reduces the relapse-rate in ulcerative colitis, 24 out of 34 patients taking 2 g. of sulphasalazine daily remained free of symptoms for a year. This result is significantly different from that obtained in a comparable group of patients who received a placebo."

### PREFERABLE TO SYSTEMIC CORTICOSTEROIDS

"We have shown that sulphasalazine, 0.5 g. four times daily, is often effective in maintaining remission . . . and it therefore appears preferable to systemic corticosteroid drugs for this purpose."

#### **REMISSION FOR ONE YEAR**

"24 patients remained in symptomatic remission for a year while taking 2 g. of sulphasalazine daily, whereas only 8 remained symptom-free in the placebo group."

#### **NORMAL MUCOSA**

"2 of the 24 patients who remained symptom-free for a year while on sulphasalazine had a haemorrhagic mucosa at the end of the trial. The remaining 22 patients had a non-haemorrhagic mucosa which in many cases appeared normal."

#### **MINIMAL SIDE EFFECTS**

"In the present trial, using a dose of 2 g. daily, only 3 patients out of 34 had to discontinue treatment because of side-effects."

#### WHITE CELL COUNT

"In the patients treated with sulphasalazine there was no difference in the haemoglobin level before and after treatment; but the mean white-cell count was lower after six months or a year than at the start of treatment, though in no patient was it less than 4500 per c.mm."


The Lancet Jan 23rd 1965 · Vol I · Pages 185-188

# SALAZOPYRIN



Literature and further information are supplied on request SINCLAIR HOUSE  $\cdot$  THE AVENUE  $\cdot$  WEST EALING  $\cdot$  LONDON W13 Distributed by Martindale Samoore Ltd

BETTER MEDICINES FOR A BETTER WORLD



Sustained relief of peptic ulcer is now possible by adjunctive treatment with Portyn\* Templets\*, the new and highly potent sustained-release anticholinergic. Each tablet contains 5 mg. benzilonium bromide, Parke-

Davis, for immediate release and 30 mg for the gradual release which provides all-night inhibition of gastric activity. Available in bottles of 15 and 100.

\*Trade mark

## PORTYN TEMPLETS

PARKE-DAVIS HOUNSLOW-LONDON-ILLEPHEAD HOLINSTOW 2361

# NEW PERGAMON PUBLICATIONS



## The Vitamin Co-factors of Enzyme Systems

F. A. Robinson, Chairman of Crookes Laboratories Ltd.

A survey of present knowledge about the group of vitamins that are now recognized as co-factors of enzymes, describing the history of each vitamin, the method by which it was isolated, how its chemical constitution was determined, and, where appropriate, how it was synthesized. The most important natural sources of each vitamin and the effects of deficiency in humans, animals, plants, insects and micro-organisms are described in detail. Dr. Robinson also discusses the nature and properties of chemical analogues and antagonists, and the function of each vitamin as a co-enzyme.

824 pages/April 1966/£8 (\$25.00)

## The Corticovisceral Theory of The Peptic Pathogenesis of Ulcer

K. M. Bykov, Director of the I.P. Pavlov Institute of Physiology, and I. T. Kurtsin, Director of the Laboratory of Corticovisceral Pathology

This is the second of a series of carefully selected translations and surveys of Russian contributions in the fields of conditioned reflexes and cerebrovisceral and behavioural physiology. The late Dr. Bykov began his work in the laboratories of Pavlov and developed a new area of research which he referred to as corticovisceral physiology and pathology. The extensive series of studies which he undertook with Dr. Kurtsin led to the corticovisceral theory of the pathogenesis of peptic ulcer, the subject of this monograph. 352 pages/June 1966/70s. (\$11.00)

### Clio Medica

Editor: Professor Marcel Florkin, President of the International Academy for the History of Medicine, Institut Leon Fredericq, Belgium Assistant Editor: Professor Erna Lensky, Institut für Geschichte der Medizin der Universität Wien, Austria

This new quarterly journal is the official organ of the International Institute for the History of Medicine, founded in August 1962 with the object of promoting studies of a high scientific standard in the history of medicine, medical sciences and other relevant disciplines, of organizing symposia to discuss special topics in this field, and of publishing a series of scholarly studies. The *Journal* contains articles on every aspect of the history of medicine and the text is liberally illustrated with contemporary drawings and documents.

Specimen copy and details of subscription rates sent on request.

## The Structure and Metabolism of The Pancreatic Islets

Edited by S. E. Brolin, University of Uppsala, Sweden

The proceedings of a symposium held at Uppsala and Stockholm (the first to be devoted entirely to pancreatic islets). The 47 illustrated papers provide a comprehensive review of the field.

Partial Contents
The growth pattern of the endocrine pancreas in mammals – B. Hellman; The hormonal role of glucagon – R. H. Unger; Quantitative histochemistry of the islets of Langerhans and acini in different species – J. M. Kissane; On the cobalt-concentrating ability of pancreatic islet tissue – F. Knutsson
552 pages/£7 (\$20.00)

## Submicroscopic Ortho- and Patho-Morphology of The Liver


in two volumes H. David, Humboldt University, Berlin

These volumes form the first comparative submicroscopic morphology of the liver, covering investigations not only of the human liver but of that of 24 animal species of varying ages. The two volumes are complementary, the first containing text and the second some 360 photomicrographs. This work will be of considerable value to electronand light-microscopists and to all engaged in medical and surgical research.

Vol. 1 412 pages/Vol. 2 361 pages/£25 per set

Pergamon Press Limited, Headington Hill Hall, Oxford; Pergamon Press Inc., 44-01 21st Street,

Long Island City, New York 11101



Whether you call it 'mucous colitis', 'spastic colon' or 'irritable colon' the clinical picture is the same – a chronic colonic disturbance with spasm/pain, disordered bowel habit (constipation or diarrhoea) and sometimes the passage of mucus per rectum.

Merbentyl (dicyclomine hydrochloride tablets B.N.F.) can relieve spasm/pain and help restore normal bowel activity in a very short space of time.

# Merbentyl tablets of dicyclomine BNF

#### Dosage

See British National Formulary 1963, Tablets – page 171.

Low basic N.H.S. Cost An average week's treatment -- 2s. 8d.

Also available with phenobarbitone and as Merbentyl Syrup.

Full clinical information available on request.

Merrell-National (Laboratories) Ltd. London W.1

Trademarks:

Merbentyl, Merrell-National



A12068



Many clinical studies have already proved the healing influence of Biogastrone on gastric ulcer. Now confirmation comes from a new source. 'A controlled double-blind trial of carbenoxolone in the treatment of radiologically confirmed gastric ulcer is reported. Radiological examination at two weeks showed a reduction in the ulcer area of 88% in the carbenoxolone group

compared with 50% in those receiving dummy tablets (P=0.01). These results show that carbenoxolone is very effective in reducing the size of the ulcer.'\* After healing, Biogastrone 25 mg. tablets are given as a safeguard against recurrence. The dosage is one tablet twice daily.

\*Brit. med. J., (1965) 2: 1274

#### Presentation:

Tablets of 50 mg. carbenoxolone sodium for initial healing. Tablets of 25 mg. to prevent recurrence.

#### More detailed information from:

Berk Pharmaceuticals Limited, Catteshall Lane, Godalming, Surrey.



## ANTIBIOTICS AND THE INTESTINAL FLORA

Treatment with broad spectrum antibiotics suppresses the normal flora, which may be replaced by abnormal resistant organisms causing diarrhoea.

This change can be prevented or reversed by the implantation of *Lactobacillus acidophilus* which has been rendered antibiotic-resistant, and is thus able to multiply freely under these abnormal conditions. The products of growth of this organism are inimical to that of many other bacteria<sup>1,2</sup> and thus contribute to its rapid preponderance over other species.

Oral treatment with antibiotic-resistant L. acidophilus in the form of ENPAC has been shown to eliminate Candida from the faeces of infants with antibiotic-induced diarrhoea, with relief of symptoms succeeded by rapid gain in weight<sup>3,4</sup> and greatly to reduce the numbers of staphylococci in the faeces of patients treated with tetracyclines.<sup>5</sup>



## LACTOBACILLUS ACIDOPHILUS PREPARATION

Samples and literature will be sent to the medical profession on request to:—
APLIN and BARRETT LIMITED · BIOLOGICALS DIVISION · TROWBRIDGE · WILTS
TEL. TRO. 4035

## TO-DAY'S DRUGS

The most up-to-date book of its kind

"... short, authoritative, and of a high standard."

Lancet (1965), 1, 893.

"... of the greatest value to doctors in general and hospital practice."

Brit. med. J. (1965), 1, 304.

"... well-planned and moderately priced book."

Medical News (1965), No. 119, 22.

"... a volume which every physician who considers himself up to date will want to have at hand."

Practitioner (1965), 194, 439.

Price 30s. (U.S.A. \$4.50) net postage 2s. 6d. (U.S.A. 50c.) extra


400 pages

Demy Octavo



## ORDER YOUR COPY NOW

The Publishing Manager, BRITISH MEDICAL JOURNAL, London W.C.1 or any leading bookseller throughout the world



## **LONGER-LASTING DRIP THERAPY**

Gastrils—first ever antacid pastilles. Gastrils do not break up in the mouth like tablets; they dissolve slowly and smoothly, so give sustained drip therapy for the treatment of peptic ulcer.

Gastrils contain a co-dried gel of aluminium hydroxide and magnesium carbonate, a buffer antacid comparable in activity to freshly prepared liquid aluminium hydroxide gel. Gastrils rapidly buffer gastric pH to an optimum level, and keep it there, without risk of acid rebound.

To relieve the tedium of constant antacid therapy, every Gastrils prescription is dispensed in equal quantities of two delicious flavours-mint and fruit. Few antacids are as effective as Gastrils—none is as good to take!

## GASTRILS

## ANTACID PASTILLES

INDICATIONS Hyperacidity, peptic ulcer, gastritis, heartburn, oesophagitis and hiatus hernia.

**FORMULA** 

Each pastille contains 500 mg. co-dried gel of aluminium hydroxide and magnesium carbonate.

COST

BASIC N.H.S. 24 individually wrapped Gastrils in carton 3/-24 individually wrapped Gastrils from dispensing pack: 2/61d. Gastrils are prescribable on form E.C.10

'Gastrils' is a registered trade mark of T.J. Smith & Nephew Ltd.



Smith & Nephew Pharmaceuticals Limited

WELWYN GARDEN CITY · HERTFORDSHIRE • ENGLAND



# Robinul® glycopyrronium bromide (glycopyrrolate ROBINS) Robinul®PH (Robinul with phenobarbitone)

Robinul acts quickly in relieving ulcer pain and establishes a near-ideal healing environment at the ulcer site. Robinul reduces or eliminates the free HCI under both basic conditions and conditions of stress by virtue of a "medical vagotomy". By exerting a more specific pharmacological action on the gastro-intestinal tract than any other organ system, blurred vision, dry mouth etc. are avoided or occur only rarely when Robinul is given in the recommended dosage range.

Indications: In addition to its primary indications for duodenal and gastric ulcer, Robinul is indicated for other G-I conditions which may benefit from anticholinergic therapy. Robinul-PH is indicated when these situations are complicated by mild anxiety or tension.

**Dosage:** Should be adjusted according to individual patient response. Average dose is 1 or 2 tablets three times a day, in the a.m., early p.m.,

and at bedtime. In acute conditions with severe manifestations, larger dosage of plain Robinul may be indicated until symptoms subside. Thereafter, dosage should be reduced to the minimum required for maintenance of symptomatic relief.

Precautions: Administer with care to patients with incipient glaucoma. Contraindicated in acute glaucoma, prostatic hypertrophy, and in the presence of urinary bladder neck obstruction and in pyloric obstruction or stenosis with significant gastric retention. Robinul-PH is contraindicated in patients sensitive to phenobarbitone or with advanced hepatic or renal disease.

Packaging: ROBINUL (pink) and ROBINUL-PH (white) tablets are available in bottles of 50 and 500.

Reference: Practitioner (1965) Vol. No. 195, Pp. 335-339; Ann N.Y. Acad Sci (1962) 99:153 Amer J. Med Sci. (1963) 246:325.

A'H'ROBINS A. H. Robins Company Limited, Horsham, Sussex.

## BRITISH





## **IOURNAL**

A comprehensive general medical journal. Essential for every medical man or student who wishes to keep informed of research in medical science, of trends in clinical medicine and of professional activities. Weekly. Annual Subscription, United Kingdom, Irish Republic, £12 12s.; Abroad £7 7s. (\$21.00).

#### and its associated specialist journals

BRITISH JOURNAL OF OPHTHALMOLOGY

Designed for the clinician and research worker and includes original articles, case notes, information on new appliances, book reviews.

Monthly. Annual Subscription, Inland £6 6s.; Overseas £7 (\$20.00).

(Combined subscription for the two Ophthalmic journals, Inland £11 11s.; Overseas £13 (\$37.00)).

OPHTHALMIC LITERATURE

Abstracts of articles on the wide field of ophthalmology published throughout the world. Each volume contains some 5,000 abstracts—covering one year.

Annual Subscription (six issues a year plus index), Inland £6 6s.; Overseas £7 (\$20.00).
(For combined subscription see above.)

JOURNAL OF CLINICAL PATHOLOGY

Original and specially commissioned articles on each branch of pathology, with prominence given to its clinical application. Technical methods, notes on new or improved apparatus, book reviews.

Annual Subscription (six issues), Inland £A 4s.; Overseas £4 15s. (\$14.00).

BRITISH HEART JOURNAL

Original work on the heart and circulation—anatomical, physiological and pathological.

Annual Subscription (six issues), Inland £4 4s.; Overseas £4 15s. (\$14.00).

ARCHIVES OF DISEASE IN CHILDHOOD

Devoted to child health and disease, with emphasis on clinical paediatrics. Original articles by paediatricians and other specialists.

Annual Subscription (six issues), Inland £4 4s.; Overseas £4 15s. (\$14.00).

BRITISH JOURNAL OF PHARMACOLOGY AND CHEMOTHERAPY
Reports on original work in all branches of the subject, including experimental chemotherapy. Biochemical and pathological studies.

Annual Subscription (9 issues, 3 vols. in 1966), Inland £15 15s.; Overseas £16 10s. (\$48.00).

JOURNAL OF NEUROLOGY, NEUROSURGERY AND PSYCHIATRY
Original articles on current research, throwing light on practice in neurology, neurosurgery and psychiatry.

Annual Subscription (six issues), Inland £4 4s.; Overseas £4 15s. (\$14.00).

ANNALS OF THE RHEUMATIC DISEASES
Original articles by authorities in the U.K., U.S.A. and elsewhere on every manifestation of the rheumatic diseases.
Summaries in English, French and Spanish.

Annual Subscription (six issues), Inland £4 4s.; Overseas £4 15s. (\$14.00).

THORAX

Original work on the anatomy, physiology and pathology of the chest and heart. Descriptions of techniques and apparatus used in modern cardiac surgery.

Annual Subscription (six issues), Inland £4 4s.; Overseas £4 15s. (\$14.00).

GUT

Journal of the British Society of Gastroenterology. A journal for physicians, surgeons, radiologists and pathologists, carrying authoritative articles on all aspects of gastroenterology.

Annual Subscription (six issues), Inland £A 4s.; Overseas £4 15s. (\$14.00).

JOURNAL OF MEDICAL GENETICS

A new journal publishing articles on clinical, statistical, pathological, cytological and biochemical aspects of medical genetics.

Quarterly. Annual Subscription, Inland £3 3s.; Overseas £3 10s. (\$10.00).

BRITISH JOURNAL OF VENEREAL DISEASES

Designed to disseminate, exchange and increase knowledge among consultants, practitioners, research workers, teachers, Designed to disseminate, exchange and increase accounts and others concerned with the problem of venereal diseases.

Quarterly. Annual Subscription, Inland £3 3s.; Overseas £3 10s. (\$10.00).

BRITISH JOURNAL OF INDUSTRIAL MEDICINE

Authoritative papers on all aspects of occupational medicine, e.g., the pneumoconioses, occupational cancers, industrial toxicology and psychology.

Quarterly. Annual Subscription, Inland £3 3s.; Overseas £3 10s. (\$10.00). toxicology and psychology.

BRITISH JOURNAL OF PREVENTIVE AND SOCIAL MEDICINE

Original work in the field, the emphasis on epidemiological methods and medical research.

Quarterly. Annual Subscription, Inland £3 3s.; Overseas £3 10s. (\$10.00).

MEDICAL AND BIOLOGICAL ILLUSTRATION

Illustration and its application to the requirements of the medical and biological sciences.


Quarterly. Annual Subscription, Inland £3 3s.; Overseas £3 10s. (\$10.00).


ABSTRACTS OF WORLD MEDICINE

Critical abstracts of important articles appearing in medical periodicals in every part of the world. Accent on what is new.

Monthly. Annual Subscription, Inland and Overseas £6 6s. (\$18.00).

Write to The Subscription Manager, BRITISH MEDICAL JOURNAL, B.M.A. House, Tavistock Square, London W.C.1 or order through any leading subscription agent or bookseller (All dollar rates quoted refer to the U.S. dollar)





#### Formula-

Each tablet: Dried Aluminium
Hydroxide Gel, B.P.C. – 3.75 gr.
Magnesium trisilicate B.P. – 3.75 gr.
Quinalbarbitone sodium B.P.–0.50 gr.
Tinct. Belladonna B.P. – 3.m.

Packings—Regular packs of 50; Dispensing packs of 500.

Samples available on personally signed request of qualified physicians only



121 NORWOOD ROAD HERNE HILL LONDON S.E.24. Phone TULse Hill 3276

