

Supplementary table

Baseline Characteristics			
Characteristics	ITT N=46	PP N=36	Excluded from PP N=10
Age (years)			
Mean \pm SD (N)	55.1 \pm 8.4 (46)	55.7 \pm 8.0 (36)	53.0 \pm 9.9 (10)
Median (Min, Max)	55.0 (31.0, 69.0)	56.5 (31.0, 69.0)	54.0 (32.0, 67.0)
Sex [% (n/N)]			
Male	63.0% (29/46)	63.9% (23/36)	60.0% (6/10)
Female	37.0% (17/46)	36.1% (13/36)	40.0% (4/10)
Height (cm)[1]			
Mean \pm SD (N)	170.8 \pm 7.2 (46)	170.5 \pm 7.3 (36)	171.9 \pm 7.1 (10)
Median (Min, Max)	172.0 (156.0, 187.5)	172.0 (156.0, 187.5)	172.5 (160.0, 181.0)
Weight (kg)			
Mean \pm SD (N)	90.3 \pm 13.1 (42)	88.9 \pm 11.3 (33)	95.2 \pm 18.4 (9)
Median (Min, Max)	90.0 (59.0, 116.0)	87.0 (64.0, 115.0)	92.0 (59.0, 116.0)
BMI (kg/m ²)[1]			
Mean \pm SD (N)	31.6 \pm 4.4 (46)	31.3 \pm 4.2 (36)	32.6 \pm 5.2 (10)
Median (Min, Max)	30.8 (25.0, 39.6)	30.7 (25.0, 39.6)	32.6 (25.1, 39.6)
Systolic Blood Pressure (mmHg)[1]			
Mean \pm SD (N)	137.4 \pm 18.2 (46)	137.1 \pm 17.2 (36)	138.6 \pm 22.4 (10)
Median (Min, Max)	135.0 (110.0, 190.0)	135.0 (110.0, 179.0)	135.0 (110.0, 190.0)
Diastolic Blood Pressure (mmHg)[1]			
Mean \pm SD (N)	85.2 \pm 9.4 (46)	84.7 \pm 9.4 (36)	86.9 \pm 9.4 (10)
Median (Min, Max)	84.0 (60.0, 102.0)	82.5 (60.0, 102.0)	90.0 (70.0, 100.0)
Duration of T2D (years)			
Mean \pm SD (N)	6.0 \pm 2.9 (46)	6.1 \pm 2.7 (36)	5.7 \pm 3.6 (10)
Median (Min, Max)	6.4 (0.1, 11.5)	6.5 (0.1, 9.8)	6.0 (0.9, 11.5)
HbA1c (%)			
Mean \pm SD (N)	8.6 \pm 0.8 (46)	8.4 \pm 0.7 (36)	9.3 \pm 1.1 (10)
Median (Min, Max)	8.4 (7.5, 11.0)	8.3 (7.5, 9.8)	9.1 (7.5, 11.0)
Fasting Plasma Glucose (mg/dL)			
Mean \pm SD (N)	195.6 \pm 48.4 (44)	195.2 \pm 42.1 (35)	197.0 \pm 71.1 (9)

Baseline Characteristics			
Characteristics	ITT N=46	PP N=36	Excluded from PP N=10
Median (Min, Max)	195.5 (85.0, 320.7)	195.0 (114.0, 274.0)	200.0 (85.0, 320.7)
Total Cholesterol (mg/dL)			
Mean \pm SD (N)	174.3 \pm 38.7 (46)	173.1 \pm 38.1 (36)	178.3 \pm 42.7 (10)
Median (Min, Max)	169.5 (98.2, 280.0)	169.5 (98.2, 280.0)	170.1 (131.5, 267.0)
LDL-C (mg/dL)			
Mean \pm SD (N)	99.0 \pm 32.8 (46)	99.2 \pm 30.0 (36)	98.3 \pm 43.2 (10)
Median (Min, Max)	92.0 (23.2, 171.0)	92.0 (48.3, 171.0)	95.0 (23.2, 170.0)
HDL-C (mg/dL)			
Mean \pm SD (N)	44.2 \pm 9.9 (46)	43.8 \pm 10.4 (36)	45.7 \pm 8.1 (10)
Median (Min, Max)	44.5 (24.0, 67.0)	42.5 (24.0, 67.0)	45.5 (30.9, 58.0)
Triglycerides (mg/dL)			
Mean \pm SD (N)	189.9 \pm 113.6 (46)	187.2 \pm 120.8 (36)	199.5 \pm 87.5 (10)
Median (Min, Max)	160.2 (52.0, 670.0)	154.5 (52.0, 670.0)	192.0 (77.9, 398.6)
Oral Anti-Diabetic Medications at Screening [% (n/N)]			
Metformin	91.3% (42/46)	91.7% (33/36)	90.0% (9/10)
Sulfonylurea	30.4% (14/46)	25.0% (9/36)	50.0% (5/10)
Region			
Europe	82.6% (38/46)	86.1% (31/36)	70.0% (7/10)
South America	17.4% (8/46)	13.9% (5/36)	30.0% (3/10)

Note: If baseline value was not available, then used screening value.