

gutjnl-2020-322540-R1

Manuscript entitled “Neutrophils interact with cholangiocytes to cause cholestatic changes in alcoholic hepatitis”

SUPPLEMENTARY TABLE

Supplementary Table S1. Clinical data of patients with nonalcoholic steatohepatitis (NASH), primary biliary cholangitis (PBC), and primary sclerosing cholangitis (PSC)

Variable	NASH	PBC	PSC
Age (years) (Mean \pm SD)	53.1 \pm 12.6 (33-73)	58.6 \pm 7.5 (47-68)	29.2 \pm 13.7 (11-54)
Sex (M/F)	3 Male; 7 Female	1 Male; 11 Female	8 Male; 2 Female
ALT (IU/L) (Mean \pm SD)	96.1 \pm 78.9 (10-209)	38.1 \pm 23.2 (8-88)	170.4 \pm 92.0 (30-305)
AST (IU/L) (Mean \pm SD)	73.7 \pm 37.6 (34-147)	35.0 \pm 11.8 (16-58)	129.4 \pm 56.8 (63-252)
ALP (IU/L) (Mean \pm SD)	122.7 \pm 79.1 (66-289)	157.9 \pm 119.2 (51-468)	371.4 \pm 272.5 (80-999)
Fibrosis Stage	F0 (0)	F0 (1)	F3 (0)
	F1 (1)	F1 (3)	F1 (0)
	F2 (6)	F2 (8)	F2 (6)
	F3 (3)	F3 (0)	F3 (4)