poster prompts can be used to serve as a point of reference. ESS should routinely be incorporated into departmental induction and education.

P28 COMPARISON OF ANXIETY AND DEPRESSION SCORES BETWEEN 2-WEEK WAIT AND BARRETT'S SURVEILLANCE ENDOSCOPY REFERRALS

Alexander Ho*, Paul Wolfson, Ashley Wilson, Áine Hogan, Hazel McBain, Margaret Duku, Vanessa Ward, Timothy Stone, Sarmed Sami, Laurence B Lovat. *Division of Surgery and Interventional Science, University College London, London, UK*

10.1136/gutjnl-2020-bsgcampus.103

Introduction BSG guidelines recommend endoscopic surveillance for patients with Barrett's oesophagus (BE), due to the 0.5% annual risk of developing oesophageal adenocarcinoma.¹ Approximately 10% of GP 2-week wait (2WW) referrals result in a cancer diagnosis, and patients on a 2WW pathway should be told of a theoretical risk of cancer.² We therefore performed a case-control study, comparing outpatients referred to endoscopy for BE surveillance (BES) and GP 2WW referrals, to ascertain the effect of possible cancer on patients' anxiety and depression under 2 different scenarios.

Methods Patients were recruited as part of the Saliva to Predict Disease Risk (SPIT) study. This is a multicentre study to improve non-invasive prediction of the risk of BE and oesophageal cancer. Ethical approval was gained from the Coventry and Warwickshire Regional Ethics Committee (17/WM/0079). Anxiety and depression was measured using the Hospital Anxiety and Depression Scale (HADS) questionnaire; this was completed at recruitment in the endoscopy department.³ This is a validated tool consisting of 14 questions, scored from 0 to 3, with 7 questions assigned to each domain. Ordinal logistic regression analysis was performed using R software v3.6.1 to account for the effect of age and gender on HADS.

Results 940 patients, split between 363 BES referrals and 577 2WW referrals were included in the final analysis. Median age was 69 for BES and 66 for NBS ($p=0.002$). 54% of patients were female in the 2WW group compared to 24% in the BES group ($p<0.001$). Accounting for both age and gender, mean HADS anxiety score was 4.76 for BES and 6.61 for 2WW ($p<0.001$, OR=1.76; 95%CI: 1.38–2.24). Mean HADS depression score was 3.57 for BES and 4.60 for 2WW ($p<0.001$, OR=1.51; 95%CI: 1.19–1.92). Interestingly, reduced age and female gender was associated with higher anxiety scores ($p<0.001$ for both), but not depression ($p=ns$).

Conclusions These results suggest that 2WW patients undergoing endoscopy have higher baseline anxiety and depression than BES patients. Most patients on a BES list would have had at least one previous endoscopy, and may have developed expectations and adaptive mechanisms to their procedure. A previous study found a reduction in depression but not anxiety scores in patients with BE and non-specific symptoms undergoing OGD.⁴ Our study partially concurs with this; it may be that 2WW patients have an additional element of anxiety compared to a cohort with non-specific symptoms, which will need further clarification.

REFERENCES

1. Fitzgerald, R. *et al. Gut* **63**, 7–42 (2014).
2. Cummings, R. *et al. BJGP* **60**, 689–690 (2010).
3. Zigmond, A. *et al. Acta Psychiatr. Scand.* **67**, 361–370 (1983).
4. Essink-Bot, M. *et al. Qual. Life Res.* **16**, 1309–1318 (2007).

P29 ENDOSCOPIC SUBMUCOSAL DISSECTION (ESD) IN THE WESTERN SETTING – IS TUNNELING TECHNIQUE THE WAY FORWARD?

Ejaz Hossain*, Pradeep Bhandari. *Portsmouth Hospital Nhs Trust, West Byfleet, UK*

10.1136/gutjnl-2020-bsgcampus.104

Introduction Since the advent of Peroral Endoscopic Myotomy (POEMS), tunneling technique has become a popular way of performing ESD.

After initial distal dissection, proximal end of the lesion is approached, creating a submucosal tunnel. The tunnel wall is then collapsed to remove the lesion. Data from a tertiary referral centre is depicted in table 1, demonstrating tunneling technique is a safe, effective and efficient way to perform ESD, specially in Western settings.

Abstract P29 Table 1

	Average size (cm square)	Mean duration (min)	En- bloc resection	Complications	
Oesophageal (N=15)	17	99	100%	Bleeding- 0	Perforation- 0
Colorectal (N=9)	36	221	100%	Bleeding- 0	Perforation- 0

P30 SERRATED POLYP DETECTION RATE IN BOWEL CANCER SCREENING COLONOSCOPY VARIES FOUR FOLD BETWEEN COLONOSCOPISTS

Hein Htet*, Ms Jane Homer, Pete Basford. *St Richard's Hospital, Chichester, UK*

10.1136/gutjnl-2020-bsgcampus.105

Introduction Sessile serrated lesions (SSLs) are precursors for 15–30% of colorectal cancers. SSLs are more subtle and difficult to detect than conventional adenomas. We aimed to analyse factors associated with clinically relevant serrated polyp detection in the UK bowel cancer screening population.

Methods Detailed analysis of the results of 1333 BCSP colonoscopies was performed. Age, gender, FIT vs. Guaiac FOBT, endoscope definition (standard definition vs. high definition), screening vs. surveillance procedure and endoscopist were evaluated in relation to serrated polyp detection rate (SPDR), adenoma detection rate (ADR), proximal adenoma detection rate (ProxADR) and advanced ADR (AdvADR). SPDR was defined as percentage of cases with any serrated polyps proximal to the sigmoid colon or serrated polyps ≥ 5 mm in the rectum or sigmoid colon. SPSS was used for statistical analysis.

Results Of 1333 colonoscopies, 119 were excluded (incomplete data, previous colectomy, site check, bowel scope colonoscopies). Overall SPDR was 16.1% (range by endoscopist 7.6 – 31.9%). Overall ADR was 59.12% (range by endoscopist 52.3 – 72%), ProxADR 35.4% (range by endoscopist 25.4 – 52.2%) and AdvADR 25.6% (range by endoscopist 19.7 – 31.9%). SPDR was significantly associated with endoscopist ($p<0.001$), but was not associated with age, gender, FIT vs Guaiac FOBT, screening vs surveillance procedure or endoscope definition. Mean significant serrated polyps per procedure was 0.254 (range by endoscopist 0.137 – 0.637). ADR was