



OPEN ACCESS

ORIGINAL ARTICLE

# ROC-king onwards: intraepithelial lymphocyte counts, distribution & role in coeliac disease mucosal interpretation

Kamran Rostami,<sup>1</sup> Michael N Marsh,<sup>2,3</sup> Matt W Johnson,<sup>2</sup> Hamid Mohaghegh,<sup>4</sup> Calvin Heal,<sup>5</sup> Geoffrey Holmes,<sup>6</sup> Arzu Ensari,<sup>7</sup> David Aldulaimi,<sup>8</sup> Brigitte Bancel,<sup>9</sup> Gabrio Bassotti,<sup>10</sup> Adrian Bateman,<sup>11</sup> Gabriel Becheanu,<sup>12</sup> Anna Bozzola,<sup>13</sup> Antonio Carroccio,<sup>14</sup> Carlo Catassi,<sup>15</sup> Carolina Ciacci,<sup>16</sup> Alexandra Ciobanu,<sup>12</sup> Mihai Danciu,<sup>17</sup> Mohammad H Derakhshan,<sup>18,19</sup> Luca Elli,<sup>20</sup> Stefano Ferrero,<sup>20</sup> Michelangelo Fiorentino,<sup>21</sup> Marilena Fiorino,<sup>14</sup> Azita Ganji,<sup>22</sup> Kamran Ghaffarzadehgan,<sup>23</sup> James J Going,<sup>24</sup> Sauid Ishaq,<sup>25</sup> Alessandra Mandolesi,<sup>15</sup> Sherly Mathews,<sup>1</sup> Roxana Maxim,<sup>17</sup> Chris J Mulder,<sup>26</sup> Andra Neefjes-Borst,<sup>26</sup> Marie Robert,<sup>27</sup> Ilaria Russo,<sup>16</sup> Mohammad Rostami-Nejad,<sup>4</sup> Angelo Sidoni,<sup>10</sup> Masoud Sotoudeh,<sup>19</sup> Vincenzo Villanacci,<sup>13</sup> Umberto Volta,<sup>21</sup> Mohammad R Zali,<sup>4</sup> Amitabh Srivastava<sup>28</sup>

For numbered affiliations see end of article.

**Correspondence to**

Dr. Kamran Rostami, Department of Gastroenterology, Milton Keynes University Hospital, Milton Keynes MK6 5LD, UK; kamran.rostami@nhs.net

Received 8 April 2017  
Revised 17 June 2017  
Accepted 22 June 2017

**ABSTRACT**

**Objectives** Counting intraepithelial lymphocytes (IEL) is central to the histological diagnosis of coeliac disease (CD), but no definitive 'normal' IEL range has ever been published. In this multicentre study, receiver operating characteristic (ROC) curve analysis was used to determine the optimal cut-off between normal and CD (Marsh III lesion) duodenal mucosa, based on IEL counts on >400 mucosal biopsy specimens.

**Design** The study was designed at the International Meeting on Digestive Pathology, Bucharest 2015. Investigators from 19 centres, eight countries of three continents, recruited 198 patients with Marsh III histology and 203 controls and used one agreed protocol to count IEL/100 enterocytes in well-oriented duodenal biopsies. Demographic and serological data were also collected.

**Results** The mean ages of CD and control groups were 45.5 (neonate to 82) and 38.3 (2–88) years. Mean IEL count was 54±18/100 enterocytes in CD and 13±8 in normal controls (p=0.0001). ROC analysis indicated an optimal cut-off point of 25 IEL/100 enterocytes, with 99% sensitivity, 92% specificity and 99.5% area under the curve. Other cut-offs between 20 and 40 IEL were less discriminatory. Additionally, there was a sufficiently high number of biopsies to explore IEL counts across the subclassification of the Marsh III lesion.

**Conclusion** Our ROC curve analyses demonstrate that for Marsh III lesions, a cut-off of 25 IEL/100 enterocytes optimises discrimination between normal control and CD biopsies. No differences in IEL counts were found between Marsh III a, b and c lesions. There was an indication of a continuously graded dose–response by IEL to environmental (gluten) antigenic influence.

**INTRODUCTION**

Coeliac disease (CD) has a large spectrum of clinical and histological presentations. The accompanying structural changes in the upper small intestinal mucosa range through normal or near-normal to severe changes where partially reduced villi amalgamate into large, lozenge-shaped surface plateaus.<sup>1</sup> As the mucosa 'flattens', these markedly hypertrophic responses are triggered by lamina propria T cells under genetic control. Intraepithelial lymphocytes (IEL) lying within the epithelium, and surveying both lumen and lamina propria, are also implicated in this process,<sup>2</sup> although their functions and prognostic value still remain somewhat undecided. An increased population of IEL has, for almost five decades, provided a sensitive histological pointer towards CD diagnosis, made more important over the years by the Marsh classification and its emphasis on the infiltrates in normal-looking, villus-bearing mucosa.<sup>3</sup>

CD is not the only intestinal condition associated with increased IEL counts, which has resulted in the view that a lymphocytosis characterising the milder Marsh I and II lesions can be dismissed as 'non-specific'. To overcome this impasse, a recent paper dealing with the differential diagnosis of these lymphocytic infiltrates was recently published by the Bucharest Consensus<sup>4</sup> and collectively defined as microscopic enteritis.

The Bucharest Consensus was primarily concerned with subtle submicroscopic changes and IEL in coeliac and non-coeliac mucosae. However, despite the numerous papers written on this subject, we were concerned that the literature lacks a definitive cut-off point to confirm the diagnosis. First, there is no universal definition for a 'normal' IEL range, and so this makes it difficult to define a specific point of departure favouring a histological diagnosis of CD.



CrossMark

**To cite:** Rostami K, Marsh MN, Johnson MW, et al. Gut Published Online First: [please include Day Month Year]. doi:10.1136/gutjnl-2017-314297

## Significance of this study

## What is already known on this subject?

- ▶ Increased population of intraepithelial lymphocytes (IEL) provides a sensitive histological pointer towards coeliac disease (CD) diagnosis.
- ▶ IEL lying within the epithelium, and surveying both lumen and lamina propria, are implicated in disease activity, but their functions and prognostic value still remain somewhat undecided.
- ▶ Submicroscopic changes of epithelial cells and the differential diagnosis of lymphocytic infiltrates were collectively defined as microscopic enteritis.

## What are the new findings?

- ▶ We show that an effective cut-off can be obtained with receiver operating characteristic curve analysis, which substantially reduces the overlap between CD and normal controls.
- ▶ We confirm that counting of IEL performed on H&E-stained material provides optimal results, compared with CD3/ $\gamma\delta$  immunostained sections.
- ▶ We have graphically represented the cumulative totals of IEL across all our biopsies, revealing a continuous 'dose-response' curve reflective of the IEL response to luminal antigen, rather than a biphasic or clonal response that might have distinguished controls from coeliac. This has never been demonstrated before.
- ▶ Finally, with a reliable, large sample, our differential counts revealed no differences between Marsh III a, b and c, indicating that the subdivision of Marsh III has no practical value.

## How might it impact on clinical practice in the foreseeable future?

- ▶ Our differential IEL counts offer unexpected insights into mucosal interpretation that will improve diagnostic accuracy. These include bringing clarity to the IEL's implications in CD histogenesis, subdivision of Marsh III, assessing the diagnostic role of immunohistochemistry and proposing an effective cut-off to separate disease control from CD.

The earliest paper<sup>5</sup> used very thick H&E sections (5–7  $\mu\text{m}$ ) which are not ideal for making accurate counts, while suggesting that IEL populations are normally distributed. An extremely high cut-off of 40 IEL/100 enterocytes was proposed. Various immunohistochemical approaches have also been deployed since that time, but never incorporated into routine practice.<sup>6</sup> Moreover, despite its apparent sophistication and derived insights, the use of immunohistochemistry can be time consuming, requiring additional laboratory techniques and technologies which, when services are under severe time and financial constraints, may not necessarily be ideal. Moreover, since these techniques have not yielded increased degrees of diagnostic precision, they have not yet become cemented into routine clinical practice.

There is therefore a very strong case for re-examining existing laboratory procedures based on H&E-stained sections, thus to improve diagnostic accuracy beyond currently held perceptions. This study aims to bring clarity to the situation. In addition, our differential IEL counts offer further unexpected insights into mucosal biopsy interpretation.

**Table 1** Frequency of symptoms/sign among patients with coeliac disease and disease control subjects

Symptoms/signs	Patients with coeliac disease (n=198)	Disease control subjects (n=203)
Weight loss	13 (6.57)*	8 (3.94)
Diarrhoea/very loose stool	68 (34.24)	13 (6.40)
Iron deficiency anaemia	54 (27.27)	64 (31.53)
Abdominal pain	26 (13.13)	26 (12.81)
Reflux	7 (3.53)	43 (21.18)
Vomiting	2 (1.01)	5 (2.46)
Bloating	16 (8.08)	10 (4.93)
Constipation	5 (2.53)	0
Thyroiditis	5 (2.53)	0
short stature	6 (3.03)	2 (0.99)
Fatigue	2 (1.01)	2 (0.99)
Hypertransaminaemia	1 (0.51)	0
Short stature	1 (0.51)	0
Epilepsy	1 (0.51)	0
Osteoporosis	1 (0.51)	0

Note: A low ferritin was found in 27% of patients with CD; however, 15% of cases had anaemia as their main presenting symptoms.

\*Number (%).

## METHODS AND MATERIALS

This case-control study was designed during the International Meeting on Digestive Pathology held in Bucharest, November 2015. It is a multicentred international study involving eight countries, comprising 19 laboratories, each of which provided histological reviews of 10 patients with CD and 10 disease control (DC) patients. The 19 participating centres provided their data from Europe, Middle East and America, which included Ancona, Bologna, Brescia, Salerno, Perugia, Milan and Sciacca (Italy); Dudley, Glasgow, Luton and Milton Keynes (UK); Mashhad and Tehran (Iran); Amsterdam (The Netherlands); Ankara (Turkey); Bucharest and Iasi (Romania); Lyon (France) and Boston (USA). In total, this provided assessments on 401 biopsy specimens and IEL counts on 198 patients with CD (142 (71.7%) women and 56 (28.3%) men) and 203 DC patients (120 (59.1%) women and 83 (40.9%) men). The CD and DC biopsies were selected if they had 4–6 well-oriented tissue samples to allow evaluation of villous/crypt ratio and the number and distribution of the lymphocytes. The number of IEL/100 epithelial cell nuclei was counted in a continuous manner, leaving out the crypt openings. Each laboratory reanalysed the quality of cases before sending the results centrally for consolidation and review. Prominent clinical features relevant to each group are listed (table 1). The inclusion criteria for recruitment were positive coeliac serology and typical histological abnormalities consistent with Marsh III changes.<sup>3,7</sup> The subjects with DC were selected from patients with iron deficiency and a negative coeliac serology. Exclusion criteria for DC included active infections, for example, *Helicobacter pylori* or bacterial overgrowth, non-steroidal anti-inflammatory drugs use, food allergy or any other small bowel disorders. In this retrospective study, H&E-stained 4–5  $\mu\text{m}$  thick sections cut from formalin-fixed paraffin-embedded archival material were used in 162 cases. Some centres additionally employed CD3 immunohistochemistry, involving 36 patients, using commercially available anti-CD3 antibodies. The CD3 antibody clone and vendor and dilutions were different in each centre.<sup>8</sup>

The project aimed to measure the number of IEL/100 enterocytes in well-oriented duodenal biopsies from patients with untreated CD compared with a corresponding number of DC

subjects. A definitive diagnosis of CD was defined as those patients demonstrating Marsh III histological abnormalities with positive serology in line with the current and previous guidelines, whereas those with seronegative Marsh III lesions were excluded.<sup>9 10</sup> A multi-institutional collaborative effort to define optimal cut-offs to diagnose CD in patients with milder mucosal changes (Marsh I–II) is being planned as follow-up to our current study.

### Ethical considerations


This study involved re-scoring of archived histology slides, and all identifiable medical information was removed and all analyses were performed using anonymised data. The data collection was in line with good clinical practice policies with approval by research and development/audit departments of countries involved. The study was also fully approved by the ethical committee of Research Institute for Gastroenterology and Liver Disease, Shahid Beheshti University of Medical Science, Tehran under the following ethical number: IR.SBMU.RIGLD.REC.1395.87.

### Statistical analysis

Data are reported as mean ( $\pm$ SD) for continuous variables and as proportion for categorical variables. Comparisons were performed using the Student's *t*-test or  $\chi^2$  test, as appropriate. Receiver operating characteristic (ROC) curves were used to evaluate sensitivity and specificity for different cut-off levels of IEL/100 in detecting CD. Statistical significance was set at  $\alpha=0.05$ .

### RESULTS

The mean ages of DC and CD groups were 38.3 (range: 2–88) and 45.5 years (range: newborn to 82), respectively, of which 59% and 71% were women. The mean IEL count was  $54\pm 18/100$  enterocytes for CD compared with  $13\pm 8$  for DC ( $p=0.0001$ ). We present data (figure 1) from IEL counts based on the largest



**Figure 1** The numerical IEL counts (non-transformed) for the entire series of 401 specimens are shown for coeliac and disease control intestinal mucosae. The overlap of 56 between IEL counts for each population is stressed. Although when the same data are accumulated, biopsy-on-biopsy (figure 3), a perfectly shaped dose–response curve is obtained. IEL, intraepithelial lymphocytes.

series of intestinal biopsies analysed to date, obtained from eight countries across three continents (North America, Europe and Middle East). In total, 401 specimens were studied, of which 198 came from well-defined patients with CD who were either endomysial antibodies (EMA) or tissue transglutaminase antibodies (tTG) antibody positive (table 1). Of the 198 patients with CD with Marsh III, 27% had iron deficiency anaemia, 13% abdominal pain, 34% diarrhoea and 6% weight loss (table 1).

The mean  $\pm$ SD for the DC was  $13\pm 7$  IEL/100 enterocyte nuclei (95% CIs 1–27). The normality of the variable was assessed visually using a histogram/assessed using a Shapiro-Wilk test which confirmed that the data are skewed to the right and distinctly non-normal. The mean IEL count for CD was  $54\pm 18$  (95% confidence limits 19–54), differing significantly from DC ( $p<0.0001$ ). The ‘normalised’ CD mean was  $51\pm 14$  (95% confidence limits 24–78). Overall, using numerical data, CD and DC biopsies showed an overlap of 56 biopsies comprising 14% of the total specimens analysed. More interestingly with log-transformation, the distributions were ‘tightened’, so that the gross overlap was then reduced to 1%, or 38 biopsies. The literature presents a variety of optimum cut-off points ranging between 20 and 40 IEL/100 enterocyte nuclei, as criteria in the diagnosis of CD (table 2). However, these levels were, surprisingly, determined from means and SD originating from some extremely small groups.<sup>11–18</sup> In this study, we had the advantage of large numbers of CD and DC biopsies which were further evaluated through ROC curve analysis (figure 2). This offered an optimal cut-off point of 25 IEL/100 enterocytes, with a sensitivity of 0.990 and specificity of 0.931, respectively; the area under the curve was 99.5%. Here, each point on the ROC curve represents a sensitivity/specificity pair for any threshold value chosen. With perfect discrimination (no overlap in the two distributions), the ROC curve should pass through the upper left-hand corner, indicating a sensitivity and specificity of 100%. Our curve does not fulfil that criterion exactly, pointing to the inevitable overlap of IEL populations between both groups.

Using our derived ROC curve, we were able to further evaluate the currently existing published extremes (20–40 IEL) in the literature (table 2). Thus, for a high cut-off of 40 IEL, the sensitivity was reduced to 0.80 with a specificity of 1.000. At this level, the false-negative rate was 19.7%, representing a total of 31 patients with CD who would have been missed. Conversely, with the lower cut-off of 20 IEL, the sensitivity was increased to 0.995, although the specificity was now reduced to 0.081. This led to an 18.2% false-positive result, which would have represented 37 DC biopsies being incorrectly labelled as ‘possible CD’. Since the actual numbers of cases misdiagnosed at these outer IEL counts are very high, their inaccuracy as histopathological guidelines is clearly evident.

Both the numerical overlapping (figure 1) and the shape of the ROC curve (figure 2) confirmed our previously published assertion<sup>1</sup> that IEL do not represent bimodally distributed populations. This was further confirmed by our cumulative IEL count, biopsy-on-biopsy (figure 3), indicative of a graded dose–response by IEL to environmental (gluten) antigenic influence.<sup>19</sup>

By taking 25 IEL/100 enterocytes as the optimal distinguishing value, a significant reduction was seen in the numerical overlap between patients with CD and DC patients (figure 1). In most previously published series, this overlap has not previously received much in-depth analysis. Given that the IEL are not bimodally distributed, any proposed cut-off between DC and CD biopsies should be seen as arbitrary. In this series, the ROC-based overlap consisted of 3 CD women as false negatives and 12 DC as false positives. Although we had expected to have

**Table 2** Summary of literature on IEL counts

Study	Methods	Number of biopsies	Upper range	Comments
Ferguson and Murray <sup>5</sup>	H&E staining, IEL/100 enterocytes, 7 µm sections	40	40	Used controls, coeliac & autoimmune conditions, incorrect about normally distributed IEL, highest IEL count recorded: 155
Batman <i>et al</i> <sup>14</sup>	H&E staining, 5 µm sections	8	33	Study of HIV enteropathy
Hayat <i>et al</i> <sup>12</sup>	H&E staining, 4 µm sections	20	25	Counts made on uninterrupted length of epithelium >500 EC, controls defined only by a 'normal' sugar permeability
Mahadeva <i>et al</i> <sup>15</sup>	H&E staining, 3 µm sections	29	22	Major interest in normal villi with IEL infiltrate. Really difficult to infer group numbers here
Kakar <i>et al</i> <sup>16</sup>	H&E staining	12	39	Interest in normal villi with IEL infiltrates
Veress <i>et al</i> <sup>11</sup>	H&E staining, 3 µm CD3+	64	H&E: 20 CD3+: 5–9	In H&E sections, if IEL to EC ratio >5:1, do CD3 count
Biagi <i>et al</i> <sup>13</sup>	H&E staining	17	45	Major interest in promoting villous tip counts
Lähdeaho <i>et al</i> <sup>20</sup>	H&E staining	76	26	No reasons given for this cut-off value: referred to Ferguson and Murray <sup>5</sup> and Kuitunen <i>et al</i> <sup>31</sup> who give no reference range
Nasseri-Moghaddam <i>et al</i> <sup>18</sup>	H&E staining, CD45+ cells	46	H&E: 46 CD45+: 47	Establishing normal criteria by histology and immunocytology
Siriweera <i>et al</i> <sup>32</sup>	H&E staining	75	8	Retrospective study on 38 control and 37 coeliac specimens. Inexplicably small upper ranges for both groups

Notes: Here are tabulated some of the principal papers concerned with IEL counts in normal and diseased intestinal mucosae. Once identified in this way, it comes as a surprise that our perceptions of normality and of the accepted distinctions between normal and abnormal are so precariously based. Furthermore, the emphasis in some papers has been on the abnormal (HIV enteropathy), or with reference to a specific aspect (such as villus tip IEL), whereas in others, the actual numbers of patients are unclear, or a worked mean IEL±SD is not given. The very small size of some of the groups analysed should also be noted, since that has a major impact on the statistical ranges and confidence limits offered as diagnostic (with permission from Marsh & Heal.<sup>22</sup>, EC, epithelial cells; IEL, intraepithelial lymphocytes.

seen an increase in IEL counts in eastern biopsies, this impression was not borne out in practice (figure 4).

There were significant differences in the results between the UK and Romania ( $p=0.011$ ) and between Italy and Romania ( $p<0.0001$ ). We did not pursue these correlations since the numbers of biopsies per group were too small to deduce any insightful conclusions that could be safely drawn.


All 198 specimens analysed in this study fulfilled Marsh III criteria, of which 132 (66.7%) specimens were graded as IIIa, IIIb or IIIc. In graded specimens, a high proportion (60; 45%) were graded as Marsh IIIc. Conversely, 34 Marsh IIIa specimens comprised only 26%, whereas the 38 remaining specimens (29%) were classified as Marsh IIIb. The grading report was not reportable for 66 of 198 specimens, because six pathologists had already ceased grading this

lesion and two others graded biopsies as a+b or b+c or even a+b+c. The IEL populations between these three groupings, however, were not significantly different (table 3).


Six groups submitted their counts as CD3+ totals, allowing the possibility of determining percentage counts for each Marsh III subdivision. Again, no differences were found (table 3) with the arithmetic mean, being 63 immunolabelled cells/100 enterocyte nuclei.

## DISCUSSION


The results of this multicentre study across three continents indicated that: (1) the optimal cut-off point between CD and DC mucosae was 25 IEL/100 enterocyte nuclei, (2) there was no substantial difference in IEL counts from countries across the



**Figure 2** The receiver operating characteristic (ROC) curve for all 401 biopsies employed in this study is shown here. If there had been no relationship between the two samples, the curve would have followed the diagonal line. Our ROC graph shows considerable between-group homogeneity, the area under the curve being 99.5%. However, the sensitivity axis does not pass through the upper-left corner, indicative of the overlap which is further examined in detail elsewhere.



**Figure 3** The cumulative IEL count, biopsy-on-biopsy, is illustrated in this diagram. The curve does not reach 401 on account of the overlapping counts between CD and DC mucosae, the overlap between biopsies being indicated by the two vertical parallel lines. The shape of the curve resembles a dose-response (to luminal antigen) and clearly demonstrates the absence of any bimodality between the two IEL populations. CD, coeliac disease; DC, disease control; IEL, intraepithelial lymphocytes.



**Figure 4** This diagram illustrates the individual measurements from biopsies sampled across the Western hemisphere (USA, UK, France (FR), Netherlands (NL), Italy (ITAL), Romania (ROM), Turkey (TURK) and Iran). Bars reflect mean $\pm$ SD for disease control (open circles) and coeliac disease (closed circles). The two sets of horizontal dotted lines indicate the overall mean $\pm$ SD for the two groups of biopsies studied. IEL, intraepithelial lymphocytes.

Western hemisphere, (3) CD IEL counts are not normally distributed, although correction by log-transformation did not materially affect outcomes, (4) the cumulative IEL population, biopsy-on-biopsy, is indicative of a graded dose-response and thus does not represent a bimodal response between CD and DC IEL, and that any proposed cut-off value is arbitrary and (5) the total number of IEL in graded Marsh III lesions (a, b and c) was identical throughout, as was also demonstrated with the (smaller) number of CD3+ labelled IEL.

The use of elevated IEL counts has, for several decades, facilitated the histological diagnosis of CD. The original study which

highlighted this advance by Ferguson and Murray<sup>5</sup> contains several misconceptions. First, the false conclusion that IEL are normally distributed. The data from our study reveal that error, together with the skewing in the CD distribution, as shown previously.<sup>19</sup> It should be noted that there was a marked difference between the overlap as expressed by the numerically bulked data (figure 1), compared with our log-transformed results (data not illustrated). Log-transformation does 'tighten up' the distributions, reducing overlapping from 47 to 38. However, irrespective of the theoretical importance of this observation, from a practical point of view, the question arises whether, in routine practice, all suspected CD counts need to be log-transformed to provide the most critically based diagnostic information. Our analyses confirm that this is unnecessary, since log-transformation does not afford any materially different result. For example, the difference for DC distribution was only 2 IEL/100 enterocytes (figure 1), and for CD was 3 IEL. This is an important outcome of our study, and certainly an important message for histopathologists.

Second, Ferguson and Murray took 40 IEL as their upper limit of normal which is one of the highest results published (table 2). Based on our ROC curve analysis, a count of 40 IEL/100 enterocytes has a markedly reduced sensitivity of 0.790, compared with 0.995, when using a 25 IEL cut-off. Our established value corresponds to that of Hayat *et al's*<sup>12</sup> study but whose outcome was based on an exceptionally low mean count of 11 IEL/100 enterocytes. Indeed, their range was 2–26 IEL, so that with a SD of 7, their upper calculated limit of normal was 25 (11+14). It is interesting to note that they studied only 20 biopsies obtained from apparently 'healthy' subjects, defined by a 'normal' sugar permeability test. Their counts were also based on enumerating IEL along an uninterrupted length of 500 enterocytes, which to us seems a technical impossibility. Nor were there any comparisons with CD biopsies, so on these three grounds, their study cannot be regarded as a critical evaluation.

Third, Ferguson and Murray introduced the methodology of counting IEL against epithelial cells, the inaccuracy of which was demonstrated by a point-counting technique.<sup>20</sup> These revelations were subsequently confirmed by computerised morphometric analysis which showed that IEL counts were likely to be overestimated twofold.

By calculating both the number of enterocytes and IEL present in a controlled unit of villous volume, the numerical discrepancy of the earlier counting technique was exposed.<sup>19, 21</sup> The flaw is in trying to obtain a meaningful result when both parameters (epithelial cell and IEL populations) are independently changing. The absolute IEL count in untreated CD mucosae is not increased, although it appears elevated since the number of enterocytes on the surface of a Marsh stage III lesion is, comparatively, markedly reduced. Conversely, with Marsh I and Marsh II stage lesions, the increase in IEL is absolute, representing a truly elevated population. Several groups have tried to establish definitive cut-off points. However, these papers exemplify the precarious background on which current practice rests, and which is clearly in need of revision. The question of a cut-off, and how to locate it, is of both theoretical and practical importance.

As evident from the published data, a range of values lying between the extremes of 20 and 40 IEL has been proposed. Beyond that, there have been no further attempts to obtain a more reliable answer. Using our ROC curve (figure 2), we were able to investigate IEL values published in the literature. At the highest notional cut-off value of 40 IEL, we predicted that there would have been 31 false negatives (coeliac cases undiagnosed), but without any false positives. On the other hand, at the lower 20 IEL level, there would have been 37 false positives and 1 (coeliac) false negative. In

**Table 3** Breakdown of Marsh III mucosal lesions

Subgroup	Biopsies (n)	H&E IEL (mean $\pm$ SD)	Biopsies (n)	CD3+ IEL (mean $\pm$ SD)
IIIa	36	54 $\pm$ 14	10	60 $\pm$ 11
IIIb	38	52 $\pm$ 17	12	67 $\pm$ 19
IIIc	63	55 $\pm$ 21	14	62 $\pm$ 31
Total	137	54 $\pm$ 18	36	63 $\pm$ 23

Statistics: H&E: IIIa vs IIIb,  $p=0.91$ ; IIIa vs IIIc,  $p=0.96$  IIIb vs IIIc,  $p=0.96$ . CD3+: IIIa vs IIIb,  $p=0.84$ ; IIIa vs IIIc,  $p=0.92$ ; IIIb vs IIIc,  $p=0.98$ .

Note. The subclassification of the Marsh III lesion has, to our knowledge, never been subject to critical analysis despite its widespread adoption, as almost a 'matter of faith'. Here we show that the IEL counts across each of these subdivisions reveal no statistical differences. This is most worrying, since the subdivisions were supposed to represent specific, detectable and progressively deteriorating pathologies during the terminal phases of mucosal flattening. Moreover, despite smaller numbers, no detectable differences in CD3+ IEL were detected between the IIIa, b and c subdivisions of the specimens analysed, in parallel with other findings<sup>28</sup>. IEL, intraepithelial lymphocytes.

comparison with our results (with 12 disease controls and 3 coeliac misdiagnosed), the values given in the literature fare badly. The important point to remember is that any cut-off is an arbitrarily fixed point. Nevertheless, given the large numbers of specimens examined in this study, together with their respective IEL populations, the results of our analysis probably represent the best result achievable and hence the most effective cut-off level of 25 IEL on H&E-stained sections for working practice.

However, in another recent publication, IEL counts were accumulated from the preceding 40-year literature, yielding 607 separate data points. In combining H&E-derived IEL counts from both studies (giving a total of 255 CD and 271 DC biopsies), ROC curve analysis provided an optimal cut-off of 26 IEL with 95.81% of the combined sample correctly identified. Marsh and Heal<sup>22</sup> also found that separate ROC curve analyses for CD3+, and  $\gamma\delta$ +, IEL counts yielded less satisfactory cut-off points. Therefore, their results are consistent with the current Bucharest conclusion that H&E-derived IEL counts are adequate for its purpose. These are extremely interesting new insights and of great help to routine histopathologists in being reassured that immunohistochemistry does not offer improved diagnostic certainty.

IEL populations represent graded intestinal responses to antigenic challenge. This has been experimentally demonstrated in treated patients after gluten challenge.<sup>19</sup> Here we have again confirmed and revealed the dose–response curve to gluten across all the specimens studied (figure 3). On inspecting our data, we were a little surprised that IEL counts did not increase across the globe from America to the near Middle East (figure 4), which we had suspected.

The statistical differences that were demonstrable are probably of little significance and were not evaluated further. It is, however, noteworthy that the IEL count is dependent on a surprisingly large number of internal and external factors<sup>1</sup> including age, genetic background, ethnicity, geocultural factors, diet, food sensitivities, parasites, chronic liver disease, infections, drugs, the maternal and local intestinal microbiomes, and recent travel.

The large numbers of biopsies employed in this work helped to reveal the extent to which false-negative and false-positive misdiagnoses are incurred. A histologically negative result for CD appears to have less impact, since the clinician can employ additional assessments including Human Leucocyte Antigen (HLA) typing, EMA, increased anti-tTG titres or even deamidated gliadin assays<sup>23</sup> to aid the diagnosis. A more difficult issue occurs when a normal biopsy is misclassified as CD, since there may be no additional pointers to avoid imposition of a lifelong gluten-free diet.

Finally, our study has cast more light on the subclassification of the Marsh stage III coeliac lesion that originated from Dutch studies<sup>24 25</sup> and later taken over by Oberhuber *et al*<sup>26</sup> in a review without citing the studies behind it. The subdivision of Marsh III was introduced to demonstrate a correlation between serology and degrees of mucosal abnormalities that led to first description of seronegative CD concept.<sup>24</sup> Within our total sample, 132 biopsies were contributory, but we were surprised that nearly one-half (60, or 46%) were classified as Marsh IIIc, whereas only 38 (29%) were considered Marsh IIIb, and even less, 34 (25%), as Marsh IIIa. It should also be observed that we found no significant differences between the IEL populations within any of these subgroups (table 3). Although this subclassification was supposed to provide a ‘standardised procedure for mucosal classification’, no empirical measurements accompanied the modifications suggested, thus leaving this to subjective interpretation. A subsequent set of ‘guidelines for pathologists’, written by pathologists,<sup>27</sup> offered micrographs of biopsies which did not correspond to those originally given. In a separate study of CD IEL and their immunohistological subtypes,<sup>28</sup> no differences in the

‘lymphogram’ in terms of the subclassification of Marsh stage III mucosae could be found. That would be inconsistent with the original proposal that each subdivision supposedly represents a progressively greater degree of mucosal involvement and damage. Previous detailed morphological studies with transmission and scanning electronic microscopy have further degraded this subclassification<sup>7</sup> by demonstrating their lack of relevance to mucosal architecture interpretation. A recent study from Australia which employed principal component and discriminant analyses<sup>29</sup> found that defined algorithms could accurately distinguish the major Marsh grades, but were unable to discriminate between any of the so-called Marsh IIIa, b and c sublesions. Together, this growing body of information seems to indicate that attempted subclassifications of Marsh stage III lesions have no intrinsic status or value, and therefore would best be abandoned.

In that regard, six contributory groups within this study had already abandoned this subclassification, whereas two other groups classified their biopsies with multiple combinations of a+b+c, indicating the impossibility, if not gross frustration, of forcing one subclassification to fit into one or the other category.

This might be an explanation for the anomalous finding that almost 50% of biopsies were classified as Marsh IIIc, with far less weight put on the other two subcategories. Clearly, if morphological, immunohistochemical or mathematically based analyses fail to reveal any major differences in the progressive degrees of tissue damage implied by this subclassification, then what value can emerge from its continued use? Equally strained is the growing habit of equating Marsh IIIa with so-called ‘partial villous atrophy’, Marsh IIIb with ‘subtotal villous atrophy’ and Marsh IIIc with ‘total villous atrophy’. Interestingly, in flat lesions, there is no evidence of an atrophic process, but only hypertrophic mucosal responses,<sup>1</sup> and a striking absence of villi.

In summary, based on a large intercontinental study of IEL, involving over 400 biopsies, we have firmly established that a cut-off point of 25 IEL/100 enterocyte nuclei provides a very effective criterion for distinguishing CD Marsh III stage biopsies from DC tissues. No obvious trends in IEL counts were found in biopsies sampled from across the three continents. The CD IEL count is not normally distributed yet, as we have shown, this does not make any substantial difference to laboratory counting in the routine work of histopathologists. We found no differences when examining IEL populations throughout the three proposed subdivisions of the Marsh III lesion, suggesting that these subdivisions (a, b, c) are spurious, lacking any useful histopathological content, while offering no diagnostic or prognostic assistance whatsoever to the gastroenterologist.

#### Author affiliations

- <sup>1</sup>Department of Gastroenterology and Pathology, Milton Keynes University Hospital, Milton Keynes, UK
- <sup>2</sup>Department of Gastroenterology, Luton and Dunstable University Hospital, Luton, UK
- <sup>3</sup>Wolfson College, University of Oxford, Oxford, UK
- <sup>4</sup>Gastroenterology and Liver Diseases Research Centre, Research Institute for Gastroenterology and Liver Diseases, Shahid Beheshti University of Medical Sciences, Tehran, The Islamic Republic of Iran
- <sup>5</sup>Centre for Biostatistics, Faculty of Biology, Academic Health Science Centre, University of Manchester, Manchester, UK
- <sup>6</sup>Department of Gastroenterology, Royal Derby Hospital, Derby, UK
- <sup>7</sup>Department of Pathology, Ankara University Medical School, Ankara, Turkey
- <sup>8</sup>Department of Gastroenterology, Warwick Hospital, Warwick, UK
- <sup>9</sup>Service de Pathologie, Centre de Biologie et Pathologie Groupe Hospitalier du Nord, Hospices Civils de Lyon, Lyon, France
- <sup>10</sup>University of Perugia Medical School, Perugia, Italy
- <sup>11</sup>Department of Cellular Pathology, University Hospital Southampton NHS Foundation Trust, Southampton, UK
- <sup>12</sup>Department of Pathology, Carol Davila University of Medicine and Pharmacy, Bucharest, Romania

<sup>13</sup>Institute of Pathology Spedali Civili, Brescia, Italy

<sup>14</sup>Internal Medicine and Pathology Unit, University of Palermo, Giovanni Paolo II Hospital, Sciacca, Italy

<sup>15</sup>Department of Pediatrics and Surgical Pathology, Università Politecnica delle Marche, Ancona, Italy

<sup>16</sup>Department of Medicine and Surgery, Scuola Medica Salernitana, University of Salerno, Salerno, Italy

<sup>17</sup>Departments of Gastroenterology and Pathology, Grigore T. Popa University of Medicine and Pharmacy, Iasi, Romania

<sup>18</sup>College of Medical, Veterinary and Life Sciences, University of Glasgow, Glasgow, UK

<sup>19</sup>Digestive Disease Research Center, Tehran University Medical Science, Tehran, Iran

<sup>20</sup>Center for Prevention and Diagnosis of Coeliac Disease and Pathology Unit, Fondazione IRCCS Ca' granda Ospedale Maggiore Policlinico, Milano, Italy

<sup>21</sup>Department of Medical and Surgical Sciences, University of Bologna and Diagnostic and Experimental, University of Bologna, Bologna, Italy

<sup>22</sup>Gastroenterology and Hepatology, Faculty of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran

<sup>23</sup>Pathology department, Razavi hospital, Mashhad, Iran

<sup>24</sup>Department of Pathology, Southern General Hospital, Lanarkshire, UK

<sup>25</sup>Dudley group of Hospitals, Birmingham City University, Birmingham, UK

<sup>26</sup>Department of Hepatogastroenterology and Pathology, Free University Medical Centre, Amsterdam, The Netherlands

<sup>27</sup>Department of Pathology and Medicine, Yale University School of Medicine, New Haven, USA

<sup>28</sup>Department of Pathology, Brigham & Women's Hospital, Boston, USA

**Correction notice** This article has been corrected since it published Online First. The affiliations for Sauid Ishaq, Masoud Sotoudeh and Stefano Ferrero have been updated.

**Acknowledgements** The Authors would like to thank Coeliac UK for making this study accessible by funding the open access publication fee.

**Contributors** Forty-one pathologists and gastroenterologists participated in this study by contributing towards pathology and clinical data, discussion on study concept, statistical analysis and writing and revising the manuscript. The study was designed during the second consensus meeting incorporated in the International Course of Pathology in Bucharest. KR designed and coordinated the study, organised the second consensus meeting with help and support from GBe. He also collected the data and wrote the first draft of the manuscript. HM performed the first statistical analysis and contributed towards data interpretation. CH performed the second analysis and MNM rewrote the draft and revised the interpretation of data. MWJ and GH critically reviewed the manuscript and MWJ contributed with clinical and histology data. GBe provided data and co-organised the consensus meeting. MHD and JGG contributed with their data, critical revision, drafting and finalising the manuscript. ASr, AE, VV, UV, MD, ACa, ANB, CCi, MRN and LE contributed in the conceptualisation and design of the study, data collection and critical revision of the manuscript. BB, GBa, ABo, CCa, ACi, SF, MIF, MaF, AG, KG, SI, AM, SM, RM, CJM, IR, ASi and MRZ contributed with their pathology and clinical data and revision of the manuscript.

**Competing interests** None declared.

**Ethics approval** Research and development/audit departments of countries involved and the ethical committee of Research Institute for Gastroenterology and Liver Diseases, Shahid Beheshti University of Medical Sciences, Tehran under the following ethical number: IR.SBMU.RIGLD.REC.1395.87.

**Provenance and peer review** Not commissioned; externally peer reviewed.

**Open Access** This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

© Article author(s) (or their employer(s) unless otherwise stated in the text of the article) 2017. All rights reserved. No commercial use is permitted unless otherwise expressly granted.

## REFERENCES

- Marsh MN, Rostami K. What is a normal intestinal mucosa? *Gastroenterology* 2016;151:784–8.
- Holmes GK, Asquith P, Stokes PL, et al. Cellular infiltrate of jejunal biopsies in adult coeliac disease in relation to gluten withdrawal. *Gut* 1974;15:278–83.
- Marsh MN. Gluten, major histocompatibility complex, and the small intestine. A molecular and immunobiologic approach to the spectrum of gluten sensitivity ('celiac sprue'). *Gastroenterology* 1992;102:330–54.
- Rostami K, Aldulaimi D, Holmes G, et al. Microscopic enteritis: Bucharest consensus. *World J Gastroenterol* 2015;21:2593–604.
- Ferguson A, Murray D. Quantitation of intraepithelial lymphocytes in human jejunum. *Gut* 1971;12:988–94.
- Järvinen TT, Kaukinen K, Laurila K, et al. Intraepithelial lymphocytes in celiac disease. *Am J Gastroenterol* 2003;98:1332–7.
- Marsh MN, Johnson MW, Rostami K. Mucosal histopathology in celiac disease: a rebuttal of Oberhuber's sub-division of Marsh III. *Gastroenterol Hepatol Bed Bench* 2015;8:99–109.
- Ubiali A, Villanacci V, Facchetti F, et al. Is TCR gamma clonality assay useful to detect early celiac disease? *J Clin Gastroenterol* 2007;41:275–9.
- Ludvigsson JF, Bai JC, Biagi F, et al. Diagnosis and management of adult coeliac disease: guidelines from the British Society of Gastroenterology. *Gut* 2014;63:1210–28.
- United European Gastroenterology. When is a coeliac a coeliac? Report of a working group of the United European Gastroenterology Week in Amsterdam, 2001. *Eur J Gastroenterol Hepatol* 2001;13:1123–8.
- Veress B, Franzén L, Bodin L, et al. Duodenal intraepithelial lymphocyte-count revisited. *Scand J Gastroenterol* 2004;39:138–44.
- Hayat M, Cairns A, Dixon MF, et al. Quantitation of intraepithelial lymphocytes in human duodenum: what is normal? *J Clin Pathol* 2002;55:393–4.
- Biagi F, Luinetti O, Campanella J, et al. Intraepithelial lymphocytes in the villous tip: do they indicate potential coeliac disease? *J Clin Pathol* 2004;57:835–9.
- Batman PA, Miller AR, Forster SM, et al. Jejunal enteropathy associated with human immunodeficiency virus infection: quantitative histology. *J Clin Pathol* 1989;42:275–81.
- Mahadeva S, Wyatt JJ, Howdle PD. Is a raised intraepithelial lymphocyte count with normal duodenal villous architecture clinically relevant? *J Clin Pathol* 2002;55:424–8.
- Kakar S, Nehra V, Murray JA, et al. Significance of intraepithelial lymphocytosis in small bowel biopsy samples with normal mucosal architecture. *Am J Gastroenterol* 2003;98:2027–33.
- Pellegrino S, Villanacci V, Sansotta N, et al. Redefining the intraepithelial lymphocytes threshold to diagnose gluten sensitivity in patients with architecturally normal duodenal histology. *Aliment Pharmacol Ther* 2011;33:697–706.
- Nasseri-Moghaddam S, Mofid A, Nouraei M, et al. The normal range of duodenal intraepithelial lymphocytes. *Arch Iran Med* 2008;11:136–42.
- Crowe PT, Marsh MN. Morphometric analysis of intestinal mucosa. VI—Principles in enumerating intra-epithelial lymphocytes. *Virchows Arch* 1994;424:301–6.
- Guix M, Skinner JM, Whitehead R. Measuring intraepithelial lymphocytes, surface area, and volume of lamina propria in the jejunal mucosa of coeliac patients. *Gut* 1979;20:275–8.
- Crowe PT, Marsh MN. Morphometric analysis of small intestinal mucosa. IV. Determining cell volumes. *Virchows Arch A Pathol Anat Histopathol* 1993;422:459–66.
- Marsh MN, Heal CJ. Evolutionary Developments in Interpreting the Gluten-Induced Mucosal Celiac Lesion: An Archimedean Heuristic. *Nutrients* 2017;9:213.
- Mooney PD, Wong SH, Johnston AJ, et al. Increased detection of celiac disease with measurement of deamidated gliadin peptide antibody before endoscopy. *Clin Gastroenterol Hepatol* 2015;13:1278–84.
- Rostami K, Kerckhaert J, von Blomberg BM, et al. SAT and serology in adult coeliacs, seronegative coeliac disease seems a reality. *Neth J Med* 1998;53:15–19.
- Rostami K, Kerckhaert J, Tiemessen R, et al. Sensitivity of antiendomysium and antigliadin antibodies in untreated celiac disease: disappointing in clinical practice. *Am J Gastroenterol* 1999;94:888–94.
- Oberhuber G, Granditsch G, Vogelsang H. The histopathology of coeliac disease: time for a standardized report scheme for pathologists. *Eur J Gastroenterol Hepatol* 1999;11:1185–94.
- Dickson BC, Streutker CJ, Chetty R. Coeliac disease: an update for pathologists. *J Clin Pathol* 2006;59:1008–16.
- De Andrés A, Camarero C, Roy G. Distal duodenum versus duodenal bulb: intraepithelial lymphocytes have something to say in celiac disease diagnosis. *Dig Dis Sci* 2015;60:1004–9.
- Charlesworth RPG, Andronicos NM, Scott DR, et al. Can the sensitivity of the histopathological diagnosis of coeliac disease be increased and can treatment progression be monitored using mathematical modelling of histological sections? – A pilot study. *Adv Med Sci* 2017;62:136–42.
- Lähdeaho ML, Kaukinen K, Collin P, et al. Celiac disease: from inflammation to atrophy: a long-term follow-up study. *J Pediatr Gastroenterol Nutr* 2005;41:44–8.
- Kuitunen P, Kosnai I, Savilahti E. Morphometric study of the jejunal mucosa in various childhood enteropathies with special reference to intraepithelial lymphocytes. *J Pediatr Gastroenterol Nutr* 1982;1:525–32.
- Siriweera E, Qi E, Yong E. Validity of intraepithelial lymphocyte counts in the diagnosis of celiac disease: A histopathological study. *Int J Celiac Dis* 2015;3:136–42.