

Table 5. Morphologic Features to Evaluate on CECT (online version only)

1. Pancreatic parenchymal necrosis: None <30% 30-50% >50%

2. Peripancreatic necrosis

3. Pancreatic/peripancreatic fluid or collections

a. Location

Intrapancreatic, where _____

Peripancreatic, where _____

b. Characteristics of fluid

Homogenous

Heterogeneous

c. Well-demarcated wall (measure thickness in mm)

No

Yes

d. Extraluminal loculated gas bubbles

No

Yes

e. Gas/fluid level

No

Yes

f. Shape of collection

Round or oval

Irregular

4. Related extrapancreatic findings

- a. Cholelithiasis
- b. Choledocholithiasis
- c. Extrahepatic biliary dilation
- d. Portal venous thrombosis/obstruction

Gastroesophageal varices

- e. Superior mesenteric venous thrombosis/obstruction
- f. Splenic vein thrombosis/obstruction

Gastric varices

- g. Arterial pseudoaneurysm

Where, describe location, size: _____

- h. Pleural effusions
- i. Ascites
- j. Inflammatory involvement of organs

Stomach

Duodenum

Jejunum

Colon

Appendix

Liver

Kidney (right / left)

Ureter (right / left)

- k. Colonic necrosis
 - l. Signs of chronic pancreatitis – pancreatic calcification
5. Unrelated intraabdominal or intrathoracic findings

Describe findings _____